

Sedentary work and risk of venous thromboembolism¹

by Camilla Ditlev Lindhardt Johannesen, MSc,² Esben Meulengracht Flachs, PhD, Niels E Ebbenhøj, DMSc, Jacob Louis Marott, MSc, Gorm Boje Jensen, DMSc, Børge G Nordestgaard, DMSc, 4 Peter Schnohr, DMSc, Jens Peter E Bonde, PhD

1. *Supplementary tables S1 and 2*

2. *Correspondence to: Camilla Ditlev Lindhardt Johannesen, Department of Occupational and Environmental Medicine, Frederiksberg and Bispebjerg Hospital, Bispebjerg Bakke 23, DK-2400 Copenhagen NV, Denmark. [E-mail: cjoh0108@regionh.dk]*

Supplementary table S1. Self-Registered Baseline Characteristics for Participants in the Copenhagen City Heart Study and the Copenhagen General Population Study, Denmark, 1977-2015.

Characteristic ^a	Self-reported occupational sitting during a workday					
	≥ 6.5 hours		3.5-6.5 hours		≤ 3.5 hours	
	N	%	N	%	N	%
Smoking status						
Current smoker	1,574	17	7,587	22	11,007	32
Former smoker	3,049	34	12,151	34	11,160	32
Never smoked	4,090	45	14,283	40	11,334	33
Missing	377	4	1,371	4	953	3
Alcohol use						
Daily intake	1,881	21	7,793	22	7,709	22
Weekly intake	3,897	43	14,651	41	11,903	35
Monthly intake	2,498	27	9,440	27	9,663	28
Never	786	9	3,349	10	4,980	14
Missing	28	0	159	0	199	1
Activity level, leisure time						

Primarily inactive	678	8	2,679	8	3,078	9
Lightly active	3,733	41	14,797	42	15,209	44
Moderate active	4,013	44	15,326	43	13,878	40
Highly active	642	7	2,472	7	2,068	6
Missing	24	0	118	0	221	1
Education						
Elementary (1-9 years)	855	9	5,199	15	11,325	33
High school (10-12 years)	5,514	61	21,453	61	17,053	50
Academic (>12 years)	2,705	30	8,665	24	5,983	17
Missing	16	0	75	0	93	0
Use of hormones ^b	935	19	3,766	20	3,991	19
Missing	44	1	222	1	300	1
Menopause ^c	2,403	48	9,623	50	11,015	54
Missing	21	0	100	1	116	1

Acronyms: N, number.

a) Only variables included in the final analyses are shown.

b) Hormones comprise oral contraception and hormone replacement therapy among women.

c) Among women only.

Supplementary table S2. Baseline BMI and Blood Samples for Participants in the Copenhagen City Heart Study and the Copenhagen General Population Study, Denmark, 1977-2015.

Characteristic ^a	Self-reported occupational sitting during a workday					
	≥ 6.5 hours		3.5-6.5 hours		≤ 3.5 hours	
	N	%	N	%	N	%
BMI ^b						
< 23.8 kg /m ²	3,290	36	12,829	36	12,234	36
23.8-27.1 kg /m ²	3,116	34	11,951	34	10,633	31
> 27.1 kg /m ²	2,665	30	10,540	30	11,489	33
Missing	19	0	72	0	98	0
Total cholesterol ^b						
< 5.1 mmol/L	3,399	37	12,699	36	11,799	34
5.1-6.0 mmol/L	2,979	33	11,545	33	11,090	32
> 6.0 mmol/L	2,632	29	10,805	30	11,248	33
Missing	80	1	343	1	317	1
CF II+VII+X ^b						
< 87 %	3,173	35	10,937	31	8,304	24
87-105 %	2,790	31	10,212	29	8,833	26
> 105 %	2,406	26	9,234	26	9,879	29
Missing	721	8	5,009	14	7,438	21
Thrombocytes ^b						
< 246 /L	2,724	30	9,759	28	8,032	23
246-299 /L	2,882	32	10,556	30	9,304	27

> 299 /L	2,876	31	10,442	29	10,087	29
Missing	608	7	4,635	13	7,031	21
Fibrinogen ^b						
< 9.6 µmol /L	3,600	40	12,656	36	10,585	31
9.6-11.5 µmol /L	2,773	30	10,242	29	9,110	26
> 11.5 µmol /L	2,216	24	8,504	24	8,796	26
Missing	501	6	3,990	11	5,963	17
CRP ^b						
< 1.1 mg /L	2,991	33	10,475	30	9,466	28
1.1-1.8 mg /L	2,940	32	10,873	31	8,711	25
> 1.8 mg /L	2,332	26	8,680	24	8,662	25
Missing	827	9	5,364	15	7,615	22
Complement C3 ^b						
< 1.0 g /L	3,011	33	10,913	31	9,000	26
1.0-1.2 g /L	2,943	32	10,235	29	9,081	26
> 1.2 g /L	2,499	28	9,475	27	9,241	27
Missing	637	7	4,769	13	7,132	21
Factor V Leiden ^c	389	4	1,633	5	1,620	5
Missing	4,299	47	15,546	44	14,683	43

Acronyms: BMI, body mass index; CF II+VII+X, coagulation factor II+VII+X; N, number.

a) Only variables included in the final analyses are shown.

b) Values are tertile cut off points

c) R506Q hetero- or homozygous