

Original article. Scand J Work Environ Health – online first

Occupational exposure to organic solvents and risk of male breast cancer: a European multicenter case–control study¹

by Nasser Laouali, MSc, Corinne Pilorget, PhD, Diane Cyr, MSc, Monica Neri, PhD, Linda Kaerlev, PhD, Svend Sabroe, PhD, Giuseppe Gorini, PhD, Lorenzo Richiardi, PhD, Maria Morales-Suàrez-Varela, PhD, Agustin Llopis Gonzalez, PhD, Wolfgang Ahrens, PhD, Karl-Heinz Jöckel, PhD, Noemia Afonso, MD, PhD, Mikael Eriksson, PhD, Franco Merletti, PhD, Jørn Olsen, PhD, Elsebeth Lynge, PhD, Pascal Guénel, MD, PhD²

1 *Supplementary table S1*

2 *Correspondence to: Pascal Guénel, MD, PhD, CESP Inserm U1018, 16 avenue Paul Vaillant-Couturier, 94807 Villejuif Cedex, France. [E-mail: pascal.guenel@inserm.fr]*

Table S1: Odds ratios for male breast cancer for exposure to organic solvents after excluding the hospital-based cancer controls									
Solvent	CES (ppm.years)	Cases (n=104)	Controls (n=1395)	Model A		Model B		Model C	
				OR	95% CI	OR	95% CI	OR	95% CI
Benzene	not exposed	82	1251	1.0	reference	1.0	reference	Not selected	
	> 0 < 0.87	13	73	2.3	1.1-5.0	3.1	0.9-10.5		
	≥0.87	9	71	2.3	1.0-5.3	3.4	0.7-16.2		
Trichloroethylene	not exposed	58	926	1.0	reference	1.0	reference	1.0	reference
	>0 <23.90	17	250	1.8	0.9-3.6	1.6	0.8-3.5	1.7	0.9-3.4
	≥23.90	29	219	2.7	1.5-4.8	2.6	1.3-5.4	2.3	1.2-4.3

Abbreviations: CES, cumulative exposure score; OR, odds ratio; CI, confidence interval

Model A: adjusted for age, country, education, BMI and alcohol drinking

Model B: adjusted for variables in Model A and all solvents.

Model C: adjusted for variables in Model B and regression model performed respectively with stepwise backward and stepwise forward variable selection.