

Scand J Work Environ Health 2002;28(2):52-70 Issue date: 2002

Effects of environmental tobacco smoke on the respiratory health of adults by Jaakkola MS, Jaakkola JJK

Affiliation: Finnish Institute of Occupational Health, Topeliuksenkatu 41 a A, FIN-00250 Helsinki, Finland. Maritta.Jaakkola@ttl.fi]

Refers to the following text of the Journal: 2002;28 suppl 2:71-83

The following articles refer to this text: 2002;28 suppl 2:3-6; 2002;28 suppl 2:71-83

Key terms: adult; asthma; chronic obstructive pulmonary disease; environmental tobacco smoke; lung function; respiratory health; respiratory infection; respiratory symptom; tobacco smoke pollution

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/12058803



This work is licensed under a Creative Commons Attribution 4.0 International License.

Effects of environmental tobacco smoke on the respiratory health of adults

by Maritta S Jaakkola, MD,¹ Jouni JK Jaakkola, MD^{2,3}

Jaakkola MS, Jaakkola JJK. Effects of environmental tobacco smoke on the respiratory health of adults. *Scand J Work Environ Health* 2002;28 suppl 2:52–70.

In this paper, current knowledge on the respiratory effects of environmental tobacco smoke among adults is synthesized, and the biological basis and methodological issues are discussed. The Medline database was searched from 1966 through October 2000. All of the related respiratory effects have been linked to both home and workplace exposures. Some evidence of a dose-response relation has also been detected for all respiratory conditions. The strongest evidence of a causal relation exists for chronic respiratory symptoms. Harmful effects on lung function have also been detected, mainly in countries and occupations with high exposure levels. There is limited evidence indicating an increased risk of its causing asthma and chronic obstructive pulmonary disease, and also for poor control of established asthma. More longitudinal studies with careful assessment of exposure are needed for better risk estimates. Despite these challenges for the future, the combination of toxicologic evidence, abundant evidence on respiratory effects among children, and the studies reviewed in this paper point to an urgent need for measures to prevent exposure to environmental tobacco smoke among adults.

Key terms asthma, chronic obstructive pulmonary disease, lung function, respiratory infections, respiratory symptoms, tobacco smoke pollution.

The first report of the consequences of exposure to environmental tobacco smoke at the workplace in relation to the lung function of adults was reported in 1980 (1). Since then, the evidence has accumulated gradually for adverse noncarcinogenic respiratory effects of exposure to environmental tobacco smoke in adulthood. The aims of this review are to synthesize current knowledge on the respiratory effects of environmental tobacco smoke among adults and to discuss the related biological basis and methodological issues. Another article in this journal synthesizes the evidence on the respiratory effects of environmental tobacco smoke among children (2). A Medline database search was carried out from 1966 through October 2000 with the Mesh-terms "Tobacco smoke pollution and exp. Respiratory track diseases". Additional material was collected from the reference lists of articles and from personal knowledge of current research.

Earlier reviews suggested that environmental tobacco smoke makes an important contribution to the total burden of environmental insults on the respiratory system, but they considered the evidence on the causality of the relations as insufficient, except for lung cancer (3-8). More recent reviews, including additional studies, have concluded that there is strong evidence of causal relations between exposure to environmental tobacco smoke and respiratory diseases in adults, especially in the case of respiratory symptoms (9, 10). Assessing exposure to environmental tobacco smoke during adulthood is more complex than such assessment in childhood, since adults are exposed from multiple sources in addition to the home environment (eg, the workplace and public places). The exposure profile may also vary considerably during different periods of adult life (11, 12). For many diseases it is important to consider the role of earlier exposures in addition to current exposure. These difficulties may have contributed to the delay in the interest in studying noncancer respiratory effects among adults.

¹ The Finnish Institute of Occupational Health, Helsinki, Finland.

² Environmental Health Program, The Nordic School of Public Health, Göteborg, Sweden.

³ Environmental Epidemiology Unit, Department of Public Health, University of Helsinki, Finland.

Reprint requests to: Dr Maritta S Jaakkola, Finnish Institute of Occupational Health, Topeliuksenkatu 41 a A, FIN-00250 Helsinki, Finland. [E-mail: Maritta.Jaakkola@ttl.fi]

Biological basis for nonmalignant respiratory effects of environmental tobacco smoke

Active smoking is a well-established cause of chronic respiratory symptoms, lung function decline, and chronic obstructive pulmonary disease (COPD) (13). Its role in determining susceptibility to respiratory infections, such as pneumonia, is also strongly supported (14). The chemical similarities between mainstream and sidestream smoke support the biological plausibility of the role of environmental tobacco smoke in causing respiratory diseases. The noncarcinogenic mechanisms of exposure to environmental tobacco smoke have been studied less than the carcinogenic effects. In this section, we discuss what has been shown or speculated to be potential mechanisms leading to nonmalignant respiratory diseases in subjects exposed to environmental tobacco smoke.

Microbiological organisms are the direct causes of infectious diseases, but exposure to environmental tobacco smoke seems to be an important determinant of susceptibility to such organisms. This susceptibility has been demonstrated very clearly for children, whereas infections have been studied less among adults (2, 9, 15). The few adult studies have linked exposure to environmental tobacco smoke to severe infections, such as pneumonia and meningitis. There may be several mechanisms underlying an increased susceptibility to infections. For example, tobacco smoke impairs host defense mechanisms by weakening immunologic responses and mucociliary clearance (13, 16, 17), and it has been shown to enhance bacterial adherence and to disrupt respiratory epithelium (18, 19).

Tobacco smoke is known to contain several irritative compounds, such as ammonia, sulfur dioxide, nitrogen oxides, acrolein, and formaldehyde (5, 20). It is biologically plausible that exposure to these substances may cause inflammatory or irritative reactions in the airways and therefore lead to respiratory symptoms and lung function impairment. Tobacco smoke has been demonstrated to induce inflammatory reactions in the airways and lung parenchyma of active smokers (21-23), and the biological effects of environmental tobacco smoke are likely to be similar to those of mainstream smoke. In addition, the impairment of host defense mechanisms and the mucociliary clearance of airways increase vulnerability to infections, and repeated respiratory infections may then predispose a person to the development of COPD (13, 17). It is also likely that lung function deficit in newborns and children attributable to exposure to environmental tobacco smoke increases the vulnerability of the lungs to additional insults later in life (2, 9).

There are several biologically plausible mechanisms through which environmental tobacco smoke could

cause and exacerbate asthma. It has been shown to increase the susceptibility to respiratory infections in childhood, and such infections have been associated with both early-onset asthma and exacerbations of underlying disease (24, 25). A more direct mechanism is also plausible, since tobacco smoke contains several irritant substances that could cause an inflammatory response in the airways, as has been shown in the case of irritant-induced occupational asthma (26). The effect of environmental tobacco smoke on asthma seems to occur via irritative rather than allergic mechanisms, and skin test positivity to tobacco is rare (6, 7, 27). Tobacco smoke may also increase epithelial permeability to environmental allergens and thus enhance allergic reactions to other inhalable allergens (28). In line with this hypothesized mechanism, passive smoking has been observed to increase total immunoglobulin E (IgE) among children and adults (29-31).

Effects of environmental tobacco smoke on the respiratory health of adults

Respiratory symptoms

Acute symptoms. The role of environmental tobacco smoke in producing acute irritative symptoms of the eyes, nose, throat, and lower airways is well established (3, 6, 7). Some experimental studies have shown significant increases in objective measurements related to symptoms, such as eye blink rate and nasal airway resistance, in response to exposure to environmental tobacco smoke (32, 33). Atopic subjects appear to be more sensitive to environmental tobacco smoke than nonatopics (7, 33, 34).

Chronic respiratory symptoms. The role of exposure to environmental tobacco smoke in the development of chronic respiratory symptoms in adults has been addressed in 12 cross-sectional (35–46) and 2 longitudinal studies (47, 48). Tables 1 and 2 summarize these studies. In addition, a study on the effects of exposure cessation on respiratory symptoms in bartenders has been reported (49). Most of the studies assessed household exposure, three studies assessed workplace exposure, and four studies assessed exposure in both of these environments. Two additional studies assessed the effects on a composite symptom score (50) and symptom severity (51).

The results of the studies are somewhat inconsistent. This inconsistency may be explained partly by methodological differences, for example, the exposure assessment approach used. In summary, the studies with a stronger power, either due to a large sample size or

Table 1. Cross-sectional	studies on env	rironmental tobacc	o smoke and	chronic	respiratory	symptoms	among adults.	(95% CI=9	95%
confidence interval, OR =	odds ratio, PR =	= prevalence ratio,	RR = risk ratio))					

Study	Study population, age	Exposure assessment	Measure Exposure of effect category compared to no exposure		Results
Lebowitz & Burrows 1976, United States (35)	1258 men and women, never smokers, ≥15 years	Household member's report of current smoking	PR	Current exposure	PR 0.66 $^{\rm a},$ 1.0 $^{\rm a},$ & 1.02 $^{\rm a}$ for cough, phlegm & wheezing, respectively
Schilling et al 1977, United States (36)	178 men, 212 women, nonsmokers	Spouse's report of current smoking	PR	Spouse smoking	Not significant for cough, phlegm or wheezing
Comstock et al 1981, United States (37)	426 men, 113 women never smokers, \geq 20 years	Household member's report of smoking 2 years earlier	RR	Smoking co- habitees	Men: RR ^b 0.96, 0.90, 1.04, & 1.08 for cough, phlegm, wheezing & dyspnea, respectively Women: RR ^b 0.17, 0.72, 1.45 & 1.79 for cough, phlegm, wheezing & dyspnea, respectively
Schenker et al 1982, United States (38)	4156 women, nonsmokers, 17–74 years	Self-report of spouse's current smoking	OR	Spouse smoking	Not significant for cough, phlegm or wheezing; OR 1.56 ° for dyspnea
Hole et al 1989, United Kingdom (39)	671 men, 1784 women, never smokers, 45–64 years	Household member's report of ever smoking	OR (95% CI)	Ever ex- posed at home	OR 1.19 (95% Cl 0.85–1.67) & 1.09 (95% Cl 0.82–1.45) for phlegm & dyspnea, respectively
Kauffmann et al 1989, United States and France (40)	726 American women, 25–69 years; 2298 French women, 25–59 years, never smokers	Spouse's report of current smoking	OR (95% CI)	Spouse smoking	American: OR 1.14 (95% CI 0.62–2.09), 1.65 (95% CI 0.72–3.78), 1.35 (95% CI 0.97–1.87) & 1.35 (95% CI 0.68–2.61) for cough, phlegm, wheezing & dyspnea, respectively French: OR 1.35 (95% CI 0.78–2.36), 0.77 (95% CI 0.29–2.03), 1.03 (95% CI 0.77–1.38) & 1.17 (95% CI 0.87–1.57) for cough, phlegm, wheezing & dyspnea, respectively
White et al 1991, United States (41)	80 men and women, never smokers, 38–65 years; subjects with home exposure excluded	Self-report of workplace exposure, verified with observations and carbon monoxide measurements in the work- place	PR - 9	Exposed at work for ≥1 years	PR 2.8 ^d , 3.4 ^d & 4.5 ^d for cough, phlegm, & dyspnea, respectively
Pope & Xu 1993, China (42)	973 women, never smokers, 20–40 years	Self-report of household members' current smoking	OR (95% CI)	1 smoker, ≥2 smokers, ≥2 smokers + coal heating	1 smoker: OR 1.02 (95% Cl 0.60–1.75), 1.43 (95% Cl 0.85–2.40), 0.93 (95% Cl 0.50–1.75) & 1.17 (95% Cl 0.61–2.25) for cough, phlegm, wheezing & dyspnea, respectively ≥2 smokers: OR 1.87 (95% Cl 0.71–4.88), 2.07 (95% Cl 0.85–5.01), 1.00 (95% Cl 0.27–3.71) & 1.46 (95% Cl 0.39–5.52) for cough, phlegm, wheezing & dyspnea, respectively ≥2 smokers + coal heating: OR 3.07 (95% Cl 1.23–7.65), 3.64 (95% Cl 1.56–8.52), 1.07 (95% Cl 0.29–4.00) & 3.55 (95% Cl 1.20–10.5) for cough, phlegm, wheezing & dyspnea, respec- tively
Ng et al 1993, Singapore (43)	1282 women, never smokers, 20–74 years	Self-report of household members' ever smoking	OR (95% CI)	≥1 light° smokers, ≥1 heavy° smokers	≥1 light smokers: OR 2.84 (95% CI 1.29–6.24), 1.50 (95% CI 0.83–2.71), 0.95 (95% CI 0.38– 2.39) & 1.23 (95% CI 0.87–1.74) for cough, phlegm, wheezing & dyspnea, respectively ≥1 heavy smokers: OR 3.79 (95% CI 1.76–8.14), 1.36 (95% CI 0.72–2.57), 2.69 (95% CI 1.23– 5.88) & 1.83 (95% CI 1.30–2.58) for cough, phlegm, wheezing & dyspnea, respectively
Leuenberger et al 1994, Switzerland (44)	1674 men, 2523 women, never smokers, 18–60 years	Self-report of exposure at home and work during the past 12 months	OR ^f (95% CI)	Any expo- sure, workplace exposure	Any exposure: OR 1.69 (95% CI 1.23– 2.31), 1.99 (95% CI 1.41–2.82) & 1.44 (95% CI 1.18–1.75) for phlegm, wheezing & dyspnea, respectively Workplace exposure: OR 1.67 (95% CI 1.23– 2.28), 2.05 (95% CI 1.42–2.96) & 1.62 (95% CI 1.29–2.03) for phlegm, wheezing & dyspnea, respectively

Table 1. Continued.

Study	Study population, age	Exposure assessment	Measure of effect	Exposure category compared to no exposure	Results
Piitulainen et al 1998, Sweden (45)	95 men, 110 women, severe α_1 -antitrypsin deficiency, never smokers, ≥20 years	Self-report of exposure ever at home and work (days/week, years of exposure)	OR (95% CI)	Exposure at home and work for ≥10 years	OR 1.6 ⁹ (95% Cl 1.3–2.4), 1.1 (95% Cl 0.7–1.6) & 1.2 (95% Cl 0.8–1.8) for phlegm, wheezing & dyspnea respectively
Lam et al 2000, Hong Kong (46)	4468 men, 728 women, never smokers, 18–58 years, police officers	Self-report of current exposure at home and work (number of smokers, number of cigarettes smoked by co-workers, hours of daily exposure)	OR (95% CI)	Exposure at home or work	Men (home): OR 1.03 (95% CI 0.68–1.56), 1.29 (95% CI 0.91–1.82) & 1.41 (95% CI 1.08–1.85) for cough, phlegm, & wheezing, respectively Men (work): OR 2.61 (95% CI 1.50–4.55), 2.05 (95% CI 1.35–3.11) & 1.76 (95% CI 1.26– 2.45) for cough, phlegm, & wheezing, respec- tively Women (home): OR 0.52 (95% CI 0.18–1.48), 0.62 (95% CI 0.25–1.51) & 1.07 (95% CI 0.57–2.03) for cough, phlegm & wheezing, respectively Women (work): OR 1.00 (95% CI 0.29–3.47), 2.57 (95% CI 0.67–9.90) & 0.55 (95% CI 0.26–1.19) for cough, phlegm & wheezing, respectively

^a Statistically nonsignificant.

^b All were statistically nonsignificant.

^c Statistically significant at P = 0.001.

^d Statistically significant at P<0.001.

Light smoker – household member who smokes or has smoked <20 cigarettes/ day, heavy smoker = household member who smokes or has smoked >20 cigarettes/day.

^t The OR values for any exposure have been taken from analyses excluding subjects with end-expiratory carbon monoxide levels of ≥7 ppm in order to reduce the misclassification of exposure; the OR values given for phlegm are actually for chronic cough and phlegm combined.

⁹ Chronic bronchitis defined as daily cough with phlegm for at least 3 months/year.

Study	Study population, age, length of follow- up	Exposure assessment	Measure of effect	Exposure compared with no exposure	Results
Schwartz & Zeger 1990, United States (47)	100 women, 18% smokers, age not given, 3 years	Self-report of presence of a smoking roommate	OR (95% CI) (controlled for persona smoking)	Smoking roommate I	Not significant for cough, OR 1.41 (95% Cl 1.08–1.85) for phlegm
Jaakkola et al 1996, Canada (48)	60 men, 57 women, never smokers, 15–40 years at the baseline, 8 years (1980/1981– 1988/1989)	Self-report of household members' smoking and exposure at work during the follow-up	OR (95% CI) per 10 cigarettes/ day	Total ex- posure (home+ work)	OR 1.55 (95% Cl 0.61–3.90), 0.69 (95% Cl 0.21–2.26), 1.15 (95% Cl 0.64–2.06) & 2.37 (95% Cl 1.25–4.51) for cough, phlegm, wheezing & dyspnea, respectively

 Table 2. Longitudinal studies on environmental tobacco smoking and chronic respiratory symptoms among adults. (95% CI = 95% confidence interval, OR = odds ratio)

longitudinal follow-up, have shown a significantly increased risk of chronic respiratory symptoms in relation to exposure to environmental tobacco smoke among adults (38, 43, 44, 46–48). In addition, some other crosssectional studies have shown a tendency towards an increased risk of symptoms related to exposure to environmental tobacco smoke, although the effects did not reach statistical significance (39, 40) or they were significant only in a sensitive subgroup (42). The excess risk of symptoms related to exposure to environmental tobacco smoke has varied between 40% and over 300%. Five studies found significant relations between workplace exposure and the occurrence of chronic respiratory symptoms when workplace exposure was assessed separately or as part of total exposure (41, 44–46, 48).

A dose-response relation was demonstrated between exposure to environmental tobacco smoke and the risk of symptoms in five studies (42–44, 46, 48). In a Swiss study (44) of nonsmoking men and women, the following symptoms showed a statistically significant relation to the quantitative estimate of exposure to environmental tobacco smoke: bronchitis symptoms, wheezing and dyspnea with the hours of daily exposure, bronchitis symptoms and dyspnea with the number of smokers to whom the subject was exposed, and wheezing with the years of exposure. In an 8-year follow-up study of nonsmoking young adults in Canada, the risk of developing dyspnea increased significantly with increasing total exposure to environmental tobacco smoke, the odds ratio (OR) being 2.37 for average exposure to 10 cigarettes/day (48). An increasing trend with increasing total exposure to environmental tobacco smoke was detected also for wheezing and cough, but the relations did not reach statistical significance.

In a study from San Francisco, California, 53 bartenders responded to an interview and performed spirometry about 1 month before and after a bill prohibiting smoking in bars and taverns came into effect (49). Workplace exposure to environmental tobacco smoke was reported to decline from a median of 28 to 2 hours per week. This decline was accompanied by a significant reduction in the prevalence of lower respiratory symptoms (wheezing, dyspnea, cough, and phlegm production) from 74% to 32% and in the prevalence of upper airway irritation symptoms (eye, nose and throat symptoms) from 77% to 19%.

A study from China showed synergistic effects of passive smoking and coal heating on the occurrence of respiratory symptoms (42). A study of a potentially susceptible group consisting of 205 never-smoking persons with severe α_1 -antitrypsin deficiency (with PiZZ genotype) from Sweden reported effects on chronic bronchitis, but not on wheezing, dyspnea, or lung function (45).

In a cross-sectional study of 2992 adults from the United Kingdom, an increasing trend for symptom severity was found across the smoking categories, the median score being 2.8 for never smokers, 4.2 for passive smokers, 3.7 for past smokers, and 5.6 for current smokers (51). The adjusted odds ratio was 1.4 [95% confidence interval (95% CI) 1.0-1.8] for severe symptoms in relation to current household exposure to environmental tobacco smoke. Another cross-sectional study of 3405 nonsmoking women from Hong-Kong reported the effects of household smoking on a composite symptom outcome, including sore throat, morning and evening cough, phlegm production during day or night, and phlegm for 3 months (50). The risk of occurrence of any symptom increased with the number of smokers at home (OR 1.16, 95% CI 0.83-1.62, for exposure to one smoker and 1.80, 95% CI 1.15-2.83, for exposure to two or more smokers).

In summary, abundant evidence from different countries indicates that exposure to environmental tobacco smoke causes chronic respiratory symptoms in adults. A dose-response relation has been shown in several studies, and the effects are biologically plausible. Exposure misclassifications due to active smoking or confounding are not likely explanations for the observed relations. Several of the studies took into account potential confounders, such as age, gender, socioeconomic status, atopy, occupational exposures, and other indoor sources of pollutants, either in the study design or in the statistical analyses. The large Swiss study excluded subjects with end-expiratory carbon monoxide over 7 parts per million (ppm) in order to exclude active smokers, but the exclusion did not essentially change the relations between exposure and chronic respiratory symptoms (44). However, more longitudinal studies are needed to throw light on temporal relations and to exclude recall bias as a potential explanation.

Lung function

Studies on the effects of environmental tobacco smoke on the lung function of adults have focused mainly on ventilatory lung function. Altogether 18 cross-sectional studies (1, 36, 37, 39, 40, 43, 52–63), 1 case-referent study (64), and 4 longitudinal studies (55, 63, 65, 66) have addressed these relations. These studies are summarized in tables 3 and 4. In addition, a study of the reduction of exposure among Californian bartenders (table 4) and another study among Canadian bartenders on workshift changes have reported lung function effects (49, 67). Three studies have addressed the effects of environmental tobacco smoke on airway hyperresponsiveness, measured with peak expiratory flow (PEF) variability or methacholine challenge (61, 62, 68).

The results of the studies have been somewhat inconsistent. Among the cross-sectional and case-referent studies, 10 studies detected significant adverse effects on forced expiratory volume in 1 second (FEV₁), indices of small airway function, or both of these parameters (1, 39, 43, 52, 55–59, 63), while 6 did not find any significant associations (36, 37, 40, 53, 60, 64). The FEV₁ deficit related to the exposure to environmental tobacco smoke was estimated to be from -50 ml in the Singapore study (43) to -257 ml in the China study (when adjusted for confounders) (59). In a meta-analysis including nine cross-sectional studies, the effect of environmental tobacco smoke on FEV₁ was -2.7% (95%) CI -4.1% - -1.2%) (63). All the cross-sectional studies but one assessed exposure from household members' smoking, while workplace exposure was assessed in eight studies (1, 53, 54, 57-59, 61, 62). Significant lung function impairment related to workplace exposure was detected in five studies (1, 54, 57-59). Four cross-sectional studies showed a significant dose-response relation between the amount of household smoking and the reduction in lung function, either FEV₁ or forced expiratory flow rate at 25-75% of forced vital capacity

Table 3. Cross-sectional and case-referent studies on environmental tobacco smoke exposure and lung function among adults. (95% CI = 95% confidence interval, cig = cigarettes, DL_{co} = carbon monoxide diffusing capacity of the lung, FEF_{25-75} = forced expiratory flow rate at 25–75% of the FVC, FEF_{75-85} = mean forced expiratory flow during 75% to 85% of the FVC, FEV_1 = forced expiratory volume in 1 second, FVC = forced vital capacity, MEF_{25} = maximal expiratory flow at 75% of the FVC, MEF_{50} = maximal expiratory flow at 50% of the FVC, MEF_{75} = maximal expiratory flow at 25% of the FVC, MMEF = mean forced expiratory flow during the middle half of the FVC, $PaCO_2$ = arterial blood carbon monoxide tension, PaO_2 = arterial blood oxygen tension, PEF = peak expiratory flow, TET = total expiratory time, VC = vital capacity)

Study	Study population, age	Exposure assessment	Lung function outcomes	Results
Schilling et al 1977, United States (36)	178 men, 212 women, nonsmokers, age not given	Spouse's report of current smoking	Level of FVC (I), FEV_1 (I), PEF (I/s), MEF_{50} (I/s), MEF_{25} (I/s)	No significant relation between exposure and lung function
White & Froeb 1980, United States (1)	400 men, 400 women, nonsmokers, mean age 46-49 years, no home exposure	Self-report of expo- sure at work for ≥20 years	Level of FVC (I), FEV ₁ (I), FEF ₂₅₋₇₅ (I/s), FEF ₇₅₋₈₅ (I/s)	Men (unexposed): FEV ₁ 3.72, FEF ₂₅₋₇₅ 3.78, FEF ₇₅₋₈₅ 1.22 Men (exposed): FEV ₁ 3.54, FEF ₂₅₋₇₅ 3.30 °, FEF ₇₅₋₈₅ 0.97 ° Women (unexposed): FEV ₁ 2.63, FEF ₂₅₋₇₅ 3.17, FEF ₇₅₋₈₅ 1.03 Women (exposed): FEV ₁ 2.47, FEF ₂₅₋₇₅ 2.72 °, FEF ₇₅₋₈₅ 0.78 °
Comstock et al 1981, United States (37)	369 men, never smokers, ≥20 years	Household member's report of smoking 2 years earlier	Prevalence ratio of FEV ₁ <80% predicted, FEV ₁ /FVC<70%	Prevalence ratio: FEV, <80% 1.42, FEV,/FVC<70% 1.19
Jones et al 1983, United States ^b (64)	205 women, non- smokers, 20–39 years	Self-report of house- hold members' smok- ing 1–2 years earlier	Odds ratio of exposure in lowest versus highest quartile of FEV ₁ % predicted	Odds ratio 0.76 (P = 0.34)
Kauffmann et al 1989 (40), 1983 (52), France and United States	France: 970 men, 2306 women, never smokers, 25–59 years; United States: 726 women, never smokers 25–69 years	Spouse's report of current smoking ,	Level of FVC (I), FEV1 (I), FEF25-75 (I/s)	French men \geq 40 years (unexposed): FVC 4.08, FEV ₁ 3.31, FEF ₂₅₋₇₅ 3.58 French men \geq 40 years (exposed °): FVC 4.11, FEV ₁ 3.19, FEF ₂₅₋₇₅ 3.02 ° French wormen \geq 40 years (unexposed): FVC 2.98, FEV ₁ 2.43, FEF ₂₅₋₇₅ 2.74 French wormen \geq 40 years (exposed °): FVC 2.89 °, FEV ₁ 2.34 °, FEF ₂₅₋₇₅ 2.57 ° American women: no significant relation between exposure and lung function
Kentner et al 1984, Germany (53)	393 men, 198 women, never smokers, median age 36-42 years	Self-report of current smoking by household members and co-worke	Level of FVC, FEF ₂₅₋₇₅ , FEF ₇₅₋₈₅ as % predicted ers	Men (unexposed): FVC 99, FEF $_{25-75}$ 99 Men (exposed): FVC 99, FEF $_{25-75}$ 101 Women (unexposed): FVC 100, FEF $_{25-75}$ 98 Women (exposed): FVC 99, FEF $_{25-75}$ 95
Salem et al 1984, Egypt (54)	517 men, 79 women, never smokers, age not given	Self-report of current and past smoking by household members or co-workers or both	Level of TET (s), FEV ₁ (l), PEF (l/min), PaO ₂ (mm Hg), PaCO ₂ (mm Hg)	Unexposed: TET 2.53, FEV ₁ 2.97, PEF 488, PaO ₂ 93.0, PaCO ₂ 37.0 Exposed ⁴ :TET 3.31 ^a , FEV ₁ 2.80, PEF 473, PaO ₂ 91.5, PaCO ₂ 39.0 ^a
Brunekreef et al 1985, The Netherlands (55)	97 women, nonsmokers, 25–60 years	Self-report of household members' smoking since1965	$\begin{array}{l} \mbox{Level of FVC (I), FEV}_1 (I), \\ \mbox{PEF (I/s), MEF}_{75} (I/s), \\ \mbox{MEF}_{50} (I/s), \mbox{MEF}_{25} (I/s), \\ \mbox{MMEF (I/s)} \end{array}$	40- to 60-year-olds (unexposed): FVC 3.45, FEV, 2.82, PEF 8.12, MEF ₇₅ 6.96 40- to 60-year-olds (exposed $^{\circ}$): FVC 3.34, FEV ₁ 2.63, PEF 6.79 a , MEF ₇₅ 5.89 a
Svendsen et al 1987, United States (56)	1245 men, never smokers, 35–57 years	Self-report of spouse's current smoking	Level of FEV_1 (I)	Unexposed: FEV_1 3.59 Exposed: FEV_1 3.49 $^{\rm a}$ for all, FEV_1 3.41 for 1–19 cig/ day, FEV_1 3.55 for ≥ 20 cig/day
Masi et al 1988, Canada (57)	133 men, 160 women, never smokers, 15–35 years	Self-report of lifetime exposure at home and work, cumulative exposure indices	Estimated effect on VC (I), FVC (I), FEV ₁ (I), PEF (I/s), FEF ₂₅₋₇₅ (I/s), DL_{c0} [mV(min \cdot mm)]	$\begin{array}{l} \mbox{Men (home, per person-years): VC 0.004, FEV_1 - 0.002 \\ \mbox{FEF}_{25-75} & -0.020, ^a \mbox{DL}_{c0} & 0.020 \\ \mbox{Men (work, per pack/day-years): VC } & -0.045, ^a \\ \mbox{FEV}_1 & -0.022, \mbox{FEF}_{25-75} & 0.024, \mbox{DL}_{c0} & -0.202 \\ \mbox{Women (home, per person-years): VC } & 0.001, \\ \mbox{FEV}_1 & -0.001, \mbox{FE}_{25-75} & -0.003, \mbox{DL}_{c0} & -0.006, \\ \mbox{FEV}_1 & -0.011, \mbox{FE}_{25-75} & -0.023, \mbox{DL}_{c0} & -0.258 \ ^a \end{array}$
Hole et al 1989, United Kingdom (39)	671 men, 1784 women, never smokers, 45–64 years	Household member's report of ever smoking	Level of FEV_1 (I)	Unexposed: FEV₁ 2.31 for men + women, FEV₁ 1.88 for women Exposed: FEV₁ 2.23 ^a for men + women, FEV₁ 1.89 for women (<15 cig/day), FEV₁ 1.83 ^a women (≥15 cig/day)

Table 3. Continued.

Study	Study population, age	Exposure assessment	Lung function outcomes	Results
Masjedi et al 1990, Iran (58)	167 men, 108 women, never smokers, 18–65 years	Self-report of current smoking by household members and co- workers	Level of FVC, FEV ₁ , PEF, FEF ₂₅₋₇₅ % predicted	Men (unexposed): FVC 102.1, FEV ₁ 102.6, PEF 100.9, FEF ₂₅₋₇₅ 104.3 Men (exposed'): FVC 97.5, ^a FEV ₁ 96.9, ^a PEF 98.8 FEF ₂₅₋₇₅ 94.4 ^a Women (unexposed): FVC 98.9, FEV ₁ 99.3, PEF 100.6, FEF ₂₅₋₇₅ 98.4 Women (exposed'): FVC 101.2, FEV ₁ 100.9 PEF 99.4, FEF ₂₅₋₇₅ 99.8
Ng et al 1993, Singapore (43)	1008 women, never smokers, 20–74 years	Self-report of house- hold members' ever smoking	Level of FEV ₁ (I) measured with a portable spirometer	Unexposed: FEV ₁ 1.83 Light ^a exposure: FEV ₁ 1.78 ^a Heavy ^g exposure: FEV ₁ 1.76 ^a
Xu & Li 1995, China (59)	131 men, 371 women, never smokers, 40–69 years	Self-report of current smoking by household members and co- workers	Level of FVC (I), FEV_1 (I)	Men (unexposed): FVC 3.69, FEV ₁ 2.96 Men (exposed ⁿ): FVC 3.21 ^a , FEV ₁ 2.61 ^a Women (unexposed): FVC 2.46, FEV ₁ 1.98 Women (exposed ⁿ): FVC 2.44, FEV ₁ 1.98
Frette et al 1996, United States (60)	176 men, 415 women, never smokers, 51–95 years	Self-report of current or past smoking by household members	Level of FVC (I), FEV1 (I), FEF25-75 (I/s)	$\begin{array}{l} \mbox{Men (unexposed): FVC 4.10, FEV_1 3.14, } \\ \mbox{FEF}_{25-75} 2.80 \\ \mbox{Men (exposed '): FVC 3.90, FEV_1 3.00, FEF}_{25-75} 2.66 \\ \mbox{Women (unexposed): FVC 2.71, FEV_1 2.13, } \\ \mbox{FEF}_{25-75} 1.99 \\ \mbox{Women (exposed '): FVC 2.72, FEV_1 2.12, } \\ \mbox{FEF}_{25-75} 2.00 \end{array}$
Casale & Pasqualetti 1997, Italy (61)	30 healthy never smokers, 30 ex- posed nonsmokers, men, mean age 43–44years	Self-report of expo- sure at home or work or both for ≥3 hours/ day, also urinary cotinine	Diurnal PEF (I/s): average level (mesor), amplitude as the absolute value and the % mesor	Unexposed: mesor 5.91, amplitude: absolute 0.37, % mesor 6.26 Exposed: mesor 5.52, amplitude: absolute 0.43, % mesor 7.79
Abbey et al 1998, United States (62)	519 men, 872 women, nonsmokers, 25–80 years	Self-report of expo- sure at home and work (in years)	Estimated effect on FEV ₁ (% predicted), FEV ₁ /VC (%), PEF lability (%)	Men (worked 10 years with smoker): FEV ₁ /VC -0.5, PEF lability 0.4 $^{\rm }$ Women (lived 10 years with smoker): FEV ₁ /VC -0.2, PEF lability 0.3
Carey et al 1999, United Kingdom (63)	683 men, 940 women, nonsmokers, 18–73 years	Self-report of current smoking by household members and past smoking of parents, salivary cotinine	FEV_1 (I) deficit (95% CI) between the top and bottom quintiles of salivary cotinine	Men: FEV, deficit -166 ml (95% Cl -295 – -37) Women: FEV, deficit -68 ml (95% Cl -145–9) Total: FEV, deficit -105 ml (95% Cl -174 – -37)

^a Significantly lower values for the exposed than for the unexposed (P<0.05).

^b Case-referent study.

^c Exposed = living with a smoker of \geq 10 grams/day.

^d Exposed = exposure at home or work or both daily for \geq 4 hours for at least 12 months.

^e Exposed = living in a home where \geq 10 cigarettes had been smoked daily since 1965.

^f Exposed = reported exposure at home or work or both.

⁹ Light exposure = ≥ 1 household members smoke or had smoked < 20 cigarettes per day, heavy exposure = ≥ 1 household members smoke or had

smoked ≥20 cigarettes per day.

^h Exposed = current smokers at home or work or both.

Exposed = has ever lived in a household with a smoker.

Significantly higher PEF variability in exposed than the unexposed (P<0.05).

(FEF₂₅₋₇₅), in at least a subgroup of the population (52, 57, 59, 63). Two additional studies found a significant effect for heavy exposure on FEV₁, but they did not show a gradient across exposure levels (39, 43).

In contrast to the cross-sectional studies, none of the longitudinal studies found significant effects of exposure to environmental tobacco smoke on the change in lung function over time (55, 63, 65, 66). The followup ranged from 2 years (65) to 17 years (55). Jaakkola and her co-workers followed a cohort of 117 young adults aged 15 to 40 years for 8 years (66). Average exposures to environmental tobacco smoke at home and in the workplace were estimated during the study period as cigarettes/day, and cumulative exposure from home and workplace sources before the beginning of the study period was estimated as cigarette-years. No significant effects were detected on the change in FEV₁ or FEF₂₅₋₇₅ over time in relation to exposure from either source. In a subgroup analysis of young subjects (25 years or younger), cumulative home exposure before the study period was statistically significantly related to a decline in FEV₁. However, this effect was too small to be considered physiologically relevant.

Table 4. Longitudinal studies on environmental tobacco smoke exposure and lung function among adults. (95% CI = 95% confidence interval, FEF_{25-75} = mean forced expiratory flow during the middle half of the forced vital capacity, FEV_1 = forced expiratory volume in 1 second, IVC = inspiratory vital capacity, PEF = peak expiratory flow)

Study	Study population, age, length of follow-up	Exposure assessment	Lung function outcomes	Results
Lebowitz 1984, United States (65)	229 adults, age not given, 2 years	Household member's report of current smoking	Mean PEF level during study period based on daily measurements	No significant relation between exposure and PEF
Brunekreef et al 1985, The Nether- lands (55)	57 women, nonsmokers, 40-60 years (at the end of follow-up), 15–17 years (1965/1967–1982)	Self-report of household members' smoking since 1965	Change in IVC (ml/year), FEV ₁ (ml/year)	Unexposed: IVC -10.1, FEV $_{\rm 1}$ -19.3 Exposed *: IVC -11.4, FEV $_{\rm 1}$ -13.9, not significant
Jaakkola et al 1995, Canada (66)	60 men, 57 women, 15-40 years (at the base- line), 8 years (1980/ 1981–1988/1989)	Self-report of average exposure at home and work during the study, lifetime exposure at home and work before the study (cumula- tive exposure)	Change in FEV ₁ (ml/year), FEF ₂₅₋₇₅ [l/(s.year)]	Exposure during study ^b (cigarettes/day): FEV ₁ 0.53 (95% Cl -0.21 – 1.28) Cumulative exposure before study ^b (cigarette-years): FEV ₁ -0.01, (95% Cl -0.03 – 0.007) No significant relations between exposure and FEF ₂₅₋₇₅ ; no evidence of modification by gender, atopy, or wheezing
Eisner et al 1998, United States (49)	38 men, 15 women, mean age 42.5 years	Self-report of workplace exposure before and 1 month after a bill pro- hibiting smoking in bars came into effect in California (in 1998)	Change in FVC (ml), FEV, (ml)	Change in lung function $^{\circ}$: 326 ml (95% Cl 9–565) for FVC & 157 ml (95% Cl 1–303) for FEV, for expo- sure of \geq 7 hours/week; 244 ml (95% Cl 0–489) for FVC & 127 ml (95% Cl -24–277) for FEV, for expo- sure of 1–6 hours/week
Carey et al 1999, United Kingdom (63)	68 men, 940 women, nonsmokers, 18–73 years at baseline, 7 years (1984/1985–1991)	Self-report of smoking by household members at baseline and follow-up, past smoking of parents	Change in FEV ₁ (mI)	Exposure started during follow-up: FEV_1 -43 (95% Cl -114 - 28), Exposure continued during study: FEV_1 25 (95% Cl -20 - 70), Mother smoked in childhood: FEV_1 -25 (95% Cl -58 - 9)

^a Exposed = living in a home where ≥10 cigarettes had been smoked daily since 1965.

^b Exposure during study = average home + work exposures during the study period (from 1980–1981 to 1988–1989); cumulative exposure before study = lifetime home + work exposures before the beginning of the study.

^c Improvement in lung function among those whose exposure ceased compared with the exposure group shown.

Two studies evaluated the effects of a change in exposure to environmental tobacco smoke among bartenders. The study of Californian bartenders showed significant improvement in lung function after workplace exposure was reduced following a bill banning smoking in bars and taverns (table 4) (49). A dose-dependent improvement was noted when those with complete cessation of workplace exposure were compared with those with moderate (1–6 hours weekly) and high $(\geq 7 \text{ hours weekly})$ exposure at the time of the followup. A study from Canada compared workshift changes in lung function between 26 nonsmoking bar workers exposed to environmental tobacco smoke and 14 nonsmoking unexposed servers (67). The bar workers experienced significant decrements in lung function over a workshift (-42 ml in FEV₁, 95% CI -83 - 0.1, and -155 ml/s in FEF₂₅₋₇₅, 95% CI -290 - -21), while servers working in restaurants where smoking was banned did not.

A study from California found a significantly higher variability in PEF among men who had worked for 10 years with a smoker (62). A study from Italy applied a cosinor analysis to evaluate the effects of exposure to environmental tobacco smoke at home and work on diurnal PEF variability (61). Passive smokers had a lower average level (mesor) and higher PEF amplitude than unexposed referents, but the differences were not statistically significant, probably due to the small sample size. In line with these results, a cross-sectional study of random population samples of adults from west and east Germany showed that current home exposure to environmental tobacco smoke increased significantly the risk of bronchial hyperresponsiveness (OR 1.37, 95% CI 1.05–1.78), defined as PD_{20} (cumulative dose of histamine at which FEV_1 declines from the basic value) smaller than or equal to 2.0 mg in methacholine challenge or a positive response to a bronchodilator test (68).

One cross-sectional study also assessed the effect of cumulative exposure to environmental tobacco smoke on the diffusing capacity of the lung (DL_{CO}) (57). Among young women, cumulative exposure at work was significantly related to a decrease in DL_{CO} . It was estimated that a woman who has never smoked but who has worked in an office with heavy exposure to environmental tobacco smoke for 10 years would have a DL_{CO} that is 3 units lower than if she has worked in a smoke-free office.

The inconsistency of the results between different studies may be explained partly by methodological

differences, such as the exposure assessment methods used, selection of the study population, and adjustment for potential confounders. However, these differences do not seem to explain the disparate findings completely. It appears that the effect of environmental tobacco smoke on the lung function of adults is dose-dependent and may not be seen in low exposures, whereas some countries, such as China, seem to have levels high enough to produce clinically relevant adverse effects. In addition, in some occupations with high exposure levels, such as bartending, the cessation of exposure seems to result in significant improvement in lung function. The sample size of most of the longitudinal studies has been relatively small, and therefore the sensitivity of the studies to detect small adverse effects may have been reduced, although this possibility has been partly compensated by a long follow-up period, which increases the power of the study. It is possible that the small effects on lung function detected in several cross-sectional studies in relation to exposure to environmental tobacco smoke are due to a susceptible group experiencing a more pronounced lung function deficit, and thus potential determinants of susceptibility should be studied in the future. Atopy and asthma are among important individual characteristics that may lead to sensitivity to environmental tobacco smoke.

In summary, several cross-sectional studies on environmental tobacco smoke and lung function in adults indicate small but significant reductions in ventilatory function parameters among subjects exposed at home or in the workplace. This effect seems to be dose-dependent, and it has been observed mainly in countries and occupations with high exposure levels. Exposure misclassification or confounding does not explain the observed relations. Most of the studies controlled for gender, age, height, socioeconomic status, or education or some combination of these variables. Other potential confounders that have been adjusted for included atopy (66), housing conditions and other indoor air pollutants (37, 43, 52, 55, 57, 59, 64, 65), occupational exposures (1, 52, 53, 57, 59, 66), upper respiratory infections (49, 62), and outdoor air pollutants (62). There is a need for more longitudinal studies to gain information on the potential effects of exposure to environmental tobacco smoke on the development of lung function during adulthood.

Asthma

There is abundant evidence of the role of environmental tobacco smoke in the development and prognosis of asthma among children (2), but a limited number of studies has addressed these relations among adults. The studies on adult asthma can be divided into etiologic studies and prognostic studies of the severity of established disease.

Induction of asthma. Environmental tobacco smoke has been addressed with respect to the induction of asthma in adulthood in five studies (43, 44, 69–71), summarized in table 5. Preliminary results from a population-based incident case-referent study from Finland have also been reported recently (72) (table 5). Two additional studies included asthma in their definition of obstructive lung disease, and they are reviewed with chronic obstructive pulmonary disease (73, 74). In addition to these studies of diagnosed asthma, studies on environmental tobacco smoke and respiratory symptoms and bronchial hyperresponsiveness, reviewed in the earlier sections, provide information relevant to early stages of asthma development.

All of the five etiologic studies concerning adults (table 5) found an increased risk of asthma in relation to exposure to environmental tobacco smoke, although this effect did not reach statistical significance in two studies (43, 70). All the studies assessed residential exposure to environmental tobacco smoke, and three also assessed workplace exposure. In the cross-sectional and case-referent studies, the risk of asthma increased by 40% to 200% in relation to exposure to environmental tobacco smoke. In two of the studies the risk of asthma increased in relation to workplace exposure, but not significantly in relation to home exposure (69, 70). A longitudinal study of Seventh-Day Adventists in the United States reported an odds ratio of 1.45 (95% CI 1.21-1.80) for asthma in relation to 10 years of exposure to environmental tobacco smoke at work (69). In a 15-year update of the follow-up of this same population, the odds ratio was 1.21 per 7 years of workplace exposure (95% CI 1.04–1.39) among women, while the effect was statistically nonsignificant for men (risk estimate was not given) (75). In the 15-year study, the subjects lost to follow-up were more likely to have asthma and be exposed to high levels of environmental tobacco smoke, and this selected loss of subjects may explain the decrease in the risk estimate between the two follow-up studies (10). The Finnish case-referent study included 231 never-smoking cases of new asthma and 487 neversmoking referents (72). The risk of asthma was significantly increased in relation to exposure to environmental tobacco smoke in the past year, both at work (OR 2.16, 95% CI 1.26–3.72) and at home (OR 4.77, 95% CI 1.29-17.7). Another study from Finland estimated that the attributable fraction of asthma mortality from exposure to environmental tobacco smoke at work was 4.5% in 1996 (76).

Clear evidence of a dose-response relation was found in four studies (44, 69, 71, 72, 75). In the longitudinal study from the United States, the risk of asthma was **Table 5.** Epidemiologic studies on environmental tobacco smoke and development of asthma among adults. [95% CI = 95% confidence interval, NS = statisically not significant (value not given), OR = odds ratio, PEF = peak expiratory flow]

Study	Study design	Study population, age	Exposure assessment	Definition of asthma	Exposure category compared with no exposure	Results
Greer et al 1993, United States (69)	Longitudinal 10-year follow- up (1977– 1987)	1414 men, 2500 women, nonsmokers, ≥25 years	Self-report of exposure at work and home (in years of exposure)	Self-report of incident physician-diagnosed asthma + wheezing or attacks of short- ness of breath	Work exposure Home exposure	Work exposure: men & women ^a OR 1.45 (95% Cl 1.21–1.80), men ^a OR 1.50 (95% Cl 1.12–2.01), women ^a OR 1.50 (95% Cl 1.17–1.92) Home exposure: no significant relation with asthma
Ng et al 1993, Singapore (43)	Cross- sectional	1282 women, never smokers, 20–74 years	Self-report of household members' ever smoking	Self-report of physi- cian-diagnosed asthma with symp- toms in the past year	Light expo- sure ^b Heavy expo- sure ^b	Light exposure: OR 0.86 (95% CI 0.34–2.21) Heavy exposure: OR 1.60 (95% CI 0.69–3.70)
Leuenberger et al 1994, Switzerland (44)	Cross- sectional	1674 men, 2523 women, never smokers, 18–60 years	Self-report of exposure at home and work during the past 12 months	Self-report of physi- cian-diagnosed asthma or current asthma (physi- cian-diagnosed asthma + wheezing or cough in past 12 months)	Any exposure °	Physician-diagnosed asthma: OR 1.42 (95% Cl 1.05–1.91) Current asthma: OR 1.62 (95% Cl 1.10– 2.37)
Flodin et al 1995, Sweden (70)	Population- based case- referent study	79 cases, 304 referents, of these never smokers: 29 cases, 159 refe- rents, 20–65 years	Self-report of exposure at work and home for \geq 3 years be- fore the diagno- sis of asthma (year of diagno- sis used for matched refe- rents)	Onset in adulthood + clinical history of reversible obstruction and positive metha- choline challenge or significant rever- sibility in beta-2 agonist test	Exposure at work and home	Exposure at work: crude OR 1.5 (95% Cl 0.8–2.6), adjusted OR 1.5 (95% Cl 0.8–2.5) Exposure at home: crude OR 0.9 (95% Cl 0.5–1.5)
Hu et al 1997, United States (71)	Cross- sectional (exposure assessment 7 years earlier)	1469 men and women, non- smokers, 20–22 years	Parents' report of smoking in a previous survey 7 years earlier	Self-report of physi- cian-diagnosed asthma, or current asthma (physician- diagnosed asthma + symptoms or asthma medication in the past year)	Maternal smok- ing, ^a paternal smoking, ^a 1 par- ent smoking, ^a 2 parents smok- ing, ^d	Physician-diagnosed asthma: maternal smoking OR 1.8 (95% Cl 1.1–3.0), paternal smoking OR 1.6 (95% Cl 1.1–2.4), 1 parent smoking OR 1.3 (95% Cl 0.9–2.0), 2 parents smoking OR 2.9 (95% Cl 1.6–5.6) Current asthma: maternal smoking OR 1.8 (95% Cl 1.0–3.2), paternal smoking OR 1.8 (95% Cl 1.1–2.7), 1 parent smoking OR 1.1 (95% Cl 0.7–1.9), 2 parents smoking OR 3.3 (95% Cl 1.7–6.4)
Jaakkola et al 2001, Finland (72)	Population- based case- referent study	231 incident cases, 487 refe- rents, never smokers, 21–63 years	Self-report of exposure at home and work as cigarettes/ day in the past 12 months or lifetime cumula- tive exposure in cigarette-years	History of asthmatic symptoms +rever- sible airflow obstruc- tion in lung function measurements (spirometry+broncho- dilation test, diurnal PEF follow-up)	Exposure in the past 12 months at home and work	At work: OR 2.16 (95% CI 1.26–3.72) At home: OR 4.77 (95% CI 1.29–17.7)

^a OR per 10 years of exposure.

b Light exposure = ≥1 household members smoke or had smoked <20 cigarettes per day, heavy exposure = ≥1 household members smoke or had smoked ≥20 cigarettes per day.</p>

^c Exposure at home or at work during the past 12 months, OR and 95% CI are from an analysis excluding subjects with end-expiratory carbon monoxide levels of ≥7 ppm in order to reduce the misclassification of exposure.

^d Mother smoking >0.5 packs/day versus mother not smoking or smoking ≤0.5 packs/day, father smoking >0.5 packs/day versus father not smoking or smoking ≤0.5 packs/day, one or two parents versus no parent smoking >0.5 packs/day.

related to the duration of workplace exposure, as discussed before (69, 75). In a Swiss cross-sectional study, the risk of physician-diagnosed asthma increased in a dose-dependent relation with the hours of exposure per day, the number of smokers to whom the subject was exposed, and the years of exposure (44). In a Finnish

study the risk of asthma showed a dose-dependent increase with cumulative lifetime exposure from work and home sources (72).

In summary, a limited number of etiologic studies on environmental tobacco smoke and adult-onset asthma has been published, and they indicate an increased

risk of asthma among persons exposed at home or at work. A dose-response relation between exposure and the risk of asthma has been found in several studies. Confounding or exposure misclassification does not seem to explain the observed relations. All the studies controlled for confounding, although the set of confounders varied. They included age, gender, education, occupational exposures, atopy, family history of asthma, and outdoor air pollution. The exclusion of subjects with an end-expiratory level of carbon monoxide of 7 ppm or more, to reduce the misclassification of some actual smokers as nonsmokers, did not have any major effect on the risk estimates in the Swiss study (44). Two studies based the definition of asthma on objective measurements (70, 72), while the others relied on selfreports of diagnosed asthma, the latter being vulnerable to both nondifferential and differential misclassification. More studies with a longitudinal design and objectively examined incident cases of asthma are needed for better risk estimates.

Severity and exacerbations of asthma. In communityand hospital-based surveys (77–79), 69–78% of asthma patients have reported that cigarette smoke aggravates symptoms. Only five epidemiologic studies (80–84) have addressed the role of environmental tobacco smoke in aggravating asthma in adults (table 6).

In a cross-sectional study from India including 200 never-smoking asthmatics, daily exposure to environmental tobacco smoke was significantly related to an increased need for asthma medication, an increased number of acute episodes of asthma and emergency department visits, and lower FEV_1 and FEF_{25-75} levels (81). Another study by this group found that, of 50 nonsmoking asthmatic women, those exposed to environmental tobacco smoke from a spouse had more severe bronchial hyperresponsiveness in histamine challenge than those unexposed (83). These two exposure groups were similar in terms of duration of asthma and exposure to biomass fuel. In a panel study of 164 asthmatic nonsmokers from the United States, daily exposure to environmental tobacco smoke was related to an increased risk of moderate or severe cough, moderate or severe breathlessness, and nocturnal asthma symptoms (80). In a health maintenance organization-based study of 619 asthmatics from the United States, regular exposure was related to worse asthma-specific quality of life, as well as to worse physical functioning and general health (82). In a 30-month follow-up, exposure to environmental tobacco smoke at the beginning of the study was related to an increased rate ratio of health care utilization due to asthma, including hospitalizations, emergency department visits, and urgency care visits. In another study of 285 nonsmoking asthmatics from northern California, self-reported regular exposure was related to greater

asthma severity, worse asthma-specific quality of life, and worse physical health status (Drs Eisner and Blanc 2001, personal communication). Asthmatics regularly exposed to environmental tobacco smoke had a significantly increased risk of emergency department visits, urgent physician visits, and hospitalizations. Asthmatics whose exposure ceased during an 18-month followup period experienced a reduction in asthma severity, improvement in physical health status, and a decrease in health care utilization. The latter two studies assessed exposure both at home and at work. All of these studies focused on patient populations recruited from physician practices or health maintenance organizations.

In Sweden a population-based study of 2065 adults was carried out as part of the European Community Respiratory Health Survey (84). The relative risk of work-associated symptomatic asthma, defined as self-reported asthma, a methacholine test positive for nonspecific airway hyperresponsiveness and reported work-related chest tightness or wheezing, was 1.7 in relation to regular workplace exposure. Statistically significant increases in the risk of work-related chest tightness were observed for the men (prevalence ratio 4.2, 95% CI 1.8–9.8) and for subjects who were atopic according to skin prick tests (prevalence ratio 2.5, 95% CI 1.3–4.8).

In addition to the epidemiologic studies, several experimental studies have been published on the acute effects of environmental tobacco smoke on adults with established asthma. These chamber exposure studies have been reviewed earlier (6, 7). The results of the experimental studies have been somewhat inconsistent, and their interpretation is hampered by small sample sizes, differences in the selection criteria applied to recruit asthmatics, variable exposure times (ranging from 1 to 6 hours), and variable methods used to assess outcome. Controlled chamber exposure studies have the strengths of measuring exposure and outcome more precisely than is usually possible in epidemiologic studies. On the other hand, they have the following weaknesses that reduce their sensitivity to detect any effects of environmental tobacco smoke on asthma: (i) only those with stable asthma can be exposed, although asthmatics in poor control are probably more sensitive to adverse effects, (ii) asthmatics with a recent respiratory infection are often excluded, although these subjects are likely to be more sensitive, and (iii) exposure periods are often short. The experimental studies suggest that there is a subpopulation of asthmatics sensitive to environmental tobacco smoke. These asthmatics experience increased respiratory symptoms, decreased lung function, and increased bronchial hyperresponsiveness in response to exposure to environmental tobacco smoke. Physiological responses of sensitive asthmatics appear to be reproducible after 2 years, and pretreatment with bronchodilating

Table 6. Epidemiologic studies on environmental tobacco smoke and exacerbations or severity of asthma among adults. (95% CI = 95% confidence interval, FEF_{25-75} = forced expiratory flow during the middle half of the FVC, FEV_1 = forced expiratory volume in 1 second, FVC = forced vital capacity, OR = odds ratio, PD_{20} = cumulative dose of histamine at which FEV_1 declined 20% from the baseline value, QOL = quality of life.)

First author, year, country	Study design	Study population, age, diagnosis of asthma	Exposure assessment	Outcomes of asthma morbidity	Results
Ostro et al 1994, United States (80)	3 months follow-up	53 men and 111 women with asthma, nonsmokers, 18–70 years, asthma diagnosed as history and signs of airway ob- struction and reversible obstruction in spirometry	Self-report in baseline ques- tionnaire (any household smoker) and daily diary (ex- posure at home or exposure to irritat- ing smoke, dust or fumes at work)	Daily diary of occur- rence and severity score of respiratory symptoms and nocturnal asthma (= awakened by asthma)	Daily exposure: moderate or severe cough OR 1.21 (95% CI 1.01–1.46), moderate or severe breathlessness OR 1.85 (95% CI 1.57–2.18), nocturnal asthma OR 1.24 (95% CI 1.00–1.53) Occupational exposures ^a : moderate or severe cough OR 1.14 (95% CI 0.98–1.33), moderate or severe breathlessness OR 1.38 (95% CI 1.19–1.59), nocturnal asthma OR 1.19 (95% CI 1.00–1.42) Chronic exposure ^b : moderate or severe breath- lessness OR 2.05 (95% CI 1.78–2.40)
Jindal et al 1994, India (81)	Cross- sectional	200 patients with asthma, never smokers, 15–50 years, asthma diagnosed as history of asthmatic symptoms and reversible airways obstruction in spirometry (excluded if acute severe asthma in preceding 2 weeks)	Self-report of smok- ing by spouse and other close contacts. minimum expo- sure 1 hour/day (or 7 hours/week) for at least 1 year	Self-report of use of bronchodilators, cor- ticosteroids, number of visits to emergency department, hospital- izations, acute epi- sodes and spirometry (FVC, FEV _{1.0} , FEF ₂₅₋₇₅ % of predicted)	Unexposed: OR 0.60, 0.33, 0.60, 8.6, 36.3 & 3.0 for visits/patient to emergency department in last year, hospitalizations/patient in last year, acute episodes/patient in last year, weeks of required steroid treatment per patient, weeks of bronchodilation maintenance in last year & weeks of absence from work in last year, respectively; mean level of % predicted for FVC, FEV ₁ & FEF ₂₅₋₇₅ 90.9, 80.8 & 75.7, respectively Exposed °: OR 0.82 °, 0.34, 1.32 °, 11.3 °, 38.3 & 3.6 ° for visits/patient to emergency department in last year, hospitalizations/patient in last year, acute episodes/patient in last year, weeks of bronchodilation maintenance in last year, weeks of required steroid treatment in last year, weeks of bronchodilation maintenance in last year, respectively; mean level of % predicted for FVC, FEV ₁ & FEF ₂₅₋₇₅ 89.4, 68.7 ° & 54.3 °, respectively
Sippel et al 1999, United States (82)	Cross- sectional and 30 months of follow-up	619 patients with asthma, smokers and nonsmokers, 15–55 years, asthma diagnosed as hospitalization for asthma within 2 years and current use of asthma medication	Self-report of regular exposure at home or work	Asthma-specific QOL, health status Follow-up (from administrative data bases): health care utilization, including hospitalizations, emergency depart- ment visits, urgency care visits	Cross-sectional (unexposed): OOL index 2.3 for breathless and 2.0 for mood, health status index 81 for physical function, 71 for bodily pain, and 65 for general health Cross-sectional (regularly exposed): QOL index 2.7 ° for breathless and 2.5 ° for mood; health status index 73 ° for physical function, 65 ° for bodily pain, and 59 ° for general health Longitudinal (regularly exposed at baseline): RR ° 2.34 (95% CI 1.80–3.05) for health care utilization
Jindal et al 1999, India (83)	Cross- sectional	50 women, 20–40 years, nonsmokers, asthma diagnosed as reversible airway obstruction in lung function tests	Self-report of exposure from spouse, exposure index = average daily hours×years of exposure	Bronchial hyperres- ponsiveness in histamine challenge, PD ₂₀ (mg/ml)	Unexposed: PD_{20} 6.1 Exposed: PD_{20} 1.7 [†] Exposure index <15: PD_{20} 3.2 Exposure index ≥15: PD_{20} 1.8 [†]
Blanc et al 1999, Sweden (84)	Cross- sectional, population- based + en- riched with symptoma- tic subjects	1013 men and 1052 wo- men, 20–44 years, 17% had self-reported asthma, smokers and nonsmokers	Self-report of regu- lar exposure at work	Reported asthma, bronchial hyperres- ponsiveness in methacholine chal- lenge and report- ed chest tightness or wheezing at work	Workplace exposure: prelevance ratio 1.7 (95% Cl 0.9–3.3)

^a Occupational exposures = daily exposure to irritating smoke, dust or fumes at work.

^b Chronic exposure = report of any household smoker in the baseline questionnaire.

^c Exposed = asthmatics with at least the minimum exposure (as defined in exposure assessment).

^d Statistically significant difference between exposed and unexposed (P<0.05).

^e Rate ratio related to regular exposure at baseline.

[†] Statistically significant difference between exposed and unexposed, or between high and low exposure groups (P<0.01).

medication seems to prevent these responses (27, 85). Determinants of this sensitivity are not known.

In summary, a limited number of epidemiologic studies and several experimental studies suggest that exposure to environmental tobacco smoke contributes to the severity and exacerbation of asthma among adults with established disease. More epidemiologic studies are needed, with focus on longitudinal design, better outcome and exposure assessment, and control for other factors affecting the stability of asthma. Experimental studies are hampered by the fact that the subgroups probably most sensitive to the adverse effects of environmental tobacco smoke cannot be exposed due to safety reasons, but studies focusing on real life type of exposures can contribute to a better understanding of the potential mechanisms of effects.

Chronic obstructive pulmonary disease

Chronic obstructive pulmonary disease (COPD) develops slowly over the years. Studies of chronic respiratory symptoms and lung function have investigated the early stages of the disease, while studies on diagnosed COPD focus on later stages. Three longitudinal studies (73, 86, 87) and three case-referent studies (74, 88, 89) have evaluated the effects of environmental tobacco smoke on the development of late stage COPD. They are summarized in table 7. A longitudinal study of Seventh-Day Adventists in California has produced several reports (73, 90-92). We were not able to identify any investigation that studied the role of environmental tobacco smoke in the exacerbation of COPD directly by measuring respiratory impairment. However, two studies assessed the impact of exposure on respiratory-related activity restrictions (93, 94).

The fact that COPD was defined in different ways in the epidemiologic studies reflects the complex nature of the disease. It overlaps to some extent with asthma, and some studies included asthma in their definition of obstructive disease (73, 74, 86, 92). The definitions used in the studies were based on symptoms or diagnoses made by a physician as reported in questionnaires, lung function measurements, diagnostic codes in mortality statistics, or a combination of these variables.

An increased risk of COPD in relation to exposure to environmental tobacco smoke was found in all the longitudinal studies. The excess risk was estimated to be from 30% to over 400%. The study from California showed the highest risk of incident COPD in relation to cumulative exposure to environmental tobacco smoke from childhood and adulthood home and workplace exposures (73). One hospital-based case-referent study from Greece (89) and one population-based case-referent study from the United States (74) showed a doseresponse relation between the amount of exposure to environmental tobacco smoke and the risk of COPD. A small hospital-based case-referent study from the United Kingdom observed an increased risk of COPD only in relation to the highest exposure level (88). Workplace exposure to enrironmental tobacco smoke was assessed only in the longitudinal study of the Seventh-Day Adventists (73, 92), while the others were limited to household exposures. The study of Seventh-Day Adventists was also the only study that evaluated the role of childhood exposure for COPD. A Finnish study estimated that the attributable fraction of COPD mortality from exposure to environmental tobacco smoke at work was 1.1% in 1996 (76).

In summary, the number of studies on environmental tobacco smoke and the development of late stage COPD is limited, but all the published studies show an excess risk in relation to exposure. A dose-response relation was found in three studies. Three studies were longitudinal in design and therefore indicated a meaningful temporal relation. Age and gender were taken into account as potential confounders in all of the studies, but otherwise control for confounding varied, including factors such as the participant's or spouse's occupation, housing quality, other indoor pollutants, and outdoor pollution. One study examined potential modification of the effects of environmental tobacco smoke by ambient air pollutants, but no significant interactions were found (73). More studies with better outcome and exposure assessment are needed before any definite conclusions can be drawn.

Two studies addressed the role of exposure to environmental tobacco smoke in respiratory-related activity restrictions (93, 94). Both studies were based on data collected by national health interview surveys conducted by the Census Bureau for the National Center for Health Statistics in the United States. In the first report, among nonsmokers, respiratory-related restricted activity in the past 2 weeks increased by an average of 1% per 1 cigarette smoked per day by a household member when the situation was compared with nonsmoking households (93). In the second report, respiratory disease exacerbation was defined as activity limitation or physician visit in the preceding 2 weeks because of chronic bronchitis, asthma, emphysema, or chronic sinusitis (94). Among a probability sample of never smokers, the risk of exacerbations of respiratory conditions increased significantly in relation to exposure to environmental tobacco smoke at home or work, with an OR of 1.44 (95% CI 1.07-1.95). These two studies suggest that exposure to environmental tobacco smoke contributes to the adverse consequences of COPD. However, there is a lack of studies evaluating more directly the role of environmental tobacco smoke as a potential trigger of COPD exacerbation.

Respiratory infections

There is convincing evidence that passive smoking enhances susceptibility to respiratory infections among children (2). For adults, surprisingly little data have been published on infections. Recently, a population-based case-referent study from the United States evaluated the relations of active smoking and exposure to environmental tobacco smoke with invasive pneumococcal disease in immunocompetent adults (95). Among a total of 228

Table 7. Epidemiologic	studies on	environmental	tobacco	smoke and	chronic obstructive	pulmonary	disease	(COPD).	(95%	CI =
95% confidence interval,	OR = odds	ratio)								

Study	Study design	Study population, age	Exposure assessment	Definition of COPD outcome	Exposure category compared to no exposure	Results
Hirayama 1981, Japan (86)	Longitudinal, 14-year follow- up (1965– 1979)	91 540 women, nonsmokers, ≥40 years	Spouse's report of smoking in the beginning of follow-up	Mortality from emphy- sema or asthma, based on diagnosis in death certificate	Spouse ex-smoker or smoked 1–19 cigarettes per day; spouse smoked ≥20 cigarettes/day	OR 1.29 ^a for spouse ex-smoker or smoking 1-19 cigarettes/day and OR 1.49 ^a for spouse smoking ≥20 cigarettes/day
Lee et al 1986, United Kingdom (88)	Hospital-based case-referent study	chronic bronchitis cases: 9 men, 17 women; referents: 133 men, 318 women, never smokers, 35–74 years	Self-report of spouse's cur- rent and past smoking habits and ETS expo- sure at work, during daily travel and during leisure time	Chronic bronchitis, diagnosed in a hospital (criteria for diagnosis not given)	Spouse's smoking during whole marriage; quantita- tive index ° 2–4 and 5–12	Men ^b : OR 0.34 for spouse's smoking during whole marriage; OR 0.83 & 1.90 for quantitative index of 2–4 & 5–12, respectively Women ^b : OR 1.22 for spouse's smoking during whole marriage; OR 1.05 & 1.03 for quantitative index of 2–4 & 5–12, respectively Combined ^b : OR 0.83 for spouse's smoking during whole marriage; OR 1.00 & 1.30 for quantitative index of 2–4 & 5–12, respectively
Kalandidi et al 1987, Greece (89)	Hospital-based case-referent study	103 COPD cases, 179 referents (visitors in the hospital), women, never smokers, 40–79 years	Self-report of spouse's current and past smo- king habits	COPD diagnosis, based ton history of dyspnea and phlegm for ≥3 years + obstruction or mixed type reduction in spirometry (no reversi- bility with broncho- dilator)	Spouse ex-smoker, spouse smokes ≤ 1 or >1 packs/day Spouse's lifelong , consumption (x 10 ³) \leq 300 or > 300	OR 0.7 (95% Cl 0.3–1.4) for spouse ex-smoker; 2.5 (95% Cl 1.3–5.0) & 1.5 (95% Cl 0.8–2.7) for spouse smoking \leq 1 & >1 packs/day, respectively; 1.3 (95 Cl 0.7–2.3) & 1.7 (95% Cl 0.8–3.4) for \leq 300 x 10 ³ & >300 x 10 ³ of spouse's lifelong consumption, respectively
Sandler et al United States (87)	Longitudinal, 12-year follow- up (1963–1975)	4162 men, 14873 women, never smokers, ≥25 years at baseline	Household members' report of smoking in the beginning of thestudy	Mortality from emphy- sema or bronchitis = underlying cause of death in death certificate	Smoking cohabitees	Men: rate ratio 0.93 (95% CI 0.16–5.32) Women: rate ratio 5.65 (95% CI 1.19–26.8)
Robbins et al 1993, United States (73)	Longitudinal, 10-year follow- up (1977–1987)	1414 men, 2500 women, nonsmokers, ≥25 years at baseline	Self-report of exposure at work and home during adult- hood and at home during childhood	Incident airways obs- tructive disease, physician diagnosed asthma + wheezing or cough and sputum for ≥3 months/year for ≥2 years or physi- cian-diagnosed emphy- sema + breathlessness	Childhood exposure only, past adult- hood exposure, child- hood + past adult- hood exposure, cur- rent adult home + work exposure, child- hood + current adult home + work expo- sure, reference cate- gory= never exposed	OR 1.09 (95% Cl 0.69–1.68), 1.33 (95% Cl 0.93–1.88),1.68 (95% Cl 1.27–2.20), 1.48 (95% Cl 0.95–2.23), & 2.03 (95% Cl1.45–2.77) for childhood exposure, childhood + past adulthood exposure, current adulthood exposure, current adult home + work exposure & childhood + current adult home + work exposure, respectively
Dayal et al 1994, United States (74)	Population- based case- referent study	219 cases, 657 referents, never smokers, age range not given	Household head's report of current smoking habits of household members	Household head's report of obstructive respira- tory disease, defined as diagnosed asthma, chronic bronchitis or emphysema	Exposure ≤1 or >1 packs/day	OR 1.16 (95% Cl 0.78–1.72) for ≤1 packs/day & 1.86 (95% Cl 1.21–2.86) for >1 packs/day

^a Risk ratio (these were statistically nonsignificant).

^b Reference category = spouse did not smoke during whole marriage or quantitative index score 0–1. All OR values were statistically nonsignificant, but the 95% CI were not given.

^c Quantitative index for regular exposure was calculated based on a sum of 0 = not at all, 1 = a little, 2 = average, 3 = a lot exposure at home, at work, during travel, and during leisure.

patients, 95% had bacteremia, 4% had meningitis, and 1% had infections at other normally sterile sites. Among nonsmokers, patients with invasive pneumococcal infections were 2.5 times as likely to be exposed to environmental tobacco smoke as referents (95% CI 1.2–5.1) when confounding by gender, race, presence of chronic illnesses, education, and living with children going to

day care were adjusted for. The odds ratios were of similar magnitude for persons exposed to smoke at home only and for those exposed only outside the home. The population attributable risk for passive smoking was 17%. A dose-response relation was observed between the hours of daily exposure and the risk of invasive pneumococcal disease. Two earlier studies have suggested that exposure to environmental tobacco smoke is related to the risk of meningococcal disease in adults (96, 97). In summary, for adults, exposure to environmental tobacco smoke has been associated with serious infections, such as bacteremic pneumococcal pneumonia and meningitis, but more studies examining the relations between environmental tobacco smoke and infections are needed.

Discussion of methodological issues

Avoidance of exposure to environmental tobacco smoke because of a health condition (eg, respiratory symptom or disease) could lead to an underestimation of the true effect of environmental tobacco smoke. Longitudinal studies recording exposure before the first symptoms and signs of the disease minimizes this type of selection bias, which is more difficult to avoid in cross-sectional and prevalent case-referent studies. In longitudinal studies, subjects lost to follow-up may introduce selection bias if the loss is related to both exposure and outcome. Attempts to maintain a high follow-up rate help to minimize this type of selection bias. The selection of referents in case-referent studies may introduce selection bias if the referents do not represent the exposure of the source population that produced the cases. Especially hospital-based referent selection is susceptible to this type of bias if the referents are chosen among a patient group whose disease is related to exposure to environmental tobacco smoke. Recruitment of all new cases and concurrent recruitment of a random sample of referents from the same source population will eliminate this type of bias. Recently there has been a tendency to carry out population-based case-referent studies to avoid this type of bias.

Information on outcome should be collected similarly from the exposed and unexposed subjects so that awareness of exposure does not influence the measurement of outcome. Any differences in the outcome assessment between the exposed and unexposed subjects will lead to information bias, which may either increase or decrease the studied relation. Nondifferential or random error will usually lead to an effect towards the null. Studies of subjective outcomes, such as respiratory symptoms, are susceptible to both differential and nondifferential outcome misclassification.

In case-referent studies, information on exposure to environmental tobacco smoke from cases and referents should be comparable. Retrospective, subjective exposure assessment may be problematic. Recall bias takes place if cases remember exposures differently from referents because of their disease. However, lung cancer studies have not supported empirically the occurrence

of recall bias in health effect studies of environmental tobacco smoke (98). Assessment of exposure to environmental tobacco smoke may be complicated, especially in adult studies, because of multiple sources of exposure (eg, home, workplace and different types of social situations). If these multiple sources are not taken into account, the reference category classified as unexposed includes subjects who, in fact, experience substantial exposure. This misclassification dilutes the risk estimate obtained. It has been estimated that ≥80% of nonsmokers have detectable cotinine levels in their body fluids (99, 100). Recently studies have started to pay attention to other sources of environmental tobacco smoke, in addition to household smoking, to achieve better risk estimates. A potentially important source of misclassification of exposure is the misclassification of smoking status due to self-reporting. Some current and former smokers may report themselves as never smokers and be at higher risk of developing smoking-related diseases. It has been suggested that smokers are more likely to marry smokers than nonsmokers are (101). If so, a bigger proportion of those classified as exposed to environmental tobacco smoke are active smokers when a comparison is being made to persons classified as unexposed. However, the proportion of ever smoking adults (current or former) who are misclassified as never smokers has been estimated to be relatively small, 3% to 7% (5, 101-103). This magnitude of misclassification is not large enough to explain the observed risks of lung diseases related to exposure to environmental tobacco smoke. Furthermore, there is no evidence of such differential misclassification in workplace exposure.

Potential confounders are determinants of the studied disease and are therefore disease-specific. Most of the studies reviewed adjusted for several of the important confounders for respiratory diseases, including age, gender, socioeconomic status, atopy, other indoor pollutants, and occupational exposures, the more recent studies taking into account potential confounding more completely. Focus has recently been on confounding by life-style factors, such as diet. However, adjustment for dietary factors has not changed essentially the risk estimates obtained in lung cancer studies, although the relation between diet and cancer has been relatively strongly established (98, 104-109). In summary, it is not likely that confounding plays a decisive role in the respiratory effects observed in relation to exposure to environmental tobacco smoke. Interactions between other life-style factors and environmental tobacco smoke have been studied less (ie, if an accumulation of several risk factors is more dangerous than the addition of risks related to individual factors separately).

Publication bias will be introduced if scientists submit manuscripts or the journals accept them on the basis of their results, typically so that studies finding no effects ("negative studies") are underrepresented. In 1994 Bero and her co-workers (110) carried out a review of published and unpublished studies of the health effects of environmental tobacco smoke to assess potential publication bias. The authors concluded that there is no evidence of publication bias against statistically nonsignificant results in the peer-reviewed literature on environmental tobacco smoke. Lung cancer in adults and respiratory diseases in children have been studied extensively, and this abundance reduces the likelihood of publication bias. However, for diseases that have been studied to a limited extent, it is important that all goodquality research be published so that an unbiased picture can be obtained. A more recent comparison of published and unpublished studies on passive smoking showed a publication delay for studies with nonsignificant results (111).

In summary, it is important to be aware of the many potential sources of bias in studies of the health effects of exposure to environmental tobacco smoke. However, especially recent studies have put a great deal of effort on avoiding these biases and adjusting for confounding. Thus an abundant amount of high-quality research exists that confirms many of the observed relations between exposure to environmental tobacco smoke and respiratory disease.

Summary

Sidestream smoke contains the same irritative and toxic compounds as mainstream smoke, and it is plausible that the biological effects of environmental tobacco smoke are similar to those of active smoking. All of the respiratory effects reviewed (ie, respiratory symptoms, lung function impairment, asthma, COPD, and pneumococcal infections) have been linked to both home and workplace exposure among adults. Some evidence of a dose-response relation has been detected for all of these conditions. Table 8 summarizes the information on the respiratory effects of exposure to environmental tobacco smoke among adults. In addition, the authors' judgment of causality, based on the number of studies, their validity, the evidence of dose-response relations, and biological plausibility, is presented.

The strongest evidence for a causal relation exists for environmental tobacco smoke and chronic respiratory symptoms. Several cross-sectional studies provide evidence that exposure to environmental tobacco smoke is related to deficits in ventilatory lung function, the effects usually being relatively small. However, longitudinal studies on lung function have not confirmed these findings. It is likely that the harmful effects of exposure to environmental tobacco smoke are dependent on the dose, since adverse effects on lung function have been detected mainly in countries and occupations with high exposure levels. A limited number of studies suggests that environmental tobacco smoke increases the risk of new asthma and contributes to poor control of established disease in adults. The evidence linking exposure to environmental tobacco smoke to COPD is also limited, but an adverse effect is supported by three longitudinal studies. There is a lack of studies on the role of environmental tobacco smoke as a potential trigger of COPD exacerbation. One study among adults linked exposure to environmental tobacco smoke to serious infections, including pneumococcal pneumonia and meningitis. There is a need for more longitudinal studies with careful assessment of exposure to provide better risk estimates for all of these respiratory conditions. New studies on asthma, COPD, and respiratory infections are needed before any definite conclusions can be drawn. However, despite these challenges being left for future research, the combination of toxicologic evidence, a large number of studies on environmental tobacco smoke and respiratory effects among children, and the adult studies reviewed in this

Table 8. Summary of environmental tobacco smoke and respiratory diseases and conditions among adults. (95% CI = 95% confidence interval, OR = odds ratio, COPD = chronic obstructive pulmonary disease, EF = effect estimate, FEV_1 = forced expiratory volume in 1 second)

Disease or condition	OR ^a or EF	95% CI	Causality ^b
Chronic respiratory symptoms			+++
Wheezing Cough Phlegm Dyspnea	OR 1.41–2.69 OR 2.61–3.79 OR 1.41–2.05 OR 1.44–4.50		
Asthma			
Induction			++
Young adults ⁰ Adults	OR 1.60–3.30 OR 1.42–1.62		
Bronchial hyperres- ponsiveness ^d	OR 1.37	1.05-1.78	+
Severity	Diverse outcomes, no summary estimate available		++
COPD			
Case-referent studies Longitudinal studies	OR 1.86–2.5 OR 1.68–5.63		++
Respiratory infections ^e	OR 2.5	1.2-5.1	+
Lung function, FEV ₁			
Cross-sectional studies Longitudinal studies	EF [†] -2.7% No significant effect	-4.1%1.2%	+

^a The range of the OR values from the studies reviewed (that showed a

significant association) or the OR and 95% CI from an individual study. ^b Causality as judged by the authors; symbols: +++ = causal relation established, ++ = strong evidence of a causal relation, + = some evidence

of a causal relation, $\vec{0}$ = no clear evidence of a causal relation.

^c From reference 71. ^d From reference 68.

From reference 95.

¹ Effect estimate and 95% CI from a meta-analysis (63): the difference in the FEV₁ between the exposed and unexposed, expressed as a percentage of the level of the unexposed group. paper point to an urgent need for measures to prevent exposure to environmental tobacco smoke among adults.

References

- White JR, Froeb HF. Small-airways dysfunction in nonsmokers chronically exposed to tobacco smoke. N Engl J Med 1980;302:720–3.
- Jaakkola JJK, Jaakkola MS. Effects of environmental tobacco smoke on the respiratory health of children. Scand J Work Environ Health 2002;28 suppl2:71–83.
- US Department of Health and Human Services, Public Health Service. The health consequences of involuntary smoking: a report of the Surgeon General. Washington (DC): US Government Printing Office, 1986.
- National Research Council, Committee on Passive Smoking. Environmental tobacco smoke: measuring exposures and assessing health effects. Washington (DC): National Academy Press, 1986.
- US Environmental Protection Agency (EPA). Respiratory health effects of passive smoking: lung cancer and other disorders. EPA, Office of Health and Environmental Assessment, Office of Research and Development. Washington (DC): EPA, 1992. EPA/600/6–90/006F.
- Tredaniel J, Boffetta P, Saracci R, Hirsch A. Exposure to environmental tobacco smoke and adult non-neoplastic respiratory diseases. Eur Respir J 1994;7:173–85.
- California Environmental Protection Agency, Office of Environmental Health Hazard Assessment. Health effects of exposure to environmental tobacco smoke. Sacramento (CA): California Environmental Protection Agency, 1997.
- Coultas DB. Passive smoking and risk of adult asthma and COPD: an update. Thorax 1998;53:381–7.
- Jaakkola MS. Environmental tobacco smoke and respiratory diseases. In: Annesi-Maesano I, Gulsvik A, Viegi G, editors. Respiratory epidemiology in Europe. Sheffield (UK): European Respiratory Society, 2000:322–83. Monograph no 15.
- Eisner MD, Blanc PD. Environmental tobacco smoke and adult asthma. In: Watson R, editor. Environmental tobacco smoke. Boca Raton (FL): CRC Press, 2001;81–105.
- Jaakkola MS, Jaakkola JJK. Assessment of exposure to environmental tobacco smoke. Eur Respir J 1997;10:2384–97.
- Jaakkola MS, Samet JM. Occupational exposure to environmental tobacco smoke and health risk assessment. Environ Health Perspect 1999;107 suppl 6:829–35.
- US Department of Health and Human Services. The health consequences of smoking: chronic obstructive pulmonary disease: a report of the Surgeon General. Washington (DC): US Department of Health and Human Services, 1984. DHHS publication, no 84–50205.
- Graham NM. The epidemiology of acute respiratory infections in children and adults: a global perspective. Epidemiol Rev 1990;12:149–78.
- Strachan DP, Cook DG. Parental smoking and lower respiratory illness in infancy and early childhood. Thorax 1997;52: 905–14.
- Green GM, Carolin D. The depressant effect of cigarette smoke on the in vitro antibacterial activity of alveolar macrophages. N Engl J Med 1967;276:421–7.
- 17. Wanner A. State of the art: clinical aspects of mucociliary

transport. Am Rev Respir Dis 1977;116:73-125.

- Fainstain V, Musher D. Bacterial adherence to pharyngeal cells in smokers, nonsmokers, and chronic bronchitis. Infect Immun 1979;26:178–82.
- Dye JA, Adler KB. Effects of cigarette smoke on epithelial cells of the respiratory tract. Thorax 1994;49:825–34.
- Hoffmann D, Hoffmann I. The changing cigarette, 1950– 1995. J Toxicol Environ Health 1997;50:307–64.
- Wright JL, Lawson LM, Pare PD, Kennedy S, Wiggs B, Hogg JC. The detection of small airways disease. Am Rev Respir Dis 1984;129:989–94.
- Thurlbeck WM. Pathophysiology of chronic obstructive pulmonary disease. Clin Chest Med 1990;11:389–403.
- Pauwels RA, Löfdahl C-G, Pride NB, Postma DS, Laitinen LA, Ohlsson SV. European Respiratory Society study on chronic obstructive pulmonary disease (EUROSCOP): hypothesis and design. Eur Respir J 1992;5:1254–61.
- Nafstad P, Magnus P, Jaakkola JJK. Early respiratory infections and childhood asthma. Pediatrics 2000;106:E38.
- Strachan DP, Cook DG. Parental smoking and childhood asthma: longitudinal and case-control studies. Thorax 1998;53:204–12.
- Tarlo SM, Broder I. Irritant-induce occupational asthma. Chest 1989;96:297–300.
- Stankus RP, Menon PK, Rando RJ, Glindmeyer H, Salvaggio JE, Lehrer SB. Cigarette smoke sensitive asthma: challenge studies. J Allergy Clin Immunol 1988;82:331–8.
- Zetterström O, Österman K, Machado L, Johansson SGO. Another smoking hazard: raised serum IgE concentrations and increased risk of occupational allergy. BMJ 1981;283: 1215–20.
- Kjellman NIM. Effect of parental smoking on IgE levels in children. Lancet 1981;1:993–4.
- Sapigni T, Biavati P, Simoni M, Viegi G, Baldacci S, Carrozzi L, et al. The Po river delta respiratory epidemiologal survey: an analysis of factors related to level of total serum IgE. Eur Respir J 1998;11:278–83.
- 31. Oryszczyn M-P, Annesi-Maesano I, Charpin D, Paty E, Maccario J, Kauffmann F. Relationships of active and passive smoking to total IgE in adults of the epidemiological study of the genetics and environment of asthma, bronchial hyperresponsiveness, and atopy (EGEA). Am J Respir Crit Care Med 2000;161:1241–6.
- Muramatsu T, Weber A, Muramatsu S, Akermann F. An experimental study on irritation and annoyance due to passive smoking. Int Arch Occup Environ Health 1983;51:305– 17.
- Bascom R, Kulle T, Kagey-Sobotka A, Proud D. Upper respiratory tract environmental tobacco smoke sensitivity. Am Rev Respir Dis 1991;143:1304–11.
- Cummings KM, Zaki A, Markello S. Variation in sensitivity to environmental tobacco smoke among adult non-smokers. Int J Epidemiol 1991;20:121–5.
- Lebowitz MD, Burrows B. Respiratory symptoms related to smoking habits of family adults. Chest 1976;69:48–50.
- Schilling RSF, Letai AD, Hui SL, Beck JB, Schoenberg JB, Bouhuys A. Lung function, respiratory disease, and smoking in families. Am J Epidemiol 1977;106:274–83.
- Comstock GW, Meyer MB, Helsing KJ, Tockman MS. Respiratory effects of household exposures to tobacco smoke and gas cooking. Am Rev Respir Dis 1981;124:143–8.
- Schenker MB, Samet JM, Speizer FE. Effect of cigarette tar content and smoking habits on respiratory symptoms in women. Am Rev Respir Dis 1982;125:684–90.

- Hole DJ, Gillis CR, Chopra C, Hawthrone VM. Passive smoking and cardiorespiratory health in a general population in the west of Scotland. BMJ 1989;299:423–7.
- Kauffman F, Dockery DW, Speizer FE, Ferris BG. Respiratory symptoms and lung function in relation to passive smoking: a comparative study of American and French women. Int J Epidemiol 1989;18:334–44.
- White JR, Froeb HF, Kulik JA. Respiratory illness in nonsmokers chronically exposed to tobacco smoke in the work place. Chest 1991;100:39–43.
- Pope CA III, Xu X. Passive cigarette smoke, coal heating, and respiratory symptoms of nonsmoking women in China. Environ Health Perspect 1993;101:314–6.
- 43. Ng TP, Hui KP, Wan CT. Respiratory symptoms and lung function effects of domestic exposure to tobacco smoke and cooking by gas in non-smoking women in Singapore. J Epidemiol Community Health 1993;47:454–9.
- 44. Leuenberger P, Schwartz J, Ackermann-Liebrich U, Blaser K, Bolognini G, Bongard JP, et al. Passive smoking exposure in adults and chronic respiratory symptoms (SAPALDIA study). Am J Respir Crit Care Med 1994;150:1222–8.
- Piitulainen E, Tornling G, Eriksson S. Environmental correlates of impaired lung function in non-smokers with severe α₁-antitrypsin deficiency (PiZZ). Thorax 1998;53:939–43.
- Lam TH, Ho LM, Hedley AJ, Adab P, Fielding R, McGhee SM, Aharonson-Daniel L. Environmental tobacco smoke exposure among police officers in Hong Kong. JAMA 2000; 284:756–63.
- Schwartz J, Zeger S. Passive smoking, air pollution, and acute respiratory symptoms in a diary study of student nurses. Am Rev Respir Dis 1990;141:62–7.
- Jaakkola MS, Jaakkola JJK, Becklake MR, Ernst P. Effect of passive smoking on the development of respiratory symptoms in young adults: an 8-year longitudinal study. J Clin Epidemiol 1996;49:581–6.
- Eisner MD, Smith AK, Blanc PD. Bartenders' respiratory health after establishment of smoke-free bars and taverns. JAMA 1998;280:1909–14.
- Wong CM, Hu ZG, Lam TH, Hedley AJ, Peters J. Effects of ambient air pollution and environmental tobacco smoke on respiratory health of non-smoking women in Hong-Kong. Int J Epidemiol 1999;28:859–64.
- Trinder PM, Croft PR, Lewis M. Social class, smoking and the severity of respiratory symptoms in the general population. J Epidemiol Community Health 2000;54:340–3.
- Kauffman F, Teshier J-F, Oriol P. Adult passive smoking in the home environment: a risk factor for chronic airflow limitation. Am J Epidemiol 1983;117:269–80.
- Kentner M, Triebig G, Weltle D. The influence of passive smoking on pulmonary function — a study of 1351 office workers. Prev Med 1984;13:656–69.
- Salem ES, El Zahby M, Senna GA, Malek A. Pulmonary manifestations among passive smokers. Bull Int Union Tuberc 1984;59:50–3.
- 55. Brunekreef B, Fischer P, Remijn B, Van Der Lende R, Schouten J, et al. Indoor air pollution and its effect on pulmonary function of adult non-smoking women, III: passive smoking and pulmonary function. Intern J Epidemiol 1985;14:227–30.
- Svendsen KH, Kuller LH, Martin MJ, Ockene JK. Effects of passive smoking in the multiple risk factor intervention trial. Am J Epidemiol 1987;126:783–95.
- Masi MA, Hanley JA, Ernst P, Becklake MR. Environmental tobacco smoke and lung function in young adults. Am Rev

Respir Dis 1988;138:296–9.

- Masjedi M-R, Kazemi H, Johnson DC. Effects of passive smoking on the pulmonary function in adults. Thorax 1990;45:27–31.
- Xu X, Li B. Exposure-response relationship between passive smoking and adult pulmonary function. Am J Respir Crit Care Med 1995;151:41–6.
- 60. Frette C, Barrett-Connor E, Clausen JL. Effect of active and passive smoking on ventilatory function in elderly men and women. Am J Epidemiol 1996;143:757–65.
- 61. Casale R, Pasqualetti P. Cosinor analysis of circadian peak expiratory flow variability in normal subjects, passive smokers, heavy smokers, patients with chronic obstructive pulmonary disease and patients with interstitial lung disease. Respiration 1997;64:251–6.
- Abbey DE, Burchette RJ, Knutsen SF, McDonnell WF, Lebowitz MD, Enright PL. Long-term particulate and other air pollutants and lung function in nonsmokers. Am J Respir Crit Care Med 1998;158:289–98.
- Carey IM, Cook DG, Strachan DP. The effects of environmental tobacco smoke exposure on lung function in a longitudinal study of British adults. Epidemiology 1999;10:319–26.
- 64. Jones JR, Higgins JTT, Higgins MW, Keller JB. Effects of cooking fuels on lung function in nonsmoking women. Arch Environ Health 1983;38:219–22.
- Lebowitz MD. Influence of passive smoking on pulmonary function: a survey. Prev Med 1984;13:645–55.
- Jaakkola MS, Jaakkola JJK, Becklake MR, Ernst P. Passive smoking and evolution of lung function in young adults: an 8-year longitudinal study. J Clin Epidemiol 1995;48:317–27.
- Dimich-Ward H, Lawson J, Chan-Yeung M. Work-shift changes in lung function in bar workers exposed to environmental tobacco smoke. Am J Respir Crit Care Med 1998; 157:A505.
- Nowak D, Heinrich J, Jörres R, Wassmer G, Berger J, Beck E, et al. Prevalence of respiratory symptoms, bronchial hyperresponsiveness and atopy among adults: west and east Germany. Eur Respir J 1996;9:2541–52.
- Greer JR, Abbey DE, Burchette RJ. Asthma related to occupational and ambient air pollutants in nonsmokers. J Occup Med 1993;35:909–15.
- Flodin U, Jönsson P, Ziepgler J, Axelson O. An epidemiologic study of bronchial asthma and smoking. Epidemiology 1995;6:503–5.
- Hu FB, Persky V, Flay BR, Richardson J. An epidemiological study of asthma prevalence and related factors among young adults. J Asthma 1997;34:67–76.
- Jaakkola MS, Piipari R, Jaakkola N, Jaakkola JJK. Environmental tobacco smoke and adult-onset asthma: A populationbased incident case-control study. Eur Respir J 2001;18 suppl 33:285s.
- Robbins AS, Abbey DE, Lebowitz MD. Passive smoking and chronic respiratory symptoms in non-smoking adults. Int J Epidemiol 1993;22:809–17.
- Dayal HH, Khuder S, Sharrar R, Trieff N. Passive smoking in obstructive respiratory diseases in an industrialized urban population. Environ Res 1994;65:161–71.
- McDonnell WF, Abbey DE, Nishino N, Lebowitz MD. Longterm ambient ozone concentration and the incidence of asthma in nonsmoking adults: the AHSMOG study. Environ Res 1999;80:110–21.
- 76. Nurminen MM, Jaakkola MS. Mortality from occupational exposure to environmental tobacco smoke in Finland.

J Occup Environ Med 2001;43:687-693.

- Dales RE, Kerr PE, Schweitzer I, Reesor K, Gougeon L, Dickinson G. Asthma management preceding an emergency department visit. Arch Intern Med 1992;152:2041–44.
- Abramson MJ, Kutin JJ, Rosier MJ, Bowes G. Morbidity, medication and trigger factors in a community sample of adults with asthma. Med J Aust 1995;162:76–81.
- 79. Tarlo SM, Broder I, Corey P, Chan-Yeung M, Ferguson A, Becker A, et al. A case-control study of the role of cold symptoms and other historical triggering factors in asthma exacerbations. Can Respir J 2000;7:42–8.
- Ostro BD, Lipsett MJ, Mann JK, Wiener MB, Selner J. Indoor air pollution and asthma: results from a panel study. Am J Respir Crit Care Med 1994;149:1400–6.
- Jindal SK, Gupta D, Singh A. Indices of morbidity and control of asthma in adult patients exposed to environmental tobacco smoke. Chest 1994;106:746–9.
- Sippel JM, Pedula KL, Vollmer WM, Buist AS, Osborne ML. Associations of smoking with hospital-based care and quality of life in patients with obstructive airway disease. Chest 1999;115:691–6.
- Jindal SK, Jha LK, Gupta D. Bronchial hyperresponsiveness of women with asthma exposed to environmental tobacco smoke. Indian J Chest Dis Allied Sci 1999;41:75–82.
- 84. Blanc PD, Ellbjär S, Janson C, Norbäck D, Norrman E, Plasche P, et al. Asthma-related work disability in Sweden: the impact of workplace exposures. Am J Respir Crit Care Med 1999;160:2028–33.
- Menon P, Stankus RP, Rando RJ, Salvaggio JE, Lehrer SB. Asthmatic responses to passive cigarette smoke: persistence of reactivity and effect of medications. J Allergy Clin Immunol 1991;88:861–9.
- Hirayama T. Non-smoking wives of heavy smokers have a higher risk of lung cancer: a study from Japan. Br Med J Clin Res Ed 1981;282:183–5.
- Sandler DP, Comstock GW, Helsing KJ, Shore DL. Deaths from all causes in non-smokers who lived with smokers. Am J Public Health 1989;79:163–7.
- Lee PN, Chamberlain J, Alderson MR. Relationship of passive smoking to risk of lung cancer and other smokingassociated diseases. Br J Cancer 1986;54:97–105.
- Kalandidi A, Trichopoulos D, Hatzakis A, Tzannes S, Saracci R. Passive smoking and chronic obstructive pulmonary disease. Lancet 1987;2:1325–6.
- 90. Euler GL, Abbey DE, Magie AR, Hodgkin JE. Chronic obstructive pulmonary disease symptom effects of long-term cumulative exposure to ambient levels of total suspended particulates and sulfur dioxide in California Seventh-day Adventist residents. Arch Environ Health 1987;42:213–22.
- Abbey DE, Petersen F, Mills PK, Beeson WL. Long-term ambient concentrations of total suspended particulates, ozone, and sulfur dioxide and respiratory symptoms in a non-smoking population. Arch Environ Health 1993;48:33–46.
- Berglund DJ, Abbey DE, Lebowitz MD, Knutsen SF, McDonnell WF. Respiratory symptoms and pulmonary function in an elderly nonsmoking population. Chest 1999;115:49–59.
- Ostro BD. Estimating the risks of smoking, air pollution, and passive smoke on acute respiratory conditions. Risk Anal 1989;9:189–96.
- 94. Mannino DM, Siegel M, Rose D, Nkuchia J, Etzel R. Environmental tobacco smoke exposure in the home and worksite and health effects in adults: results from the 1991 National Health Interview Survey. Tob Control 1997;6:296–305.

- Nuorti JP, Butler JC, Farley MM, Harrison LH, McGeer A, Kolczak MS, et al. Cigarette smoking and invasive pneumococcal disease. N Engl J Med 2000;342:681–9.
- Stuart JM, Cartwright KAV, Dawson JA, Rickard J, Noah ND. Risk factors for meningococcal disease: a case control study in Southwest England. Community Med 1988;10:139– 46.
- 97. Tappero JW, Reporter R, Wenger JD, Ward BA, Reeves MW, Missbach TS, et al. Menongococcal disease in Los Angeles County, California, and among men in the county jails. N Engl J Med 1996;335:833–40.
- Fontham ETH, Correa P, Reynolds P, Wu-Williams A, Buffler PA, Greenberg RS, et al. Environmental tobacco smoke and lung cancer in nonsmoking women: a multicenter study. JAMA 1994;271:1752–9.
- Riboli E, Preston-Martin S, Saracci R, Haley NJ, Trichopoulos D, Becher H, et al. Exposure of nonsmoking women to environmental tobacco smoke: a 10-country collaborative study. Cancer Causes Control 1990;1:243–52.
- 100. Pirkle JL, Flegal KM, Bernert JT, Brody DJ, Etzel RA, Maurer KR. Exposure of the US population to environmental tobacco smoke: the third national health and nutrition examination survey, 1988 to 1991. JAMA 1996;275:1233–40.
- 101. Wu AH. Exposure misclassification bias in studies of environmental tobacco smoke and lung cancer. Environ Health Perspect 1999;107 suppl 6:873–7.
- 102. Nyberg F, Isaksson I, Harris JR, Pershagen G. Misclassification of smoking status and lung cancer risk from environmental tobacco smoke in never-smokers. Epidemiology 1997;8:304–9.
- 103. Hacksaw AK, Law MR, Wald NJ. The accumulated evidence on lung cancer and environmental tobacco smoke. BMJ 1997;315:980–8.
- 104. Hirayama T. Dietary habits are of limited importance in influencing the lung cancer risk among Japanese females who never smoked. In: Bieva CJ, Courtois Y, Govaerts M, editors. Present and future of indoor air quality: proceedings of the Brussels conference, February 1989. Amsterdam: Elsevier Science, 1989.
- 105. Kalandidi A, Katsouyanni K, Voropoulou N, Bastas G, Saracci R, Trichopoulos D. Passive smoking and diet in the etiology of lung cancer among non-smokers. Cancer Causes Control 1990;1:15–21.
- 106. Alavanja MCR, Brown CC, Swanson C, Brownson RC. Saturated fat intake and lung cancer risk among nonsmoking women in Missouri. J Natl Cancer Inst 1993;85:1906–16.
- 107. Mayne ST, Janerich DT, Greenwald P, Chorost S, Tucci C, Zaman MB, et al. Dietary beta-carotene and lung cancer risk in US nonsmokers. J Natl Cancer Inst 1994;86:33–8.
- 108. Cardenas VM, Thun MJ, Austin H, Lally CA, Clark WS, Greenberg S, et al. Environmental tobacco smoke and lung cancer mortality in the American Cancer Society's cancer prevention study II. Cancer Causes Control 1997;8:57–64.
- 109. Boffetta P, Agudo A, Ahrens W, Benhamou E, Banhamou S, Darby SC, et al. Multicenter case-control study of exposure to environmental tobacco smoke and lung cancer in Europe. J Natl Cancer Inst 1998;90:1440–50.
- 110. Bero LA, Glantz SA, Rennie D. Publication bias and public health policy on environmental tobacco smoke. JAMA 1994;272:133–6.
- 111. Misakian AL, Bero LA. Publication bias and research on passive smoking. Comparison of published and unpublished studies. JAMA 1998;280:250–3.