

Scand J Work Environ Health 1993;19(2):7-72 Issue date: 1993

Health risk evaluation of nitrogen oxides.

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/8209198



Print ISSN: 0355-3140 Electronic ISSN: 1795-990X

SUMMARY

The nitrogen oxides discussed in this document are nitric oxide (NO) and nitrogen dioxide (NO₂). In combustion processes, NO is primarily formed and subsequently oxidized to NO₂ in the atmosphere. Traffic is the major source to human outdoor exposure to NO and NO₂ in urban areas. In homes with gasfueled stoves or water heaters, the concentration of NO₂ is generally higher indoors than outdoors.

Upon inhalation, NO₂ penetrates deep into the lung. In animal experiments, NO₂ causes biochemical and morphological changes in lung tissue, affects host defense mechanisms, and causes decreased pulmonary function. The lowest observed effect level after prolonged exposure is 560 μ g·m⁻³.

In short-term controlled studies on humans, NO₂ causes bronchoconstriction and an increase in bronchial responsiveness. A meta-analysis indicates a lowest observed effect level of 200 μ g·m⁻³ for increased bronchial responsiveness in asthmatics.

Epidemiologic studies on children living in homes with gas stoves suggest an increased risk of lower respiratory illness at average indoor NO₂ concentrations of about 40-80 μ g·m³, with short-term peaks exceeding 1000 μ g·m³. The studies focusing on outdoor exposure provide some evidence of increased respiratory disease rates for children and nonsmoking adults living in areas with long-term average NO₂ concentrations of 30-100 μ g·m⁻³; however, the specific role of NO₂ is not clear.

Based on controlled studies on humans, a 1-h guideline value of 100 μ g·m⁻³ for NO₂ in ambient air (eg, as the 99th percentile) is recommended. This value would correspond to a long-term average (half-year mean) of about 40 μ g·m⁻³. The epidemiologic evidence is not considered sufficient for a long-term guideline for NO₂.

NO is formed endogenously for purposes such as signaling in the nervous system, the mediation of vasodilation, and the mediation of cytotoxicity in macrophages. After the inhalation of high concentrations of NO, lung vasodilator effects occur. No health-based guideline value for NO in ambient air is suggested.