



Scand J Work Environ Health 2018;44(4):403-413

<https://doi.org/10.5271/sjweh.3728>

Published online: 18 Apr 2018, Issue date: 01 Jul 2018

Night work and hypertensive disorders of pregnancy: a national register-based cohort study

by [Hammer P](#), [Flachs E](#), [Specht I](#), [Pinborg A](#), [Petersen S](#), [Larsen A](#), [Hougaard K](#), [Hansen J](#), [Hansen Å](#), [Kolstad H](#), [Garde A](#), [Bonde JP](#)

In this nationwide study, we investigated the association of hypertensive disorders of pregnancy with different dimensions of night work objectively assessed through payroll data. Our results indicate that working consecutive night shifts during the first 20 pregnancy weeks increases the risk of hypertensive disorders by 41%, which may be considered when providing recommendations on organization of night work during pregnancy.

Affiliation: Department of Occupational and Environmental Medicine, Bispebjerg University Hospital, Bispebjerg Bakke 23, indgang 20F, 2400 Copenhagen, Denmark. paulahammer@dadlnet.dk

Refers to the following texts of the Journal: [2006;32\(6\):502-514](#)

[2007;33\(4\):304-317](#) [1989;15\(6\):395-403](#) [2010;36\(2\):96-108](#)

[2013;39\(4\):325-334](#) [2015;41\(3\):268-279](#) [2017;43\(1\):59-67](#)

The following articles refer to this text: [2019;45\(6\):577-587](#);

[2020;46\(6\):557-569](#); [2022;48\(7\):520-529](#)

Key terms: [circadian disruption](#); [cohort study](#); [gestational hypertension](#); [hypertension](#); [hypertensive disorder](#); [night work](#); [payroll data](#); [preeclampsia](#); [pregnancy](#); [register-based cohort study](#); [shift work](#); [shift worker](#); [work schedule](#); [working time](#)

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/29669140

Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](#).



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Night work and hypertensive disorders of pregnancy: a national register-based cohort study

by Paula Hammer, MD,¹ Esben Flachs, MSc, PhD,¹ Ina Specht, MSc, PhD,² Anja Pinborg, MD, PhD,³ Sesilje Petersen, MSc, PhD,¹ Ann Larsen, MSc, PhD,⁴ Karin Hougaard, MSc, PhD,^{4,5} Johnni Hansen, MSc, PhD,⁶ Åse Hansen, MSc, PhD,^{4,5} Henrik Kolstad, PhD,⁷ Anne Garde, PhD,^{4,5} Jens Peter Bonde, MD, PhD^{1,5}

Hammer P, Flachs E, Specht I, Pinborg A, Petersen S, Larsen A, Hougaard K, Hansen J, Hansen Å, Kolstad H, Garde A, Bonde JP. Night work and hypertensive disorders of pregnancy: a national register-based cohort study. *Scand J Work Environ Health* 2018;44(4):403–413. doi:10.5271/sjweh.3728

Objective The aim of this study was to investigate whether night work expressed by number and duration of night shifts, number of consecutive night shifts, and number of quick returns during the first 20 weeks of pregnancy is a risk factor for hypertensive disorders of pregnancy (HDP).

Methods The study population comprised Danish workers in public administration and hospitals who gave birth between 2007 and 2013. Exposure was assessed objectively through payroll data. Information on the outcome was retrieved from the National Patient Register. We performed logistic regression on the risk for HDP according to night work adjusted for age, body mass index (BMI), parity, socioeconomic status, and sickness absence prior to pregnancy.

Results Among 18 724 workers, 60% had at least one night shift during the first 20 weeks of pregnancy. The prevalence of HDP was 3.7%. Among night workers, the risk of HDP grew with increasing number of consecutive night shifts [odds ratio (OR) 1.41, 95% confidence interval (CI) 1.01–1.98] and of quick returns after night shifts (OR 1.28, 95% CI 0.87–1.95). Among obese women (body mass index ≥ 30 kg/m²), those who worked long night shifts and longer spells of consecutive night shifts, and had the highest number of quick returns after night shifts, had a 4–5 fold increased risk of HDP compared to day workers.

Conclusion Working consecutive night shifts and quick returns after night shifts during the first 20 pregnancy weeks was associated with an increased risk of HDP, particularly among obese women.

Key terms circadian disruption; gestational hypertension; hypertension; payroll data; preeclampsia; shift work; shift worker; work schedule; working time.

Around 14% of the female European workers <50 years engage in night work (1). Several studies have investigated adverse pregnancy outcomes in relation to work schedules during pregnancy (2–6), but studies focusing on the pregnant women's health are sparse (7–9).

Hypertensive disorders of pregnancy (HDP) including preeclampsia and gestational hypertension occur in around 8% and 5% of pregnancies worldwide and in Denmark, respectively, and are a major cause of morbidity and mortality (10–12). It is suggested that the incidence of HDP has increased over time probably

due to advanced maternal age and increased occurrence of obesity and diabetes in mothers (13, 14). The pathophysiology of HDP is not fully elucidated but seems to involve maternal, fetal and placental factors (14–19).

Night work, including both fixed night shifts and shift work, may influence the risk of HDP in several ways. Psychosocial factors related to night work, such as low job control and work-life conflict, have been associated with cardiovascular diseases including hypertension (20, 21). Another mechanism is through behavioral changes induced by night work affecting sleep, smoking

¹ Department of Occupational and Environmental Medicine, Bispebjerg University Hospital, Copenhagen, Denmark.

² The Parker Institute, Bispebjerg and Frederiksberg University Hospital, Copenhagen, Denmark.

³ Department of Obstetrics and Gynecology, Hvidovre University Hospital, Copenhagen, Denmark.

⁴ National Research Centre for the Working Environment, Copenhagen, Denmark.

⁵ University of Copenhagen, Department of Public Health, Copenhagen, Denmark.

⁶ Danish Cancer Society Research Center, Copenhagen, Denmark.

⁷ Department of Occupational Medicine, Danish Ramazzini Centre, Aarhus University Hospital, Aarhus, Denmark.

Correspondence to: Paula EC Hammer, Bispebjerg Bakke 23, indgang 20F, 2400 Copenhagen, Denmark. [E-mail: paulahammer@dadlnet.dk]

habits, physical activity, diet and body mass (20, 22). Furthermore several physiological mechanisms including circadian disruption, hormonal changes, altered lipids and increased inflammation markers have been proposed linking night work with cardiovascular diseases (20, 22, 23). Melatonin, one of the main hormones affected by circadian disruption, is also produced in the placenta and plays a crucial role in maternal, fetal and placental physiology acting as an anti-inflammatory and immunomodulatory hormone, as well as a regulator of apoptosis (24–32). Furthermore the circadian oscillation of blood pressure is controlled in part by melatonin (33, 34). An altered circadian pattern of blood pressure has been reported in HDP, and as a result melatonin has been studied for its potential use in the treatment of preeclampsia (35, 36).

The few studies that have been conducted on the association between night and shift work with HDP revealed conflicting results (37–40). A major limitation of these studies is the crude assessment of work schedules. For instance in three (37, 38, 40) out of four studies it was not clear whether their definition of shift work included night shifts.

Payroll data provides accurate information on work schedules for a large population overcoming hereby the limitations related to exposure assessment in prior studies (41, 42).

The primary aim of this study was to investigate whether night work expressed by number and duration of night shifts, number of consecutive night shifts and number of quick returns during pregnancy is related to increased risk of HDP. We furthermore investigated whether age, body mass index (BMI) and socioeconomic status (SES) modified the effect of night work on the risk of HDP.

Methods

Design

We conducted a prospective register-based cohort study with information from three Danish national registries linked on individual level through the civil registration number given to all residents in Denmark since 1968.

The Danish Working Hour Database (DWHD), a national payroll database covering more than 250 000 employees in the Danish administrative regions including all hospital employees, provided the source population. It includes daily information on time of start and end of all workdays, sickness absence, paid and unpaid leave, occupation and place of employment from January 2007 to December 2015 (41, 43). Pregnancy information and covariates were identified from the Danish

Medical Birth Registry, which contains information from all home and hospital births in Denmark from 1973 onwards (44). Outcome variables were identified from the Danish National Patient Registry, which provides data on inpatients in Danish hospitals since 1977 and on outpatients since 1994 (45).

Study cohort

Women from DWHD who gave birth at least once between 2007 and 2013 were identified (N=42 485 women with 60 482 births). We excluded women ≤ 18 and ≥ 50 years (N=15), multiple pregnancies (N=2957), pregnancies conceived in 2006 (N=6403), and pregnancies from women without registrations in DWHD of any night or day shift during the first 20 pregnancy weeks (N=26 481). This was the exposure time because gestational hypertension is, by definition, diagnosed after 20 pregnancy weeks (14). To avoid clustering effects, each woman contributed with only one pregnancy, the first during the study period (N=5902 pregnancies excluded), leaving 18 724 women eligible for analyses (figure 1).

Exposure

Exposure definitions are in line with recent studies using payroll data (41, 46).


Shifts, including on-call shifts, lasting ≥ 3 hours were defined as day (start time after 06:00 and end time before 21:00 hours), evening (end time after 21:00 and before 02:00 hours), night (any start and end time including working hours between 23:00 and 06:00 hours) and early morning (start time between 03:00 and 06:00 hours).

A night worker was defined by working ≥ 1 night shift and a day worker by working ≥ 1 day shift but no night, evening or early morning shifts during the first 20 pregnancy weeks.

Consecutive night shifts. Categories of consecutive night shifts were 0 (only single night shifts), 2–3 (at least one spell of 2–3 consecutive night shifts and no spells of ≥ 4 consecutive night shifts), and ≥ 4 (at least one spell of ≥ 4 consecutive night shifts) during the first 20 pregnancy weeks.

Quick returns. We defined quick returns as intervals between shifts lasting < 11 hours (47). Quick returns after night shifts were defined as a recovery period of < 28 hours after a night shift (46). Categories of number of quick returns and quick returns after night shift were 0, 1–4 and ≥ 5 quick returns during the first 20 pregnancy weeks.

Duration of night shifts. Long night shift workers were defined by working ≥ 1 long night shift (≥ 12 hours) during the first 20 pregnancy weeks.



Number of night shifts. Number of night shifts was analyzed in categories of 1–19 or ≥ 20 (roughly corresponding to ≥ 1 night shift/week during the first 20 pregnancy weeks).

Covariates

Age (<30, 30–35, >35 years), BMI (<18.5, 18.5–24.9, 25–29.9, ≥ 30 kg/m²), parity (1, 2, ≥ 3) and smoking (nonsmoker, former smoker, smoker) registered by the midwife or family doctor at the first antenatal visit were retrieved from the Danish Medical Birth Registry. Classification of SES into high, low or medium was derived from Statistics Denmark. It was based on DISCO-88, the Danish version of the International Standard Classification of Occupations (ISCO-88) (48), in the calendar years 2007–2009, and DISCO-08, the Danish version of ISCO-08 (49), in the calendar years 2010–2013.

Sickness absence three months prior to pregnancy was expressed as the sum of all days registered with ≥ 3 hours of sickness absence in DWHD during this period. It was categorized as 0, <10 and ≥ 10 days.

Missing values for parity, smoking and SES represented only 1.4%, 2.9% and 0.2% respectively. Missing

values for BMI (4.4%) were evenly distributed across exposure categories. Missing values of sickness absence three months prior to conception (7.9%) occurred when the woman's employment covered by DWHD had <3 months prior to conception.

Outcome

The outcome of HDP was defined by ICD-10 codes (50): hypertension (I10–I15), gestational hypertension (O12, I13, I16) and pre-eclampsia and eclampsia (O14, I15).

Statistical analysis

We computed odds ratios (OR) with 95% confidence intervals (CI) for HDP according to different dimensions of night work during the first 20 weeks of pregnancy by logistic regression. Model 1 refers to crude analyses and model 2 is adjusted for age, BMI, parity, smoking, SES and sickness absence three months prior to pregnancy categorized as described above. Because of too few cases, it was not possible to adjust the analyses for cases of prior HDP (N=287), prior diabetes (N=17), and current

gestational diabetes (N=202). Model 3 is further adjusted for number of night shifts in the analyses of consecutive night shifts, quick returns and duration of night shifts.

In all analyses, except for interaction analyses, we made comparisons of night workers with day workers and comparisons within night workers. In the latter, night workers in the lowest category of exposure (1–19 night shifts, duration of night shift of <12 hours, night workers without consecutive night shifts and night workers without quick returns) were used as the reference group.

We investigated whether the association between night work and HDP was modified by age, BMI and SES by a likelihood ratio test comparing models with main effects only with models that in addition included an interaction term, ie, the product of the combined effect. We used a level of significance of 5%.

Gestational length was used to identify conception date. There were only 330 (0.6%) pregnancies with missing values for gestational length but the proportion of still births among these was statistically significant higher (15.4%) than among other pregnancies (0.4%). We therefore substituted the missing values by the mean value of gestational length for live (278 days) and still

(220 days) births, respectively.

We performed the following sensitivity analyses: (i) restricted to nulliparous women (N=9 660), (ii) with pre-eclampsia as the outcome (N=18 724), and (iii) restricted to the first trimester as the exposure time (N=18 158). In the latter analysis, night workers had at ≥1 night shift and day workers ≥1 day shift but no night, evening or early morning shifts during the first 12 pregnancy weeks instead of 20 weeks applied in the main analysis.

All analyses were done with the SAS 9.4 software (SAS Institute, Cary, North Carolina, United States).

Results

In our cohort of 18 724 pregnant women, 11 193 were classified as night workers and 7531 as day workers (table 1). The most frequent occupations were nurse (44%), physician (13%), medical secretary (7%), physio/occupational therapist (5%) and laboratory technician (4%) reflecting that the majority of the workers covered by the DWHD are employed at hospitals. Characteristics of day

Table 1. Characteristics of pregnant workers in public administration and hospitals in Denmark, 2007–2013. [SD=standard deviation.]

	Day work ^a (N=7531)				Night work ^b (N=11 193)			
	N	%	Mean	SD	N	%	Mean	SD
Age (years)			31.8	4.2			30.7	3.9
Body mass index (kg/m ²)			24.0	7.6			23.9	7.9
Parity								
1	3462	46.0			6198	55.4		
2	2596	34.5			3156	28.2		
≥3	1356	18.0			1701	15.2		
Smoking								
Non smoker	6952	92.3			10 319	92.2		
Former smoker	106	1.4			230	2.1		
Smoker	273	3.6			309	2.8		
Socioeconomic status								
High	1844	24.5			2618	23.4		
Medium	3848	51.1			7575	67.7		
Low	1821	24.2			988	8.8		
Most frequent ^c occupations								
Nurse	1071	14.2			6,857	61.3		
Physician	659	8.8			2,000	17.9		
Nurse assistant	208	2.8			390	3.5		
Laboratory technician	503	6.7			259	2.3		
Midwife	10	0.1			248	2.2		
Medical secretary	1314	17.5			58	0.5		
Physio- and ergo therapist	977	13.0			30	0.3		
Cleaning and kitchen staff	391	5.2			8	0.07		
Psychologist	463	6.2			1	0.01		
Shifts during the first 20 weeks of pregnancy								
Day			65.9	23.8			37.9	17.2
Night							11.2	9.1
Evening							9.2	9.7
Early morning							0.03	0.8
Weekly working hours ^d			23.8	8.9			25.1	7.7
Sickness absence days 3 months prior to pregnancy			2.9	7.2			2.6	5.6

^a ≥1 day shift and no night, evening or early morning shift during the first 20 pregnancy weeks.

^b ≥1 night shift during the first 20 pregnancy weeks.

^c Occupations with ≥100 subjects.

^d Paid and unpaid leave excluded.

and night workers were rather similar. Day workers had a total of 496 024 day shifts during the first 20 weeks of pregnancy. Night workers had a total of 652 858 shifts being 65% day shifts and 19% night shifts. Only 113 women (1%) worked fixed night shifts. They had higher BMI (mean 25.2 kg/m², SD 5.0), higher proportion of women with parity ≥ 3 (27%), higher proportion of current smokers (4.4%) and higher proportion of women with low SES (43%) compared to the other night workers in

the cohort. The prevalence of HDP was 3.6% among day workers and 3.8% among night workers.

Women working ≥ 1 spell of ≥ 4 consecutive night shifts during the first 20 pregnancy weeks had higher risk of HDP compared to night workers without consecutive night shifts (OR 1.41, 95% CI 1.01–1.98), see table 2. We furthermore observed a statistically significant trend of increasing risk with increasing number of consecutive night shifts. Women with spells of exclu-

Table 2. Odds ratios (OR) of hypertensive disorders of pregnancy by consecutive night shifts during the first 20 pregnancy weeks among workers in public administration and hospitals in Denmark, 2007–2013. [CI=confidence interval]

Consecutive night shifts	Women		Cases		Model 1 ^a		Model 2 ^b	
	N	%	N	%	OR	95% CI	OR	95% CI
All workers (N=18 724)								
Day work ^c	7531	40.2	270	3.6	1.00	Referent	1.00	Referent
0	4003	21.4	132	3.3	0.92	0.74–1.13	0.85	0.67–1.08
2–3	5225	27.9	205	3.9	1.10	0.91–1.32	0.97	0.79–1.20
≥ 4	1965	10.5	89	4.5	1.28	0.99–1.62	1.13	0.86–1.48
P for trend						0.05		0.62
Night workers ^d (N=11 193)								
0	4003	35.8	132	3.3	1.00	Referent	1.00	Referent
2–3	5225	46.7	205	3.9	1.20	0.96–1.50	1.22	0.92–1.62
≥ 4	1965	17.6	89	4.5	1.39	1.06–1.83	1.41	1.01–1.98
P for trend						0.02		0.04

^a Crude analysis.

^b Adjusted for categories of age, body mass index, smoking, socioeconomic status, parity and sickness absence three months prior to pregnancy.

^c ≥ 1 day shift and no night, evening or early morning shift during the first 20 pregnancy weeks.

^d ≥ 1 night shift during the first 20 pregnancy weeks.

Table 3. Odds ratios (OR) of hypertensive disorders of pregnancy by number of quick returns^a and quick returns after a night shift^b during the first 20 pregnancy weeks among workers in public administration and hospitals in Denmark, 2007–2013. [CI=confidence interval]

	Women		Cases		Model 1 ^c		Model 2 ^d	
	N	%	N	%	OR	95% CI	OR	95% CI
Quick returns								
All workers (N=18 724)								
Day work ^e	7531	40.2	270	3.6	1.00	Referent	1.00	Referent
0	3817	20.4	128	3.4	0.93	0.75–1.15	0.92	0.72–1.16
1–4	5123	27.4	203	4.0	1.11	0.92–1.34	1.00	0.81–1.23
≥ 5	2253	12.0	95	4.2	1.18	0.93–1.50	0.94	0.72–1.22
P for trend					0.10		0.76	
Night workers ^f (N=11 193)								
0	3817	34.1	128	3.4	1.00	Referent	1.00	Referent
1–4	5123	45.8	203	4.0	1.19	0.95–1.49	1.12	0.87–1.45
≥ 5	2253	20.1	95	4.2	1.27	0.97–1.66	1.07	0.79–1.46
P for trend					0.07		0.64	
Quick returns after a night shift								
All workers (N=18 724)								
Day work ^e	7531	40.2	270	3.6	1.00	Referent	1.00	Referent
0	1023	5.5	39	3.8	1.07	0.75–1.48	0.84	0.55–1.23
1–4	4569	24.4	160	3.5	0.98	0.80–1.19	0.86	0.69–1.07
≥ 5	5601	29.9	227	4.1	1.14	0.95–1.36	1.06	0.87–1.29
P for trend					0.26		0.74	
Night workers ^f (N=11 193)								
0	1023	9.1	39	3.8	1.00	Referent	1.00	Referent
1–4	4569	40.8	160	3.5	0.92	0.65–1.33	1.03	0.69–1.59
≥ 5	5601	50.0	227	4.1	1.07	0.76–1.53	1.28	0.87–1.95
P for trend					0.30		0.05	

^a < 11 hours between two consecutive shifts.

^b < 28 hours between a night shift and the consecutive shift.

^c Crude analysis.

^d Adjusted for categories of age, body mass index, smoking, socioeconomic status, parity and sickness absence three months prior to pregnancy.

^e ≥ 1 day shift and no night, evening or early morning shift during the first 20 pregnancy weeks.

^f ≥ 1 night shift during the first 20 pregnancy weeks.

Table 4. Odds ratios (OR) of hypertensive disorders of pregnancy by duration of night shifts during the first 20 pregnancy weeks among workers in public administration and hospitals in Denmark, 2007–2013. [CI=confidence interval]

Duration of night shifts	Women		Cases		Model 1 ^a		Model 2 ^b	
	N	%	N	%	OR	95% CI	OR	95% CI
All workers (N=18 724)								
Day work ^c	7531	40.2	270	3.6	1.00	Referent	1.00	Referent
<12 hours ^d	5734	30.6	214	3.7	1.04	0.87–1.25	0.94	0.76–1.16
≥12 hours ^e	5459	29.2	212	3.9	1.09	0.90–1.31	1.00	0.81–1.23
P for trend					0.37		0.92	
Night workers ^f (N=11 193)								
<12 hours	5734	51.2	214	3.7	1.00	Referent	1.00	Referent
≥12 hours	5459	48.8	212	3.9	1.04	0.86–1.27	1.08	0.85–1.36

^a Crude analysis.^b Adjusted for categories of age, body mass index, smoking, socioeconomic status, parity and sickness absence three months prior to pregnancy.^c ≥1 day shift and no night, evening or early morning shift during the first 20 pregnancy weeks.^d ≥1 night of <12 hours and no night shifts of ≥12 hours.^e ≥1 night shift of ≥12 hours.^f ≥1 night shift during the first 20 pregnancy weeks.**Table 5.** Odds ratios (OR) of hypertensive disorders of pregnancy by number of night shifts during the first 20 pregnancy weeks among workers in public administration and hospitals in Denmark, 2007–2013. [CI=confidence interval]

Number of night shifts	Women		Cases		Model 1 ^a		Model 2 ^b	
	N	%	N	%	OR	95% CI	OR	95% CI
All workers, N=18 724								
Day work ^c	7531	40.2	270	3.6	1.00	Referent	1.00	Referent
1–19	9560	51.1	360	3.8	1.05	0.90–1.24	0.94	0.78–1.12
≥20	1633	8.7	66	4.0	1.13	0.85–1.48	1.09	0.81–1.45
P for trend					0.35		0.96	
Night workers ^d , N=11 193								
1–19	9560	85.4	360	3.8	1.00	Referent	1.00	Referent
≥20	1633	14.6	66	4.0	1.08	0.82–1.40	1.15	0.86–1.52

^a Crude analysis.^b Adjusted for categories of age, body mass index, smoking, socioeconomic status, parity and sickness absence three months prior to pregnancy.^c ≥1 day shift and no night, evening or early morning shift during the first 20 pregnancy weeks.^d ≥1 night shift during the first 20 pregnancy weeks.

sively 2–3 consecutive night shifts had, on average, 4 consecutive night shifts in total. While women with ≥1 spell of ≥4 consecutive night shifts had, on average, 14 consecutive night shifts in total. Hence these categories express both length of spells and total number of consecutive night shifts.

As shown in table 3, we observed a statistically significant trend of increasing risk of HDP with increasing number of quick returns after night shifts. However, the risk estimate for the highest exposed group, those with ≥5 quick returns after a night shift (on average 10.4 quick returns) during the first 20 pregnancy weeks, did not reach statistical significance (OR 1.28, 95% CI 0.87–1.95).

Table 4 presents the results for long-night-shift workers compared to day workers (OR 1.00, 95% CI 0.81–1.23), and compared to short-night-shift workers (OR 1.08, 95% CI 0.85–1.36). Of all long night shifts, 40% lasted 17–24 hours and 34% lasted 9–16 hours, while 62% of all short night shifts lasted ≤8 hours.

Table 5 presents the results for women who worked ≥20 night shifts during the first 20 pregnancy weeks (on average 28 night shifts) compared to day workers (OR 1.13, 95% CI 0.85–1.48), and compared to women

working 1–19 night shifts (OR 1.15, 95% CI 0.86–1.52).

Further adjustment for number of night shifts (model 3) did not substantially change the results in the analyses of consecutive night shifts, quick returns and duration of night shifts.

The association between night work and HDP was modified by BMI (P-value for multiplicative interaction 0.03). As presented in table 6, analysis among women with BMI ≥30 kg/m² revealed that those who worked ≥4 consecutive night shifts had substantially increased risk of HDP compared to day workers (OR 5.31, 95% CI 1.98–14.22). The corresponding risk for women with BMI <25 kg/m² was OR 1.02, 95% CI 0.73–1.41. Further adjustment for BMI among obese women did not change the results. A similar increase was observed for all exposures among obese women (see supplementary tables S1–S4, www.sjweh.fi/show_abstract.php?abstract_id=3728). Due to low statistical power we were unable to make stratified comparisons within night workers only. We found no interaction of any of the analyzed exposures with maternal age or SES.

Overall sensitivity analyses slightly attenuated the estimates across all exposures. The effect of consecutive night shifts during the first 20 pregnancy weeks was con-

Table 6. Odds ratios (OR) of hypertensive disorders of pregnancy by consecutive night shifts during the first 20 pregnancy weeks stratified^a by body mass index (BMI) among workers in public administration and hospitals in Denmark, 2007–2013. [CI=confidence interval]

Consecutive night shifts	Women		Cases		Model1 ^b		Model2 ^c	
	N	%	N	%	OR	95% CI	OR	95% CI
BMI <25 kg/m ² (N=12 815)								
Day work	5119	40.0	201	3.9	1.00	Referent	1.00	Referent
0	2952	23.0	96	3.3	0.82	0.64–1.05	0.77	0.59–1.02
2–3	3545	27.7	154	4.3	1.11	0.90–1.38	0.95	0.75–1.21
≥4	1199	9.4	56	4.7	1.20	0.89–1.62	1.02	0.73–1.41
BMI 25–29 kg/m ² (N=3501)								
Day work	1419	40.5	52	3.7	1.00	Referent	1.00	Referent
0	644	18.4	24	3.7	1.02	0.62–1.67	0.93	0.55–1.56
2–3	994	28.4	40	4.0	1.10	0.72–1.68	0.97	0.63–1.51
≥4	444	12.7	16	3.6	0.98	0.56–1.74	0.94	0.53–1.69
BMI ≥ 30 kg/m ² (N=1588)								
Day work	671	42.3	7	1.0	1.00	Referent	1.00	Referent
0	229	14.4	10	4.4	4.33	1.63–11.52	3.47	1.15–10.52
2–3	435	27.4	8	1.8	1.78	0.64–4.94	1.60	0.53–4.83
≥4	253	15.9	13	5.1	5.14	2.03–13.03	5.31	1.98–14.22

^a P-value for multiplicative interaction 0.0317.^b Crude analysis.^c Adjusted for categories of age, smoking, socioeconomic status, parity and sickness absence three months prior to pregnancy.

sistent although not statistically significant throughout sensitivity analyses within night workers (OR 1.39, 95% CI 0.94–2.05 restricted to nulliparous women, OR 1.40, 95% CI 0.91–2.15 with pre-eclampsia as the outcome, and OR 1.36, 95% CI 0.96–1.93 with the first trimester as the exposure time). Regarding the question on possible selection out of night work during pregnancy, we identified only 580 women (5%) who worked at least one night shift during the first trimester and changed to fixed day work during the second trimester. These women had similar age (mean 31 years), BMI (mean 23.7 kg/m²) and smoking habits (2.8% current smokers) as the rest of the cohort but presented a higher proportion of physicians (37%).

Discussion

To our knowledge, this is the first study to investigate the association between HDP with different dimensions of night work objectively assessed through payroll data. In our study, workers with ≥4 consecutive night shifts during the first 20 pregnancy weeks had higher risk of HDP compared to night workers without consecutive night shifts (OR 1.41, 95% CI 1.01–1.98). We furthermore observed a dose–response gradient for number of consecutive night shifts and the risk of HDP. The fact that this effect was observed in comparisons within night workers strengthens the evidence of a causal effect as the group of night workers is more homogeneous. These analyses may therefore be less susceptible to the healthy worker effect present in comparisons of night versus day workers. In fact, we observed higher risk estimates in comparisons within night workers for all the expo-

sure. Comparisons within night workers may be more appropriate from an epidemiological point of view. On the other hand, analyses restricted to night workers exclude an unexposed group and some selection bias regarding different dimensions of night work remains. Previous studies have shown that individual preferences related to both personal (chronotype, sleep flexibility, social context) (51–53) and occupational (work content, demands and environment) (54) factors vary substantially among night workers resulting in differences in adaptation to night work. Accordingly we found that workers with fixed night work during the first 20 pregnancy weeks differed in BMI, parity, smoking habits and SES compared to the other night workers in the cohort. Compared to the background Danish population, our cohort presented lower prevalence of smoking during pregnancy (3% versus 12%) (55) and lower proportion of overweight women (19% versus 46%) (56), which may reflect a more health promoting behavior among healthcare professionals.

Our findings are in accordance with recent studies focusing on consecutive night shifts rather than solely on the number of night shifts. For example increasing the number of consecutive night shifts has been associated with progressive changes in hormones involved in circadian regulation, such as melatonin, cortisol, thyroxine and prolactin (30, 31, 57). Such changes have been observed down to three consecutive night shifts (58, 59). Furthermore, it has been suggested that at least two days off work are required to allow for circadian readjustment following 2–4 consecutive night shifts (31, 60). In our cohort, the majority of hospital employees had rotating shifts with different schedules nearly every week which do not fulfill this recommendation. Hence, in this context, working consecutive night shifts may

lead both to circadian disruption and to insufficient recovery. Our findings of increasing odds ratios of HDP with increasing number of quick returns after night shifts also support the potential effect of insufficient recovery after a night shift.

In our data, BMI modified the effect of night work on the risk of HDP, as obese women who worked longer night shifts, longer spells of consecutive night shifts and had the highest number of quick returns after night shifts had 4–5 fold increased risk of HDP compared to day workers. It is known that pre-pregnancy BMI is an important risk factor for HDP independent of weight gain during pregnancy (17, 61, 62). Even though these results are based on few cases, they are consistent across exposures. Obese women neither had higher proportion of workers with fixed night shifts nor a gradient of increasing BMI from day to night workers.

We hypothesized that women who worked night shifts during the first trimester and changed working schedule to only day work during the second trimester due to health problems might cause bias towards the null as the exposure time in the main analysis was 20 weeks. However, sensitivity analysis resetting exposure time to the first 12 pregnancy weeks indicated no such bias. On the other hand, analysis restricted to the first trimester excludes a possible effect of night work during the second trimester, which may in part explain the attenuation of the estimates. Even though the physiopathology of HDP seems to be related with placenta development in the beginning of pregnancy (14), demographic and lifestyle factors on the second and third trimester of pregnancy seem also to influence the risk of HDP (63).

We found no statistically significant association between HDP with any of the analyzed dimensions of night work compared to day workers, suggesting that the effect of night work on the risk of HDP is related to the way night shifts are organized rather than the mere presence of night shifts. This can be in part due to differences in work content and work environment between day and night workers. We did not observe the presence of more pronounced risk factors for HDP among night workers compared to day workers. Actually our cohort of night workers had lower BMI, a lower proportion of smokers and a lower proportion of workers with low SES than day workers. Similar to our results, three out of four previous studies that compared shift workers with day workers found no association between with HDP (37, 39, 40). Wergeland & Strand (38) reported an increased prevalence of pre-eclampsia among shift workers, but only among parous women.

The main strengths of our study are the large and national sample size, the objective and detailed exposure assessment, and the use of validated and objective registries for identification of covariates and outcomes, which makes information bias and selection in and out

of the study unlikely. Furthermore, we evaluated different dimensions of night work within night workers and restricted the exposure time to specific periods of pregnancy. Some limitations include a lack of information on workload during night shifts, such as the possibility for sleep during on call shifts, and on chronotype and personal preferences of the participants. The latter is especially relevant because night work is compulsory for the majority of occupations in our cohort. Additionally, our study design did not account for the healthy worker effect, where women with health problems in general tend to choose day work. As our cohort comprises primarily healthcare professionals, our results may not apply for pregnant workers in other occupations.

Ideally future studies on health effects of night work during pregnancy should combine objectively assessed work schedules with information on chronotype and personal preferences, work content and environment, and should perform comparisons both with day workers and within groups of night workers.

Concluding remarks

In this nationwide study of Danish pregnant workers in the public health sector with objectively assessed work schedules, working consecutive night shifts and quick returns after night shifts during the first 20 pregnancy weeks was associated with an increased risk of HDP, in particular among obese women. Possible ways for avoiding such risk when organizing night work during pregnancy are favoring single night shifts or short spells of consecutive night shifts and reducing quick returns by allowing for adequate recovery time following night shifts.

Acknowledgments

This work was supported by The Danish Working Environment Research Fund grant 31-2015-03 2015001705.

The establishment of the DWHD has been financed by research grants from The Danish Working Environment Research Fund (23-2012-09), The Nordic Program on Health and Welfare – Nordforsk (74809) and The National Research Centre for the Working Environment. The Danish administrative regions have partially financed the transfer of data to the cohort.

The Danish administrative regions are acknowledged for the participation and willingness to provide data to the DWHD. Jens Worm Begtrup, Lisbeth Nielsen and Anders Ørberg are thanked for valuable work with data management.

The authors declare no conflicts of interest.

References

- Parent-Thirion A, Biletta I, Cabrita J, Vargas O, Vermeulen G, Wilczynska A et al. Eurofond (2016), Sixth European Working Conditions Survey - Overview report. Luxembourg; 2016.
- Bonde JP, Jørgensen KT, Bonzini M, Palmer KT. Miscarriage and occupational activity: a systematic review and meta-analysis regarding shift work, working hours, lifting, standing, and physical workload. *Scand J Work Environ Health* 2013 Jul;39(4):325–34. <http://dx.doi.org/10.5271/sjweh.3337>.
- Stocker LJ, Macklon NS, Cheong YC, Bewley SJ. Influence of shift work on early reproductive outcomes: a systematic review and meta-analysis. *Obstet Gynecol* 2014 Jul;124(1):99–110. <http://dx.doi.org/10.1097/AOG.0000000000000321>.
- Zhu JL, Hjollund NH, Andersen AM, Olsen J. Shift work, job stress, and late fetal loss: The National Birth Cohort in Denmark. *J Occup Environ Med* 2004 Nov;46(11):1144–9. <http://dx.doi.org/10.1097/01.jom.0000145168.21614.21>.
- Zhu JL, Hjollund NH, Boggild H, Olsen J. Shift work and subfecundity: a causal link or an artefact? *Occup Environ Med* 2003 Sep;60(9):E12. <http://dx.doi.org/10.1136/oem.60.9.e12>.
- Zhu JL, Hjollund NH, Olsen J; National Birth Cohort in Denmark. Shift work, duration of pregnancy, and birth weight: the National Birth Cohort in Denmark. *Am J Obstet Gynecol* 2004 Jul;191(1):285–91. <http://dx.doi.org/10.1016/j.ajog.2003.12.002>.
- Bonzini M, Palmer KT, Coggon D, Carugno M, Cromi A, Ferrario MM. Shift work and pregnancy outcomes: a systematic review with meta-analysis of currently available epidemiological studies. *BJOG* 2011 Nov;118(12):1429–37. <http://dx.doi.org/10.1111/j.1471-0528.2011.03066.x>.
- Palmer KT, Bonzini M, Harris EC, Linaker C, Bonde JP. Work activities and risk of prematurity, low birth weight and pre-eclampsia: an updated review with meta-analysis. *Occup Environ Med* 2013 Apr;70(4):213–22. <http://dx.doi.org/10.1136/oemed-2012-101032>.
- Chau YM, West S, Mapedzahama V. Night work and the reproductive health of women: an integrated literature review. *J Midwifery Womens Health* 2014 Mar-Apr;59(2):113–26. <http://dx.doi.org/10.1111/jmwh.12052>.
- Kuklina EV, Ayala C, Callaghan WM. Hypertensive disorders and severe obstetric morbidity in the United States. *Obstet Gynecol* 2009 Jun;113(6):1299–306. <http://dx.doi.org/10.1097/AOG.0b013e3181a45b25>.
- Duley L. The global impact of pre-eclampsia and eclampsia. *Semin Perinatol* 2009 Jun;33(3):130–7. <http://dx.doi.org/10.1053/j.semperi.2009.02.010>.
- Behrens I, Basit S, Melbye M, Lykke JA, Wohlfahrt J, Bundgaard H et al. Risk of post-pregnancy hypertension in women with a history of hypertensive disorders of pregnancy: nationwide cohort study. *BMJ* 2017 Jul;358:j3078. <http://dx.doi.org/10.1136/bmj.j3078>.
- Wallis AB, Safflas AF, Hsia J, Atrash HK. Secular trends in the rates of preeclampsia, eclampsia, and gestational hypertension, United States, 1987–2004. *Am J Hypertens* 2008 May;21(5):521–6. <http://dx.doi.org/10.1038/ajh.2008.20>.
- Naderi S, Tsai SA, Khandelwal A. Hypertensive Disorders of Pregnancy. *Curr Atheroscler Rep* 2017 Mar;19(3):15–0648. <http://dx.doi.org/10.1007/s11883-017-0648-z>.
- Sibai B, Dekker G, Kupferminc M. Pre-eclampsia. *Lancet* 2005 Feb;365(9461):785–99. [http://dx.doi.org/10.1016/S0140-6736\(05\)71003-5](http://dx.doi.org/10.1016/S0140-6736(05)71003-5).
- Gaillard R, Bakker R, Steegers EA, Hofman A, Jaddoe VW. Maternal age during pregnancy is associated with third trimester blood pressure level: the generation R study. *Am J Hypertens* 2011 Sep;24(9):1046–53. <http://dx.doi.org/10.1038/ajh.2011.95>.
- Gaillard R, Steegers EA, Hofman A, Jaddoe VW. Associations of maternal obesity with blood pressure and the risks of gestational hypertensive disorders. The Generation R Study. *J Hypertens* 2011 May;29(5):937–44. <http://dx.doi.org/10.1097/HJH.0b013e328345500c>.
- Zhou A, Xiong C, Hu R, Zhang Y, Bassig BA, Triche E et al. Pre-Pregnancy BMI, Gestational Weight Gain, and the Risk of Hypertensive Disorders of Pregnancy: A Cohort Study in Wuhan, China. *PLoS One* 2015 Aug;10(8):e0136291. <http://dx.doi.org/10.1371/journal.pone.0136291>.
- Silva LM, Coolman M, Steegers EA, Jaddoe VW, Moll HA, Hofman A et al. Low socioeconomic status is a risk factor for preeclampsia: the Generation R Study. *J Hypertens* 2008 Jun;26(6):1200–8. <http://dx.doi.org/10.1097/HJH.0b013e3282fcc36e>.
- Puttonen S, Härmä M, Hublin C. Shift work and cardiovascular disease - pathways from circadian stress to morbidity. *Scand J Work Environ Health* 2010 Mar;36(2):96–108. <http://dx.doi.org/10.5271/sjweh.2894>.
- Härmä M. Workhours in relation to work stress, recovery and health. *Scand J Work Environ Health* 2006 Dec;32(6):502–14. <http://dx.doi.org/10.5271/sjweh.1055>.
- Kecklund G, Axelsson J. Health consequences of shift work and insufficient sleep. *BMJ* 2016 Nov;355:i5210. <http://dx.doi.org/10.1136/bmj.i5210>.
- Vyas MV, Garg AX, Iansavichus AV, Costella J, Donner A, Laugsand LE et al. Shift work and vascular events: systematic review and meta-analysis. *BMJ* 2012 Jul;345:e4800. <http://dx.doi.org/10.1136/bmj.e4800>.
- Nakamura Y, Tamura H, Kashida S, Takayama H, Yamagata Y, Karube A et al. Changes of serum melatonin level and its relationship to feto-placental unit during pregnancy. *J Pineal Res* 2001 Jan;30(1):29–33. <http://dx.doi.org/10.1034/j.1600-079X.2001.300104.x>.
- Reiter RJ, Tan DX, Korkmaz A, Rosales-Corral SA. Melatonin and stable circadian rhythms optimize maternal, placental and fetal physiology. *Hum Reprod Update* 2014 Mar-Apr;20(2):293–307. <http://dx.doi.org/10.1093/humupd/dmt054>.

26. Reiter RJ, Tan DX, Tamura H, Cruz MH, Fuentes-Broto L. Clinical relevance of melatonin in ovarian and placental physiology: a review. *Gynecol Endocrinol* 2014 Feb;30(2):83–9. <http://dx.doi.org/10.3109/09513590.2013.849238>.
27. Soliman A, Lacasse AA, Lanoix D, Sagrillo-Fagundes L, Boulard V, Vaillancourt C. Placental melatonin system is present throughout pregnancy and regulates villous trophoblast differentiation. *J Pineal Res* 2015 Aug;59(1):38–46. <http://dx.doi.org/10.1111/jpi.12236>.
28. Tamura H, Nakamura Y, Terron MP, Flores LJ, Manchester LC, Tan DX et al. Melatonin and pregnancy in the human. *Reprod Toxicol* 2008 Apr;25(3):291–303. <http://dx.doi.org/10.1016/j.reprotox.2008.03.005>.
29. Boivin DB, Boudreau P. Impacts of shift work on sleep and circadian rhythms. *Pathol Biol (Paris)* 2014 Oct;62(5):292–301. <http://dx.doi.org/10.1016/j.patbio.2014.08.001>.
30. Jensen MA, Garde AH, Kristiansen J, Nabe-Nielsen K, Hansen AM. The effect of the number of consecutive night shifts on diurnal rhythms in cortisol, melatonin and heart rate variability (HRV): a systematic review of field studies. *Int Arch Occup Environ Health* 2016 May;89(4):531–45. <http://dx.doi.org/10.1007/s00420-015-1093-3>.
31. Jensen MA, Hansen AM, Kristiansen J, Nabe-Nielsen K, Garde AH. Changes in the diurnal rhythms of cortisol, melatonin, and testosterone after 2, 4, and 7 consecutive night shifts in male police officers. *Chronobiol Int* 2016 Aug;11:1–13. <https://doi.org/10.1080/07420528.2016.1212869>.
32. Morris CJ, Aeschbach D, Scheer FA. Circadian system, sleep and endocrinology. *Mol Cell Endocrinol* 2012 Feb;349(1):91–104. <http://dx.doi.org/10.1016/j.mce.2011.09.003>.
33. Simko F, Pechanova O. Potential roles of melatonin and chronotherapy among the new trends in hypertension treatment. *J Pineal Res* 2009 Sep;47(2):127–33. <http://dx.doi.org/10.1111/j.1600-079X.2009.00697.x>.
34. Solocinski K, Gumz ML. The Circadian Clock in the Regulation of Renal Rhythms. *J Biol Rhythms* 2015 Dec;30(6):470–86. <http://dx.doi.org/10.1177/0748730415610879>.
35. Marseglia L, D'Angelo G, Manti S, Reiter RJ, Gitto E. Potential utility of melatonin in preeclampsia, intrauterine fetal growth retardation, and perinatal asphyxia. *Reprod Sci* 2016 Aug;23(8):970–7. <http://dx.doi.org/10.1177/1933719115612132>.
36. Aversa S, Pellegrino S, Barberi I, Reiter RJ, Gitto E. Potential utility of melatonin as an antioxidant during pregnancy and in the perinatal period. *J Matern Fetal Neonatal Med* 2012 Mar;25(3):207–21. <http://dx.doi.org/10.3109/14767058.2011.573827>.
37. Nurminen T. Shift work, fetal development and course of pregnancy. *Scand J Work Environ Health* 1989 Dec;15(6):395–403. <http://dx.doi.org/10.5271/sjweh.1833>.
38. Wergeland E, Strand K. Working conditions and prevalence of pre-eclampsia, Norway 1989. *Int J Gynaecol Obstet* 1997 Aug;58(2):189–96. [http://dx.doi.org/10.1016/S0020-7292\(97\)00083-0](http://dx.doi.org/10.1016/S0020-7292(97)00083-0).
39. Haelterman E, Marcoux S, Croteau A, Dramaix M. Population-based study on occupational risk factors for preeclampsia and gestational hypertension. *Scand J Work Environ Health* 2007 Aug;33(4):304–17. <http://dx.doi.org/10.5271/sjweh.1147>.
40. Chang PJ, Chu LC, Hsieh WS, Chuang YL, Lin SJ, Chen PC. Working hours and risk of gestational hypertension and pre-eclampsia. *Occup Med (Lond)* 2010 Jan;60(1):66–71. <http://dx.doi.org/10.1093/occmed/kqp119>.
41. Garde AH, Hansen J, Kolstad HA, Larsen AD, Hansen AM. How do different definitions of night shift affect the exposure assessment of night work? *Chronobiol Int* 2016;33(6):595–8. <http://dx.doi.org/10.3109/07420528.2016.1167729>.
42. Härmä M, Koskinen A, Ropponen A, Puttonen S, Karhula K, Vahtera J et al. Validity of self-reported exposure to shift work. *Occup Environ Med* 2017 Mar;74(3):228–30. <http://dx.doi.org/10.1136/oemed-2016-103902>.
43. Vistisen HT, Garde AH, Frydenberg M, Christiansen P, Hansen AM, Andersen J et al. Short-term effects of night shift work on breast cancer risk: a cohort study of payroll data. *Scand J Work Environ Health* 2017 Jan;43(1):59–67. <http://dx.doi.org/10.5271/sjweh.3603>.
44. Knudsen LB, Olsen J. The Danish Medical Birth Registry. *Dan Med Bull* 1998 Jun;45(3):320–3.
45. Schmidt M, Schmidt SA, Sandegaard JL, Ehrenstein V, Pedersen L, Sørensen HT. The Danish National Patient Registry: a review of content, data quality, and research potential. *Clin Epidemiol* 2015 Nov;7:449–90. <http://dx.doi.org/10.2147/CLEP.S91125>.
46. Härmä M, Ropponen A, Hakola T, Koskinen A, Vanttola P, Puttonen S et al. Developing register-based measures for assessment of working time patterns for epidemiologic studies. *Scand J Work Environ Health* 2015 May;41(3):268–79. <http://dx.doi.org/10.5271/sjweh.3492>.
47. European Union's Working Time Directive (2003/88/EC) of the European Parliament and Council. 2003.
48. Statistics Denmark DISCO-88 - Danmarks Statistik Fagklassifikation. 1996.
49. Statistics Denmark DISCO-08 - Danmark Statistik Fagklassifikation. 2011.
50. International Statistical Classification of Diseases and Related Health Problems 10th Revision. 5th ed: World Health Organization, Geneva; 2011.
51. Saksvik IB, Bjorvatn B, Hetland H, Sandal GM, Pallesen S. Individual differences in tolerance to shift work--a systematic review. *Sleep Med Rev* 2011 Aug;15(4):221–35. <http://dx.doi.org/10.1016/j.smrv.2010.07.002>.
52. Gamble KL, Motsinger-Reif AA, Hida A, Borsetti HM, Servick SV, Ciarleglio CM et al. Shift work in nurses: contribution of phenotypes and genotypes to adaptation. *PLoS One* 2011 Apr;6(4):e18395. <http://dx.doi.org/10.1371/journal.pone.0018395>.
53. Nabe-Nielsen K, Jensen MA, Hansen AM, Kristiansen J,

- Garde AH. What is the preferred number of consecutive night shifts? results from a crossover intervention study among police officers in Denmark. *Ergonomics* 2016 Oct;59(10):1392–402. <http://dx.doi.org/10.1080/00140139.2015.1136698>.
54. Nabe-Nielsen K, Tüchsen F, Christensen KB, Garde AH, Diderichsen F. Differences between day and nonday workers in exposure to physical and psychosocial work factors in the Danish eldercare sector. *Scand J Work Environ Health* 2009 Jan;35(1):48–55. <http://dx.doi.org/10.5271/sjweh.1307>.
 55. Ekblad M, Gissler M, Korkeila J, Lehtonen L. Trends and risk groups for smoking during pregnancy in Finland and other Nordic countries. *Eur J Public Health* 2014 Aug;24(4):544–51. <http://dx.doi.org/10.1093/eurpub/ckt128>.
 56. Country Profiles on Nutrition, Physical Activity and Obesity in the 53 WHO European Region Member States. 2013.
 57. Chang YS, Chen HL, Wu YH, Hsu CY, Liu CK, Hsu C. Rotating night shifts too quickly may cause anxiety and decreased attentional performance, and impact prolactin levels during the subsequent day: a case control study. *BMC Psychiatry* 2014 Aug;14:218. <http://dx.doi.org/10.1186/s12888-014-0218-7>.
 58. Leung M, Tranmer J, Hung E, Korsiak J, Day AG, Aronson KJ. Shift Work, Chronotype, and Melatonin Patterns among Female Hospital Employees on Day and Night Shifts. *Cancer Epidemiol Biomarkers Prev* 2016 May;25(5):830–8. <http://dx.doi.org/10.1158/1055-9965.EPI-15-1178>.
 59. Dumont M, Paquet J. Progressive decrease of melatonin production over consecutive days of simulated night work. *Chronobiol Int* 2014 Dec;31(10):1231–8. <http://dx.doi.org/10.3109/07420528.2014.957304>.
 60. Niu SF, Chung MH, Chu H, Tsai JC, Lin CC, Liao YM et al. Differences in cortisol profiles and circadian adjustment time between nurses working night shifts and regular day shifts: A prospective longitudinal study. *Int J Nurs Stud* 2015 Jul;52(7):1193–201. <http://dx.doi.org/10.1016/j.ijnurstu.2015.04.001>.
 61. Ruhstaller KE, Bastek JA, Thomas A, McElrath TF, Parry SI, Durnwald CP. The Effect of early excessive weight gain on the development of hypertension in pregnancy. *Am J Perinatol* 2016 Oct;33(12):1205–10. <http://dx.doi.org/10.1055/s-0036-1585581>.
 62. Gudnadóttir TA, Bateman BT, Hernández-Díaz S, Luque-Fernandez MA, Valdimarsdóttir U, Zoega H. Body mass index, smoking and hypertensive disorders during pregnancy: a population based case-control study. *PLoS One* 2016 Mar;11(3):e0152187. <http://dx.doi.org/10.1371/journal.pone.0152187>.
 63. Gaillard R, Arends LR, Steegers EA, Hofman A, Jaddoe VW. Second- and third-trimester placental hemodynamics and the risks of pregnancy complications: the Generation R Study. *Am J Epidemiol* 2013 Apr;177(8):743–54. <http://dx.doi.org/10.1093/aje/kws296>.

Received for publication: 2 February 2018