



Scand J Work Environ Health 2020;46(2):152-160

<https://doi.org/10.5271/sjweh.3861>

Published online: 07 Nov 2019, Issue date: 01 Mar 2020

Shift work and the risk of knee osteoarthritis among Chinese workers: a retrospective cohort study

by [Zhou M](#), [Yang S](#), [Guo Y](#), [Wang D](#), [Qiu W](#), [Wang B](#), [Mu G](#), [Cao L](#), [Zhang Z](#), [Chen W](#)

Few studies have estimated the association between shift work and the risk of knee osteoarthritis (KOA). This retrospective cohort study showed that shift work was independently associated with the elevated risk of KOA. The longer the duration of shift work employment, the greater the risk, but the risk gradually decreased after leaving shift work.

Affiliation: School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, China. wchen@mails.tjmu.edu.cn

Refers to the following text of the Journal: [2002;28\(6\):394-401](#)

The following article refers to this text: [2024;50\(5\):317-328](#)

Key terms: [China](#); [Chinese worker](#); [circadian rhythm](#); [cohort study](#); [knee osteoarthritis](#); [metabolic disorder](#); [occupational health](#); [retrospective cohort study](#); [shift rotation](#); [shift work](#); [shift worker](#); [worker](#)

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/31696238



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Shift work and the risk of knee osteoarthritis among Chinese workers: a retrospective cohort study

by Min Zhou, PhD,^{1,2} Shijie Yang, PhD,^{1,2} Yanjun Guo, PhD,^{1,2} Dongming Wang, PhD,^{1,2} Weihong Qiu, MS,^{1,2} Bin Wang, PhD,^{1,2} Ge Mu, PhD,^{1,2} Limin Cao, PhD,^{1,2} Zhuang Zhang, PhD,^{1,2} Weihong Chen, PhD^{1,2}

Zhou M, Yang S, Guo Y, Wang D, Qiu W, Wang B, Cao L, Zhang Z, Chen W. Shift work and the risk of knee osteoarthritis among Chinese workers: a retrospective cohort study. *Scand J Work Environ Health*. 2020;46(2):152–160. doi:10.5271/sjweh.3861

Objectives Shift work is associated with metabolic disorders as it disturbs circadian rhythms. However, the potential association of shift work with knee osteoarthritis (KOA), a metabolic-related disease, has not been confirmed. The objective of this study was to quantify the association of shift work with the risk of KOA.

Methods This retrospective cohort study comprising 13 741 retired workers from the Dongfeng Motor Corporation. KOA was defined by knee pain and clinical X-ray radiographs. Occupational history, including job description and shift work, was collected from face-to-face interviews using questionnaires. Cox proportional hazards regression models were used to estimate exposure–response relationships.

Results During the 590 085 person-years of follow-up, a total of 847 cases of KOA (incidence of KOA was 143.5 per 100 000 person-years) were identified. After adjusting for potential confounders, shift work was independently associated with an elevated risk of KOA [hazard ratio (HR) 1.19, 95% confidence interval 1.03–1.36]. Compared with daytime workers, the risk of KOA increased with prolonged periods of shift work; the HR of KOA for participants with 1–9 years, 10–19 years, and ≥ 20 years shift work were 1.03 (95% CI 0.84–1.26), 1.19 (95% CI 0.98–1.46), and 1.42 (95% CI 1.15–1.76), respectively. The HR for KOA associated with shift work gradually decreased as the period after finishing shift work increased.

Conclusion Our results indicated that shift work could be an independent risk factor for KOA.

Key terms China; circadian rhythm; metabolic disorder; occupational health; shift rotation; shift worker.

Shift work – including night shifts, rotating shifts, and irregular schedules – is becoming increasingly common worldwide due to the rapid progress of industrialization and service demands. As much as 20% of the total active population participate in shift work schedules in industrialized countries worldwide (1). This employment practice typically divides a day into shifts, setting periods of time during which different groups of workers take up their posts to make use of or provide services throughout all 24 hours of each day.

Normally the body has fixed circadian rhythms, also known as the circadian clock, which presents in stable sleeping and feeding patterns. In shift work, the time of work activities, sleeping, and even diet for workers need to be constantly changed. Studies have reported that shift

work, particularly night shifts, can disrupt the circadian clock (2, 3). The circadian clock must be reset on a routine basis to ensure synchrony between the organism and the environment, as well as to maintain internal homeostasis including metabolism, inflammation, and hormone production (4). When the circadian clock is disturbed by shift work, an imbalance in internal homeostasis will occur. Previous studies have suggested that shift work increases the risk of chronic metabolic-related diseases (5–7). Furthermore, Berenbaum reported that the circadian clock regulated the homeostasis and function of musculoskeletal tissues including muscles, tendons, ligaments, bone, cartilage, subchondral bone, and synovial. Thus, disturbance of homeostasis of musculoskeletal tissues could occur due to disruption of the circadian clock (8).

¹ Department of Occupational and Environmental Health, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China.

² Key Laboratory of Environment and Health in Ministry of Education & Ministry of Environmental Protection, and State Key Laboratory of Environmental Health (Incubating), School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China.

Correspondence to: Weihong Chen, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, China [E-mail: wchen@mails.tjmu.edu.cn].

Knee osteoarthritis (KOA) – manifested as degenerative arthritis – is a disease of the entire joint, involving cartilage, synovial tissue, tendons, ligaments, and subchondral bone (9). Common and disturbing symptoms of KOA include pain or aching stiffness in or around the knee, localized swelling around the knee, and/or limited motion of the knee. KOA most commonly affects people aged ≥ 45 years. In 2016, it was reported that 301.57 million people worldwide were affected by osteoarthritis, and 14.70 million new cases were recorded (10). Moreover, KOA is currently the leading cause of lower-limb disability in adults and results in a huge medical burden worldwide. Results of the 2016 Global Burden of Disease Study showed that osteoarthritis caused 16.28 million disability-adjusted life-years in 195 countries (11). Therefore, potential risk factors for KOA have received widespread attention.

Published reports have suggested that KOA is influenced by obesity, aging, and overloaded external force on knee joints such as heavy loads or repeated bending from occupational activities or sports (12–14). Recently, evidence is emerging that systemic metabolism plays a key role in the onset and progression of KOA (15). Several studies have demonstrated that metabolic disorders are important risk factors of KOA (16–18). Our previous research also showed that hyperlipidemia was positively independently associated with an elevated risk of KOA in a cohort study (19). However, whether metabolic disorders caused by the disturbance of the circadian clock following shift work is related to KOA remains to be established. Few studies have focused on the association of shift work with the risk of KOA.

In order to fill the gap in knowledge, we conducted the present retrospective cohort study among 13 741 retired workers from the Dongfeng Motor Corporation (20). The objective of this study was to quantify the association of shift work with the risk of KOA and to evaluate the persistent association of shift work with the risk of KOA after leaving shift work.

Methods

Study population

A group of 14 438 retired workers from Dongfeng Motor Corporation, ie, all of the participants that underwent a physical examination at the Dongfeng Central Hospital, were recruited in this study. In 2013, trained professionals interviewed all participants face-to-face and collected information on demographics, knee pain, medical history (osteoarthritis, rheumatoid arthritis, and accidental injury), occupational history (job titles, calendar years of each job for each individual's full duration

of employment, job content, and shift work experience), and lifestyle (smoking, alcohol drinking, and physical exercise). After the interview, all participants underwent examination of the knee joint including tenderness, range of motion, extension test, and McMurray's test (21). A total of 13 741 participants were included after excluding individuals with incomplete information on occupational history or knee health status, knee surgery caused by accidental injury that was associated with secondary KOA, history of rheumatoid arthritis, diagnosis of KOA prior to the start of employment, diagnosis of KOA prior to the start of shift work.

This study was approved by the Medical Ethics Committee of Dongfeng General Hospital, Dongfeng Motor Corporation, and the School of Public Health, Tongji Medical College, Huazhong University of Science and Technology. All participants signed written informed consent.

Follow-up

The first year of employment for each participant was retrospectively assigned as the beginning of the follow-up period. The end of the follow-up period was defined as the year of physician-diagnosed KOA or 2013, whichever came first.

Assessment of KOA

Self-reported knee pain was defined when one or both knee(s) met at least one of the following conditions: (i) pain or aching stiffness in or around the knee on most days for ≥ 1 month during the preceding 12 months; (ii) persistent pain or aching stiffness in or around the knee within the past week.

Information regarding the clinical diagnosis of KOA and the year of diagnosis was collected from the questionnaires and confirmed by insurance records and treatment information. Clinical KOA cases were only defined if the participant had knee pain (as defined above) and the bilateral weight-bearing anteroposterior X-ray radiographs showed a Kellgren & Lawrence grade ≥ 2 in at least one knee (22).

Assessment of shift work

Shift work was identified as working with a schedule involving unusual working hours as opposed to the normal daytime work schedule, ie, 08:00–17:00 hours for ≥ 1 year. There were two common kinds of shift work for the participants in this study: two-shifts including day work (08:00–17:30) and night work (17:30–02:30) shifts on a weekly rotation; three-shifts during which three crews of workers succeeded each other at 08:00, 16:00, and 00:00 hours. The workers in any kind of shift work took turns

to work in the early morning and at night. Shift work duration was calculated by the starting year of shift work to the earliest year out of the following: the end of shift work, the first year of clinical KOA diagnosis, or 2013, and was categorized into 1–9, 10–19, and ≥ 20 years.

Years after leaving shift work was the duration from the year leaving shift work to the end of the follow-up period. The duration of leaving shift work was divided into tertiles based on the duration of shift work.

Assessment of covariates

Body weight and height was measured to the nearest 0.1 kg and 0.1 cm, with participants wearing light indoor clothing and no shoes. Body mass index (BMI) was calculated from weight in kilograms divided by body height in squared meters.

Information regarding smoking status (current/former/never), drinking status (current/former/never), and physical exercise was collected from the questionnaires. Participants who had been smoking as much as one cigarette per day for ≥ 6 months were considered current smokers, and those who had been drinking alcohol as often as once per week for ≥ 6 months were considered current drinkers. Physical exercise was defined as regular exercise of ≥ 20 minutes per day over the past 6 months (20).

Based on the job held for the longest duration, the work posture was grouped into sitting, standing, squatting or kneeling, or bending if such posture was maintained for more than half of one work shift (ie, ≥ 4 hours for most cases) according to the job content description. The occupational chemicals (eg, industrial dust) level for each job title in the workplace came from the company occupational hazard monitoring records, and was categorized into “yes” when the concentrations of chemicals were higher than the national occupational chemicals exposure limit. Early retirement was defined as retiring before the statutory age of retirement (55 years for male workers and 50 years for females).

Statistical analyses

Basic characteristics of participants were presented as means and standard deviation (SD) for normally distributed continuous variables and as numbers and percentages for categorical variables. Continuous variables were compared using ANOVA analysis across shift work duration, whereas categorical variables were compared using Chi-square test.

Cox proportional hazards regressions were used to evaluate the association of shift work with KOA, and hazard ratios (HR) and 95% confidence intervals (CI) were calculated. We used age as the time variable to define the risk sets for incidence, and we used cases of KOA as the outcome variable in regression analyses.

The basic model 1 only included shift work. Model 2 also included gender and BMI (continuous) as they were reported to be significantly associated with the risk of KOA (23). Model 3 also included lifestyle variables (smoking, drinking, and physical exercise). Model 4 was further adjusted for working years. Model 5 was further adjusted for work posture and occupational chemicals, for the reason that shift work experience may vary according to different jobs. Models were conducted with daytime workers as the reference group.

Multivariable stepwise Cox proportional hazards regressions that started with relevant factors (including gender, BMI, smoking, drinking, physical exercise, working years, work postures, occupational chemicals, and shift work) and ended with variables that were statistically significantly associated with a P-value < 0.05 in a stepwise selection procedure, were used to explore the independent association between shift work and risk of KOA.

To investigate the combined effects of shift work and other factors on KOA, we estimated HR by crossed dichotomized shift work (shift worker = A+; daytime worker = A-) and each of the other factors (eg, obesity = B+; normal or low body weight = B-). The relative excess risk due to interaction (RERI) (calculated as $HR_{A+B+} - HR_{A+B-} - HR_{A-B+} + 1$) was used to evaluate additive interaction (24). Multiplicative interaction was examined by adding an interaction term of shift work (A+/A-) and each of the other factors (B+/B-) to the Cox proportional hazards regressions model.

To evaluate the persistent role of shift work in the risk of KOA, the HR of KOA associated with shift work was calculated, respectively, at different levels of duration of leaving shift work. Linear trend tests were conducted by including the median value for each level of duration of leaving shift work as a continuous variable in the models.

Statistical tests were two-sided with a significance set at $P < 0.05$. All statistical analyses were performed using SAS 9.4 software (SAS Institute, Cary, NC, USA).

Results

The basic characteristics of the participants in 2013 are shown in table 1. A total of 13 741 participants (females 7453, 54.2%) were included in the analysis and the mean age was 64.4 (SD 8.6) years old. A total of 5390 (39.2%) workers had engaged in shift work, including 1877 (13.7%) workers who had undergone shift work for 1–9 years, 1859 (13.5%) for 10–19 years, and 1654 (12.0%) for ≥ 20 years. Participants with a longer duration of shift work were more likely to be males, current smokers, older, have worked in awkward postures, and been exposed to occupational chemicals.

Table 1. Basic characteristics of study population according to shift work. [BMI=body mass index; KOA=knee osteoarthritis; NA=not applicable; SD=standard deviation]. Continuous variables were compared using ANOVA analysis across shift work duration, whereas categorical variables were compared using Chi-square test.

| Variables | Total | | Duration of shift work | | | | | | | | P-value |
|--------------------------|--------------|------------|------------------------|------------|-------------|------------|-------------|------------|-------------|------------|---------|
| | | | Never | | 1–9 years | | 10–19 years | | ≥20 years | | |
| | N (%) | Mean (SD) | N (%) | Mean (SD) | N (%) | Mean (SD) | N (%) | Mean (SD) | N (%) | Mean (SD) | |
| Female gender | 7453 (54.2) | | 4512 (54.0) | | 1208 (64.4) | | 1068 (57.4) | | 665 (40.2) | | <0.001 |
| Age (years) | | 64.4 (8.6) | | 65.4 (8.2) | | 62.9 (8.6) | | 63.5 (8.1) | | 64.7 (7.5) | <0.001 |
| BMI (kg/m ²) | | 24.2 (3.3) | | 24.2 (3.3) | | 24.2 (3.3) | | 24.2 (3.4) | | 24.2 (3.3) | 0.836 |
| Smokers | | | | | | | | | | | <0.001 |
| Non | 9690 (70.7) | | 5966 (71.6) | | 1407 (75.0) | | 1292 (69.7) | | 1025 (62.0) | | |
| Current | 2155 (15.8) | | 1259 (15.1) | | 270 (14.4) | | 291 (15.7) | | 335 (20.3) | | |
| Former | 1868 (13.6) | | 1105 (13.3) | | 198 (10.6) | | 271 (14.6) | | 294 (17.8) | | |
| Drinkers | | | | | | | | | | | <0.001 |
| Non | 9354 (68.3) | | 5784 (69.5) | | 1328 (70.9) | | 1244 (67.1) | | 998 (60.4) | | |
| Current | 3473 (25.3) | | 2035 (24.4) | | 439 (23.4) | | 479 (25.8) | | 520 (31.5) | | |
| Former | 50 (6.4) | | 507 (6.1) | | 105 (5.6) | | 130 (7.0) | | 134 (8.1) | | |
| Physical exercise | 12379 (90.1) | | 7504 (89.9) | | 1695 (90.3) | | 1666 (89.6) | | 1514 (91.5) | | 0.180 |
| Work postures | | | | | | | | | | | <0.001 |
| Sitting | 4781 (34.9) | | 3473 (41.8) | | 551 (29.5) | | 406 (21.9) | | 351 (21.3) | | |
| Standing | 6011 (43.9) | | 3218 (38.7) | | 917 (49.0) | | 983 (53.0) | | 893 (54.1) | | |
| Bending | 1868 (13.6) | | 1016 (12.2) | | 254 (13.6) | | 335 (18.1) | | 263 (15.9) | | |
| Squatting or kneeling | 1022 (7.5) | | 600 (7.2) | | 148 (7.9) | | 130 (7.0) | | 144 (8.7) | | |
| Occupational chemicals | 1728 (12.6) | | 883 (10.6) | | 244 (13.0) | | 298 (16.0) | | 303 (18.3) | | <0.001 |
| Shift work | | | | | | | | | | | |
| Daytime workers | 8351 (60.8) | | 8351 (100.0) | | NA | | NA | | NA | | <0.001 |
| Two shifts | 4079 (29.7) | | NA | | 1489 (79.3) | | 1383 (74.4) | | 1207 (73.0) | | |
| Three shifts | 1311 (9.5) | | NA | | 388 (20.7) | | 476 (25.6) | | 447 (27.0) | | |
| Early retirement years | | 1.7 (3.5) | | 1.6 (3.5) | | 2.2 (4.0) | | 2.0 (3.5) | | 1.4 (2.8) | <0.001 |
| KOA | 874 (6.4) | | 531 (6.4) | | 114 (6.1) | | 120 (6.5) | | 109 (6.6) | | 0.933 |

During the 590 085 person-years of follow-up, we identified 847 cases of KOA. The incidence of KOA in this cohort was 143.5 per 100 000 person-years. After adjusting for gender and BMI, the HR of KOA among shift workers was 1.15 (95% CI 1.01–1.32) when compared with daytime workers. Controlling for lifestyles and occupational characteristics did not change this association. Different shift types showed similar associations with the risk of KOA; the fully adjusted HR for two shifts and three shifts were 1.19 (95% CI 1.02–1.38) and 1.19 (95% CI 0.94–1.50), respectively. Both continuous and categorical analyses suggested positive exposure–response relationship between prolonged periods of shift work and the risk of KOA. Compared with daytime workers, the HR of KOA among participants with 1–9, 10–19, and ≥20 years of shift work were 1.03 (95% CI 0.84–1.26), 1.19 (95% CI 0.98–1.46), and 1.42 (95% CI 1.15–1.76), respectively, after adjusting for gender, BMI, lifestyles, and occupational characteristics. In the fully adjusted model, each 5-year increase in the duration of shift work was associated with a 5% increase in the risk of KOA (table 2).

In the stepwise multivariate Cox proportional hazards regression analyses, the following associations remained stable: between shift work and KOA (HR 1.16, 95% CI 1.01–1.33), between categorical shift work dura-

tion and KOA (HR 1.02, 1.16, 1.36 for 1–9, 10–19, and ≥20 years of shift work), and between continuous shift work duration and KOA (HR 1.04, 95% CI 1.01–1.08) (table 3).

The results of the combined effects of shift work with other factors (gender, obesity, physical exercise, smoking, drinking, work posture, occupational chemicals, and early retirement) in relation to the risk of KOA are shown in table 4. Compared with participants without shift work and lower BMI, the HR for KOA among those with shift work and lower BMI, without shift work and higher BMI, and with shift work and higher BMI were 1.17 (95% CI 0.43–3.23), 2.18 (95% CI 1.16–4.07), and 2.60 (95% CI 1.38–4.88), respectively. Women with shift work showed the highest risk of KOA (HR 2.62, 95% CI 2.08–3.30) compared with men with daytime work. However, neither the multiplicative interaction terms nor the RERI were statistically significant, suggesting that the hypothesis of multiplicative interaction or additive interaction between shift work and other risk factors cannot be rejected.

We then estimated the persistent role of shift work in the risk of KOA after leaving shift work. As shown in figure 1, among participants with ≥20 years of shift work, the HR of KOA associated with shift work showed a decreasing trend as the duration of leaving shift work

Table 2. Hazard ratios (HR) and 95% confidence intervals (CI) for knee osteoarthritis (KOA) by shift work. [NA=not applicable].

| Variables | KOA | Model 1 ^a | | Model 2 ^b | | Model 3 ^c | | Model 4 ^d | | Model 5 ^e | |
|---|-----------|----------------------|-----------|----------------------|-----------|----------------------|-----------|----------------------|-----------|----------------------|-----------|
| | N (%) | HR | 95% CI | HR | 95% CI | HR | 95% CI | HR | 95% CI | HR | 95% CI |
| Shift work | | | | | | | | | | | |
| No | 531 (6.4) | 1 | reference | 1 | reference | 1 | reference | 1 | reference | 1 | reference |
| Yes | 343 (6.4) | 1.14 | 0.99–1.31 | 1.15 | 1.01–1.32 | 1.14 | 1.00–1.31 | 1.16 | 1.01–1.33 | 1.19 | 1.03–1.36 |
| Shift work duration (categorical) | | | | | | | | | | | |
| Daytime workers (years) | 531 (6.4) | 1 | reference | 1 | reference | 1 | reference | 1 | reference | 1 | reference |
| 1–9 | 114 (6.1) | 1.13 | 0.92–1.38 | 1.03 | 0.84–1.27 | 1.03 | 0.84–1.27 | 1.02 | 0.83–1.25 | 1.03 | 0.84–1.26 |
| 10–19 | 120 (6.5) | 1.17 | 0.96–1.43 | 1.16 | 0.95–1.41 | 1.15 | 0.95–1.41 | 1.16 | 0.95–1.41 | 1.19 | 0.98–1.46 |
| ≥20 | 109 (6.6) | 1.13 | 0.92–1.38 | 1.29 | 1.05–1.58 | 1.28 | 1.04–1.57 | 1.36 | 1.10–1.68 | 1.42 | 1.15–1.76 |
| Shift work duration (per 5-year increase) | NA | 1.02 | 0.99–1.05 | 1.04 | 1.01–1.07 | 1.04 | 1.01–1.07 | 1.04 | 1.01–1.08 | 1.05 | 1.02–1.09 |
| Shift work types | | | | | | | | | | | |
| Daytime workers | 531 (6.4) | 1 | reference | 1 | reference | 1 | reference | 1 | reference | 1 | reference |
| Two shifts | 259 (6.4) | 1.14 | 0.99–1.33 | 1.15 | 0.99–1.34 | 1.15 | 0.99–1.33 | 1.16 | 1.01–1.35 | 1.19 | 1.02–1.38 |
| Three shifts | 84 (6.4) | 1.14 | 0.91–1.44 | 1.14 | 0.91–1.44 | 1.14 | 0.90–1.44 | 1.16 | 0.92–1.46 | 1.19 | 0.94–1.50 |

^a Crude HR.^b Adjusted for gender, and body mass index (BMI) (continuous).^c Model 2 + smoking, drinking, and physical exercise.^d Model 3 + working years.^e Model 4 + work posture, and occupational chemicals.**Table 3.** Hazard ratios (HR) and 95% confidence intervals (CI) for knee osteoarthritis (KOA) from multivariate stepwise cox proportional hazards regression model. [BMI=body mass index; NA=not applicable].

| Variables | KOA | Model 1 ^a | | Model 2 ^b | | Model 3 ^c | |
|--|-----------|----------------------|-----------|----------------------|-----------|----------------------|-----------|
| | N (%) | HR | 95% CI | HR | 95% CI | HR | 95% CI |
| Shift work | | | | | | | |
| No | 531 (6.4) | 1 | reference | | | | |
| Yes | 343 (6.4) | 1.16 | 1.01–1.33 | | | | |
| Shift work duration (categorical) | | | | | | | |
| Daytime workers (years) | 531 (6.4) | | | 1 | reference | | |
| 1–9 | 114 (6.1) | | | 1.02 | 0.83–1.25 | | |
| 10–19 | 120 (6.5) | | | 1.16 | 0.95–1.41 | | |
| ≥20 | 109 (6.6) | | | 1.36 | 1.10–1.68 | | |
| Shift work duration (per 5-years increase) | NA | | | | | 1.04 | 1.01–1.08 |
| Gender | | | | | | | |
| Male | 264 (4.2) | 1 | reference | 1 | reference | 1 | reference |
| Female | 610 (8.2) | 2.67 | 2.18–3.27 | 2.67 | 2.18–3.27 | 2.67 | 2.18–3.28 |
| BMI (kg/m ²) | NA | 1.07 | 1.06–1.09 | 1.07 | 1.05–1.09 | 1.07 | 1.05–1.09 |
| Drinkers | | | | | | | |
| Non | 638 (6.8) | 1 | reference | 1 | reference | 1 | reference |
| Current | 186 (5.4) | 1.44 | 1.20–1.72 | 1.43 | 1.19–1.72 | 1.43 | 1.20–1.72 |
| Former | 50 (5.7) | 1.41 | 1.04–1.92 | 1.40 | 1.03–1.91 | 1.40 | 1.03–1.91 |

^a Initially included gender, BMI (continuous), smoking, drinking, physical exercise, working years, work postures, occupational chemicals, and shift work.^b Initially included gender, BMI (continuous), smoking, drinking, physical exercise, working years, work postures, occupational chemicals, and duration of shift work (categorical).^c Initially included gender, BMI (continuous), smoking, drinking, physical exercise, working years, work postures, occupational chemicals, and duration of shift work (continuous).

increased, with a P-value for trend of 0.009; however, the HR decreased without statistical significance among those with 1–9 or 10–19 years of shift work (P-values for trend 0.162 and 0.312, respectively).

In order to prove the credibility of our results, we examined the association of gender, BMI, lifestyle, awkward work postures, and working years with KOA. As shown in supplementary table S1 (www.sjweh.fi/show_abstract.php?abstract_id=3861), female gender (HR 2.87, 95% CI 2.48–3.32), obesity (HR 1.09, 95% CI 1.06–1.11), standing (HR 1.20, 95% CI 1.03–1.39) or bending work posture (HR 1.19, 95% CI 0.96–1.47),


and early retirement (HR 1.61, and 4.07 for 1–4 and ≥5 years of early retirement, respectively) were risk factors of KOA.

Discussion

In the present study, we identified a positive association between shift work and the risk of KOA among 13 741 retired workers. There was an exposure–response relationship between prolonged shift work duration and the

Table 4. Combined effect of shift work and other potential factors on the risk of knee osteoarthritis (KOA). [BMI=body mass index; CI=confidence interval; HR=hazard ratio; RERI=relative excess risk due to interaction].

| | Daytime workers | | | Shift workers | | | Test for multiplicative interaction | | Test for additive interaction | |
|---------------------------------------|-----------------|------|-----------|---------------|------|-----------|-------------------------------------|-----------|-------------------------------|------------|
| | KOA N (%) | HR | 95% CI | KOA N (%) | HR | 95% CI | HR | 95% CI | RERI | 95% CI |
| Gender ^a | | | | | | | | | | |
| Male | 158 (4.1) | 1 | reference | 106 (4.3) | 1.20 | 0.94–1.54 | | | | |
| Female | 373 (8.3) | 2.25 | 1.81–2.81 | 237 (8.1) | 2.62 | 2.08–3.30 | 0.97 | 0.72–2.81 | 0.21 | -0.01–0.43 |
| BMI (kg/m ²) ^a | | | | | | | | | | |
| <24 | 10 (3.8) | 1 | reference | 6 (3.5) | 1.17 | 0.43–3.23 | | | | |
| ≥24 | 521 (6.4) | 2.18 | 1.16–4.07 | 337 (6.5) | 2.60 | 1.38–4.88 | 1.02 | 0.37–2.82 | 0.25 | -0.85–1.35 |
| Physical exercise ^a | | | | | | | | | | |
| No | 58 (6.9) | 1 | reference | 34 (6.6) | 1.16 | 0.76–1.78 | | | | |
| Yes | 473 (6.3) | 1.06 | 0.80–1.39 | 309 (6.3) | 1.26 | 0.95–1.68 | 1.03 | 0.66–1.62 | 0.05 | -0.44–0.54 |
| Smoking ^b | | | | | | | | | | |
| No | 435 (7.3) | 1 | reference | 269 (7.2) | 1.17 | 1.01–1.36 | | | | |
| Yes | 96 (4.1) | 0.93 | 0.72–1.20 | 74 (4.5) | 1.14 | 0.87–1.51 | 1.06 | 0.75–1.48 | 0.05 | -0.34–0.23 |
| Drinking ^b | | | | | | | | | | |
| No | 397 (6.9) | 1 | reference | 241 (6.8) | 1.14 | 0.97–1.34 | | | | |
| Yes | 134 (5.3) | 1.38 | 1.11–1.70 | 102 (5.6) | 1.76 | 1.39–2.22 | 1.13 | 0.83–1.53 | 0.10 | -0.24–0.45 |
| Work posture ^a | | | | | | | | | | |
| Sitting | 305 (6.2) | 1 | reference | 270 (6.6) | 1.30 | 1.11–1.54 | | | | |
| Others | 226 (6.5) | 1.29 | 1.08–1.55 | 73 (5.6) | 1.24 | 0.95–1.61 | 0.76 | 0.56–1.03 | 0.27 | -0.01–0.55 |
| Occupational chemicals ^a | | | | | | | | | | |
| No | 482 (6.4) | 1 | reference | 283 (6.2) | 1.15 | 0.99–1.33 | | | | |
| Yes | 49 (5.6) | 0.81 | 0.60–1.09 | 60 (7.1) | 1.28 | 0.97–1.68 | 1.44 | 0.96–2.16 | 0.32 | -0.10–0.74 |
| Early retirement ^a | | | | | | | | | | |
| No | 308 (5.4) | 1 | reference | 171 (5.2) | 1.13 | 0.94–1.37 | | | | |
| Yes | 223 (8.3) | 2.15 | 1.80–2.58 | 172 (8.2) | 2.46 | 2.02–3.00 | 1.01 | 0.77–1.32 | 0.17 | -0.33–0.68 |

^a Models were adjusted for working years.^b Models were adjusted for working years and gender.**Figure 1.** Hazard ratios for knee osteoarthritis (KOA) according to duration of shift work and the years after leaving shift work, compared to daytime workers. The model was adjusted for age, gender body mass index (continuous), smoking status, drinking status, physical exercise, working years, and work posture. Tests of linear trend were conducted by including the median value for each duration level of leaving shift work as a continuous variable in the models.

elevated risk of KOA. However, the association of shift work with KOA gradually decreased as the duration of leaving shift work increased.

Although several cross-sectional studies have shown that shift work is positively associated with musculoskeletal disorders, including low-back and shoulder pain (25–27), few studies have estimated the association between shift work and KOA. Our results reported here have notable public health implications. Shift work remains a vital component of the modern economy, and the percentage of shift workers has continued to increase in many countries (28). An increased risk of KOA among shift workers in this study has indicated that the adverse health effects of shift work were diverse. Considering the increasing prevalence and burden of KOA, and the significant reduction in quality of life and work productivity induced by KOA (11, 29), further action should be carried out to avoid or regulate shift work.

Although the mechanisms for the increased risk of KOA due to shift work remain unclear, several factors may be underlying in this association. First, the circadian clock regulates various rhythmic biological processes, including wake-sleep cycles, cell cycles, energy metabolism, and hormone secretion. Shift work is generally associated with chronic circadian clock misalignment (30). This circadian misalignment directly disturbs the homeostasis of cartilage, bone, and tendons (31–33). Circadian misalignment has also been found to result in metabolic disorders and inflammatory consequences (34, 35), which have been linked to cartilage and bone homeostasis (36). Second, shift work may influence KOA through sleep quality. In our previous reports, shift work was associated with reduced sleep quality (37). We further analyzed the mediating role of sleep quality in the relationship between shift work and KOA and found that poor sleep quality significantly mediated the total effects of shift work on KOA by 15.84%. Poor sleep may directly reduce the recovery time of knee problems. Third, shift workers tended to have different dietary habits from daytime workers, ie, greater energy density, increased saturated fat, and decreased dietary fiber were reported among shift workers (38, 39). A high intake of fat and saturated fatty acids was associated with the onset and progression of KOA, while dietary fiber was inversely associated with KOA worsening (40, 41).

Furthermore, our findings also suggested that the effect of shift work on the risk of KOA could decrease after leaving shift work. The risk of KOA may return to the daytime worker level after 25 years of ceasing shift work for participants that had ever been engaged in 1–9 years of shift work, while it took 20 years for those with 10–19 years of shift work, and it was reported to take <5 years for those with ≥ 20 years of shift work. These results suggest that the first few years could be the most

critical time for shift workers to adapt themselves to irregular circadian clocks.

The strengths of this study include the large sample size, detailed occupational history, and detailed information on a wide range of potential confounders. We are aware of the limitations of this study. First, we did not provide knee X-ray examinations for each participant although they could have a free one if they had knee symptoms and the company was willing pay for such examinations. Second, information regarding shift work was self-reported, which may lead to misclassifications of the exposure. However, we checked the shift work information (yes/no) used in the present study with data collected in 2008, and the agreement (kappa coefficient) was 0.76 (95% CI 0.75–0.78). Third, although information regarding lifestyle was collected over the past year, the individuals may have changed their lifestyle over time. Fourth, we failed to include retired workers who died before the 2013 recruitment or moved to other cities, and this may have caused selection bias. But according to company records, 87% of retired employees attended this medical examination.

In conclusion, the present study showed that shift work was independently associated with an elevated risk of KOA. These findings have substantial implications for preventing KOA by modifying shift work policies.

Acknowledgments

The authors are indebted to Dr Stephen S Bao at the Washington State Department of Labor and Industries for providing valuable comments.

Funding

This work was supported by the National Natural Science Foundation of China (grant number 81573121) and the Fundamental Research Funds for the Central Universities, Huazhong University of Science and Technology (grant number 2016JCTD116). The funders played no role in study design; data collection, analysis, or interpretation; the writing of the manuscript; or the decision to approve publication of the finished manuscript.

Disclosure of interest

The authors declare no conflicts of interest.

References

1. Eurofound. Working time and work–life balance in a life course perspective, Eurofound, Dublin. 2012.
2. Ball LJ, Palesh O, Kriegsfeld LJ. The Pathophysiologic Role of Disrupted Circadian and Neuroendocrine Rhythms in Breast Carcinogenesis. *Endocr Rev* 2016 Oct;37(5):450–66. <https://doi.org/10.1210/er.2015-1133>.
3. Wright KP Jr, Bogan RK, Wyatt JK. Shift work and the assessment and management of shift work disorder (SWD). *Sleep Med Rev* 2013 Feb;17(1):41–54. <https://doi.org/10.1016/j.smrv.2012.02.002>.
4. Van Laake LW, Lüscher TF, Young ME. The circadian clock in cardiovascular regulation and disease: Lessons from the Nobel Prize in Physiology or Medicine 2017. *Eur Heart J* 2018 Jun;39(24):2326–9. <https://doi.org/10.1093/eurheartj/ehx775>.
5. De Bacquer D, Van Risseghem M, Clays E, Kittel F, De Backer G, Braeckman L. Rotating shift work and the metabolic syndrome: a prospective study. *Int J Epidemiol* 2009 Jun;38(3):848–54. <https://doi.org/10.1093/ije/dyn360>.
6. Hublin C, Partinen M, Koskenvuo K, Silventoinen K, Koskenvuo M, Kaprio J. Shift-work and cardiovascular disease: a population-based 22-year follow-up study. *Eur J Epidemiol* 2010 May;25(5):315–23. <https://doi.org/10.1007/s10654-010-9439-3>.
7. Pan A, Schernhammer ES, Sun Q, Hu FB. Rotating night shift work and risk of type 2 diabetes: two prospective cohort studies in women. *PLoS Med* 2011 Dec;8(12):e1001141. <https://doi.org/10.1371/journal.pmed.1001141>.
8. Berenbaum F, Meng QJ. The brain-joint axis in osteoarthritis: nerves, circadian clocks and beyond. *Nat Rev Rheumatol* 2016 Sep;12(9):508–16. <https://doi.org/10.1038/nrrheum.2016.93>.
9. Felson DT. Clinical practice. Osteoarthritis of the knee. *N Engl J Med* 2006 Feb;354(8):841–8. <https://doi.org/10.1056/NEJMc051726>.
10. GBD 2016 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017 Sep;390(10100):1211–59. [https://doi.org/10.1016/S0140-6736\(17\)32154-2](https://doi.org/10.1016/S0140-6736(17)32154-2).
11. GBD 2016 DALYs and HALE Collaborators. Global, regional, and national disability-adjusted life-years (DALYs) for 333 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017 Sep;390(10100):1260–344. [https://doi.org/10.1016/S0140-6736\(17\)32130-X](https://doi.org/10.1016/S0140-6736(17)32130-X).
12. Haukka E, Ojajärvi A, Takala EP, Viikari-Juntura E, Leino-Arjas P. Physical workload, leisure-time physical activity, obesity and smoking as predictors of multisite musculoskeletal pain. A 2-year prospective study of kitchen workers. *Occup Environ Med* 2012 Jul;69(7):485–92. <https://doi.org/10.1136/oemed-2011-100453>.
13. Solidaki E, Chatzi L, Bitsios P, Coggon D, Palmer KT, Kogevinas M. Risk factors for new onset and persistence of multi-site musculoskeletal pain in a longitudinal study of workers in Crete. *Occup Environ Med* 2013 Jan;70(1):29–34. <https://doi.org/10.1136/oemed-2012-100689>.
14. Lau EC, Cooper C, Lam D, Chan VN, Tsang KK, Sham A. Factors associated with osteoarthritis of the hip and knee in Hong Kong Chinese: obesity, joint injury, and occupational activities. *Am J Epidemiol* 2000 Nov;152(9):855–62. <https://doi.org/10.1093/aje/152.9.855>.
15. Mobasheri A, Rayman MP, Gualillo O, Sellam J, van der Kraan P, Fearon U. The role of metabolism in the pathogenesis of osteoarthritis. *Nat Rev Rheumatol* 2017 May;13(5):302–11. <https://doi.org/10.1038/nrrheum.2017.50>.
16. Schett G, Kleyer A, Perricone C, Sahinbegovic E, Iagnocco A, Zwerina J et al. Diabetes is an independent predictor for severe osteoarthritis: results from a longitudinal cohort study. *Diabetes Care* 2013 Feb;36(2):403–9. <https://doi.org/10.2337/dc12-0924>.
17. Haugen IK, Ramachandran VS, Misra D, Neogi T, Niu J, Yang T et al. Hand osteoarthritis in relation to mortality and incidence of cardiovascular disease: data from the Framingham heart study. *Ann Rheum Dis* 2015 Jan;74(1):74–81. <https://doi.org/10.1136/annrheumdis-2013-203789>.
18. Niu J, Clancy M, Aliabadi P, Vasani R, Felson DT. Metabolic Syndrome, Its Components, and Knee Osteoarthritis: The Framingham Osteoarthritis Study. *Arthritis Rheumatol* 2017 Jun;69(6):1194–203. <https://doi.org/10.1002/art.40087>.
19. Zhou M, Guo Y, Wang D, Shi D, Li W, Liu Y et al. The cross-sectional and longitudinal effect of hyperlipidemia on knee osteoarthritis: results from the Dongfeng-Tongji cohort in China. *Sci Rep* 2017 Aug;7(1):9739. <https://doi.org/10.1038/s41598-017-10158-8>.
20. Wang F, Zhu J, Yao P, Li X, He M, Liu Y et al. Cohort Profile: the Dongfeng-Tongji cohort study of retired workers. *Int J Epidemiol* 2013 Jun;42(3):731–40. <https://doi.org/10.1093/ije/dys053>.
21. Jackson JL, O'Malley PG, Kroenke K. Evaluation of acute knee pain in primary care. *Ann Intern Med* 2003 Oct;139(7):575–88. <https://doi.org/10.7326/0003-4819-139-7-200310070-00010>.
22. Kellgren JH, Lawrence JS. Radiological assessment of osteoarthrosis. *Ann Rheum Dis* 1957 Dec;16(4):494–502. <https://doi.org/10.1136/ard.16.4.494>.
23. Reijman M, Pols HA, Bergink AP, Hazes JM, Belo JN, Lieveense AM et al. Body mass index associated with onset and progression of osteoarthritis of the knee but not of the hip: the Rotterdam Study. *Ann Rheum Dis* 2007 Feb;66(2):158–62. <https://doi.org/10.1136/ard.2006.053538>.
24. Hosmer DW, Lemeshow S. Confidence interval estimation of interaction. *Epidemiology* 1992 Sep;3(5):452–6. <https://doi.org/10.1097/00001648-199209000-00012>.

25. Takahashi M, Matsudaira K, Shimazu A. Disabling low back pain associated with night shift duration: sleep problems as a potentiator. *Am J Ind Med* 2015 Dec;58(12):1300–10. <https://doi.org/10.1002/ajim.22493>.
26. Lipscomb JA, Trinkoff AM, Geiger-Brown J, Brady B. Work-schedule characteristics and reported musculoskeletal disorders of registered nurses. *Scand J Work Environ Health* 2002 Dec;28(6):394–401. <https://doi.org/10.5271/sjweh.691>.
27. Choobineh A, Soltanzadeh A, Tabatabaee H, Jahangiri M, Khavaji S. Health effects associated with shift work in 12-hour shift schedule among Iranian petrochemical employees. *Int J Occup Saf Ergon* 2012;18(3):419–27. <https://doi.org/10.1080/10803548.2012.11076937>.
28. Eurofound. Sixth European Working Conditions Survey - Overview report (2017 update), Publications Office of the European Union, Luxembourg. 2017.
29. Agaliotis M, Fransen M, Bridgett L, Nairn L, Votrubec M, Jan S et al. Risk factors associated with reduced work productivity among people with chronic knee pain. *Osteoarthritis Cartilage* 2013 Sep;21(9):1160–9. <https://doi.org/10.1016/j.joca.2013.07.005>.
30. Gumenyuk V, Howard R, Roth T, Korzyukov O, Drake CL. Sleep loss, circadian mismatch, and abnormalities in reorienting of attention in night workers with shift work disorder. *Sleep (Basel)* 2014 Mar;37(3):545–56. <https://doi.org/10.5665/sleep.3494>.
31. Gossan N, Zeef L, Hensman J, Hughes A, Bateman JF, Rowley L et al. The circadian clock in murine chondrocytes regulates genes controlling key aspects of cartilage homeostasis. *Arthritis Rheum* 2013 Sep;65(9):2334–45. <https://doi.org/10.1002/art.38035>.
32. Dudek M, Meng QJ. Running on time: the role of circadian clocks in the musculoskeletal system. *Biochem J* 2014 Oct;463(1):1–8. <https://doi.org/10.1042/BJ20140700>.
33. McDearmon EL, Patel KN, Ko CH, Walisser JA, Schook AC, Chong JL et al. Dissecting the functions of the mammalian clock protein BMAL1 by tissue-specific rescue in mice. *Science* 2006 Nov;314(5803):1304–8. <https://doi.org/10.1126/science.1132430>.
34. Scheer FA, Hilton MF, Mantzoros CS, Shea SA. Adverse metabolic and cardiovascular consequences of circadian misalignment. *Proc Natl Acad Sci USA* 2009 Mar;106(11):4453–8. <https://doi.org/10.1073/pnas.0808180106>.
35. Guo B, Yang N, Borysiewicz E, Dudek M, Williams JL, Li J et al. Catabolic cytokines disrupt the circadian clock and the expression of clock-controlled genes in cartilage via an NFκB-dependent pathway. *Osteoarthritis Cartilage* 2015 Nov;23(11):1981–8. <https://doi.org/10.1016/j.joca.2015.02.020>.
36. Aspdren RM, Scheven BA, Hutchison JD. Osteoarthritis as a systemic disorder including stromal cell differentiation and lipid metabolism. *Lancet* 2001 Apr;357(9262):1118–20. [https://doi.org/10.1016/S0140-6736\(00\)04264-1](https://doi.org/10.1016/S0140-6736(00)04264-1).
37. Guo Y, Liu Y, Huang X, Rong Y, He M, Wang Y et al. The effects of shift work on sleeping quality, hypertension and diabetes in retired workers. *PLoS One* 2013 Aug;8(8):e71107. <https://doi.org/10.1371/journal.pone.0071107>.
38. Nea FM, Kearney J, Livingstone MB, Pourshahidi LK, Corish CA. Dietary and lifestyle habits and the associated health risks in shift workers. *Nutr Res Rev* 2015 Dec;28(2):143–66. <https://doi.org/10.1017/S095442241500013X>.
39. Bonnell EK, Huggins CE, Huggins CT, McCaffrey TA, Palermo C, Bonham MP. Influences on Dietary Choices during Day versus Night Shift in Shift Workers: A Mixed Methods Study. *Nutrients* 2017 Feb;9(3):9. <https://doi.org/10.3390/nu9030193>.
40. Lu B, Driban JB, Xu C, Lapane KL, McAlindon TE, Eaton CB. Dietary Fat Intake and Radiographic Progression of Knee Osteoarthritis: Data From the Osteoarthritis Initiative. *Arthritis Care Res (Hoboken)* 2017 Mar;69(3):368–75. <https://doi.org/10.1002/acr.22952>.
41. Dai Z, Niu J, Zhang Y, Jacques P, Felson DT. Dietary intake of fibre and risk of knee osteoarthritis in two US prospective cohorts. *Ann Rheum Dis* 2017 Aug;76(8):1411–9. <https://doi.org/10.1136/annrheumdis-2016-210810>.

Received for publication: 17 June 2019