Scand J Work Environ Health 1986;12(3):216-220

<https://doi.org/10.5271/sjweh.2155>

Issue date: Jun 1986

Chronic effects of dichloromethane on amino acids, glutathione and phosphoethanolamine in gerbil brain.

by [Briving C](#), [Hamberger A](#), [Kjellstrand P](#), [Rosengren L](#), [Karlsson JE](#), [Haglid KG](#)

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/3749836

This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Chronic effects of dichloromethane on amino acids, glutathione and phosphoethanolamine in gerbil brain

by Carin Briving, PhD,¹ Anders Hamberger, MD,¹ Per Kjellstrand, PhD,²
Lars Rosengren, MD,¹ JE Karlsson,¹ Kenneth G Haglid, MD¹

BRIVING C, HAMBERGER A, KJELLSTRAND P, ROSENGREN L, HAGLID KG. Chronic effects of dichloromethane on amino acids, glutathione and phosphoethanolamine in gerbil brain. *Scand J Work Environ Health* 12 (1986) 216–220. Mongolian gerbils were exposed to dichloromethane for three months by continuous inhalation at 210 ppm. Total free tissue amino acids, glutathione, and phosphoethanolamine were determined in the vermis posterior of the cerebellum and the frontal cerebral cortex. These two brain areas were chosen because humans occupationally exposed to dichloromethane have shown abnormalities in the electroencephalogram of the frontal part of the cerebral cortex. This study showed that long-term exposure of gerbils to dichloromethane (210 ppm) for three months leads to decreased levels of glutamate, γ -aminobutyric acid, and phosphoethanolamine in the frontal cerebral cortex, while glutamine and γ -aminobutyric acid are elevated in the posterior cerebellar vermis.

Key terms: experimental work, frontal cerebral cortex, long-term dichloromethane exposure, posterior cerebellar vermis.

Dichloromethane (methylene chloride, CH_2Cl_2) is used for a variety of purposes, as a solvent, a degreasing agent, an aerosol propellant, a paint remover, and as an extractant in the food industry. The adverse effects associated with dichloromethane exposure are primarily neurological and cardiovascular, while hepatotoxicity and nephrotoxicity are less pronounced. The effects on the central nervous system (CNS) are related to the anesthetic properties of the solvent. In cases of accidental human exposure, CNS symptoms include fatigue, nausea, headache, labored breathing, and unconsciousness. Behavioral alterations following acute exposure have also been described (29). Few experimental studies have concerned the chronic effects of dichloromethane on the CNS. Intermittent exposure of rats to dichloromethane (500–1 000 ppm) for two weeks causes neurochemical changes, including signs of increased protein breakdown in the cerebrum (26). We have previously shown that long-term exposure to moderate levels of trichloroethylene and perchloroethylene alters uptake and levels of amino acids in discrete brain regions of the gerbil (4).

This study reports on dichloromethane-induced effects on the nervous system of gerbils exposed for three months to the solvent at 210 ppm. The Swedish hygienic standard value is 70 ppm. The present work is part of an extended study on discrete brain areas, previously shown to be sensitive to chlorinated aliphatic solvents (8, 23). Concentrations of free amino

acids, glutathione, and phosphoethanolamine have been followed in the search for functional correlates to the recently reported alterations in cell composition of the CNS upon chronic exposure to dichloromethane (23).

Materials and methods

Animal exposure

The material consisted of four groups of Mongolian gerbils (eight animals in each), all the exposed animals being sex-matched littermates of the control animals. The animals were four months old when the exposure started. Two groups were exposed to 210 ppm of dichloromethane by continuous inhalation. Two control groups were exposed to air in identical chambers. The environmental temperature was $22 \pm 2^\circ\text{C}$. Commercial laboratory rat chow (Astra/Ewos, Sweden) and water were freely available.

The solvent (Uddeholm AB, Sweden) was vaporized at 60°C and diluted with filtered ($0.3 \mu\text{m}$) air. The concentration of the solvent in the inhalation chambers was monitored with a Miran 1A spectrophotometer equipped with a multipath gas cell, and it was kept within $\pm 10\%$ of the desired level. The exposure was interrupted for 1–2 h per week for changes of water, bedding material, and food (9).

After three weeks of exposure, eight exposed and eight control animals were removed from the inhalation chamber and immediately bled for the determination of carboxyhemoglobin percent saturation. After three months, eight exposed and their eight littermate controls were decapitated in the order control-exposed. The brains were rapidly removed and dissected as previously described (8). The vermis posterior of the cere-

¹ Institute of Neurobiology, University of Göteborg, S-400 33 Göteborg, Sweden.

² Institute of Zoophysiology, University of Lund, S-223 62 Lund, Sweden.

Reprint requests to: Dr KG Haglid, Institute of Neurobiology, University of Göteborg, POB 33 031, S-400 33 Göteborg, Sweden.

bellum (lobulus VI—X) and the frontal cerebral cortex were used for the determination of total free-tissue amino acids, glutathione, and phosphoethanolamine.

Tissue content of free amino acids

The dissected brain areas were extracted in 70 % ethanol for 24 h at +4°C. The resulting extracts were stored at -20°C until analyzed. The free amino acids were measured with reversed-phase liquid chromatography (Varian 5000 LC) after precolumn derivatization (10, 15). Briefly, a sample of the diluted tissue extract was mixed with borate buffer (pH 12), o-phthaldialdehyde, and mercaptoethanol. The mixture was injected onto a nucleosil 5 C₁₈ column after 1 min. The effluent was monitored fluorimetrically (FS 970 LC fluorimeter, Schoeffel Instrument). The mobile phase was a gradient from 30—100 % methanol in 0.04 M phosphate buffer, pH 5.2. Protein was determined according to Lowry et al (15).

Determination of carboxyhemoglobin

Four male littermate pairs of gerbils were continuously exposed to dichloromethane (210 ppm) for three weeks.

Four male littermate pairs of gerbils served as controls exposed to clean air. At the end of the exposure, the animals were removed from the inhalation chambers and immediately bled. The carboxyhemoglobin percent saturation was determined according to Collins et al (6) as modified by Lundh et al (16).

Statistical evaluation

The statistical analysis was performed with the aid of the Student's t-test for dependent series.

Results

Body and brain weights

The body and whole brain weights, as well as the weights of the dissected tissue pieces, did not differ significantly between the control and solvent-exposed groups (not shown).

Levels of free amino acids, proteins, glutathione and phosphoethanolamine

The protein concentrations per wet weight in different brain areas were not significantly altered after expo-

Figure 1. Content of different amino acids, glutathione, and phosphoethanolamine in the frontal cerebral cortex of gerbils immediately after continuous exposure to dichloromethane (210 ppm) for three months. Eight animals have been studied in each group, and the contents of amino acids have been estimated by the high-pressure liquid chromatographic technique. The results are given as the percentage of the control value. The statistical evaluation has been carried out with the aid of the Student's paired t-test (* $p < 0.05$). (X = control, Y = exposed, GLT = glutathione, ASP = aspartate, GLU = glutamate, SER = serine, GLN = glutamine, GLY = glycine, p-ET = phosphoethanolamine, TAU = taurine, ALA = alanine, GABA = γ -aminobutyric acid, a-ABA = α -aminobutyric acid, MET = methionine, VAL = valine)

VERMIS POSTERIOR

Figure 2. Amino acids, glutathione, and phosphoethanolamine content in the posterior cerebellar vermis after continuous exposure to dichloromethane (210 ppm) for three months. The result has been obtained from the same animals as in figure 1. Statistical evaluations are also the same as in figure 1 (* $p < 0.05$, ** $p < 0.01$).

sure to dichloromethane. The means of the free amino acids in the posterior vermis of control gerbils were somewhat lower (4) than in earlier studies on rat (10). Glutamate, phosphoethanolamine, and γ -aminobutyric acid (GABA) were significantly decreased in the frontal cerebral cortex after the exposure to dichloromethane (figure 1). A different pattern of alterations was found in the cerebellar posterior vermis, eg, increased levels of glutamine and γ -aminobutyric acid (figure 2).

Carboxyhemoglobin content

After three weeks of exposure at 210 ppm the carboxyhemoglobin percent saturation was significantly increased [11.5 (SE 0.4)] when compared to that of the controls [0.43 (SE 0.01)] ($p \leq 0.001$).

Discussion

Long-term exposure to dichloromethane induces effects which differ markedly in two brain areas. When gerbils are exposed to a higher dose of dichloromethane (350 ppm) for two months, long-lasting astrogliotic changes are induced and are traceable four months after exposure, especially in the frontal cerebral cortex (23).

After inhalation, dichloromethane is rapidly absorbed and distributed to the major body compartments. The absorbed solvent is mainly eliminated by pulmonary excretion of the unaltered parent compound. In the rat, about 50 % of the absorbed dichloromethane is metabolized before elimination at higher exposure concentrations, whereas essentially all may be metabolized at lower exposure concentrations (18). Two major metabolic pathways of biotransformation have been demonstrated in rat liver. The first pathway described follows oxidation by the microsomal cytochrome P-450 monooxygenases and ultimately yields carbon monoxide (12). The second follows conjugation to glutathione and leads to the formation of formaldehyde (1). Carbon monoxide and carbon dioxide are the main metabolites in exhaled air (18). A covalent binding of metabolites of ^{14}C -dichloromethane has been demonstrated in rat liver preparations (2).

One of the toxicologic hazards has been suggested to be brain dysfunction induced by carbon monoxide (3). The carboxyhemoglobin levels were significantly elevated among gerbils exposed to dichloromethane for three weeks. The relationship between the metabolism of dichloromethane and neurotoxicity of the compound is not fully understood. As both the major pathways of dichloromethane metabolism are present

to some extent in rat brain (1, 17), local metabolite formation could be of importance in explaining the neurotoxicity of the compound. One metabolite, formaldehyde, formed after conjugation to glutathione, is a well-recognized toxin which is known to bind covalently to tissue macromolecules (5). However, the levels of glutathione were unaffected in the exposed animals. Furthermore, the toxicologic hazards of dichloromethane in rodent brain could depend on the lipophilic properties of the compound or on its intrinsic toxicity. Dichloromethane is, however, less lipid-soluble than other chlorinated organic solvents with known neurotoxicologic potentials, such as trichloroethylene and tetrachloroethylene (25).

In man, certain brain areas are more vulnerable to chronic dichloromethane exposure, eg, the anterior parts of the cerebral cortex (28). Our previous studies also indicate that different cerebellar brain areas of the gerbil are less sensitive towards dichloromethane exposure (23). The observed decreased levels of glutamate, γ -aminobutyric acid, and phosphoethanolamine in the frontal cerebral cortex after dichloromethane exposure are of interest in comparison with our previous results indicating that the frontal cerebral cortex is especially vulnerable (23). It has recently been reported that the γ -aminobutyric acid contents are markedly reduced in cerebral and cerebellar cortex in patients with encephalopathy upon repeated hemodialysis (20). The significance of the effect on phosphoethanolamine is presently not clear. However, this primary amine is extensively released from CNS areas in experimental epilepsy, hypoglycemia, and ischemia (7, 13, 24). The increased levels of glutamine in the posterior cerebellar vermis could reflect an activation of astrocytic glia, as glutamine synthetase is localized exclusively in astrocytes (19). Alternatively, the increased glutamine could be secondary to the hyperammonemia caused by alterations in the liver due to dichloromethane. However, the liver toxicity of dichloromethane is considered to be low (21). The present results are in accordance with those of Honma et al (11), ie, increased CNS glutamine after one month's exposure to various levels of perchloroethylene (400–800 ppm) or in the posterior cerebellar vermis after longer exposure periods at lower concentrations (4). The increased levels of γ -aminobutyric acid in the posterior cerebellar vermis after dichloromethane exposure might indicate solvent interaction with inhibitory neuronal pathways with consequences for the cerebellar regulation of body movements. It has been shown that acute exposure to dichloromethane results in a depression of the vestibulo-oculomotor reflex (VOR) in rats (27). This effect could be due to interactions with the cerebellar inhibition of this reflex (22). Consequently, our findings indicate that brain regions in the gerbil show specific patterns with regard to alterations in amino acid levels, as well as phosphoethanolamine, upon dichloromethane exposure. It is of particular interest that the neurotransmitter group of amino acids

is affected. The decreased levels of γ -aminobutyric acid, glutamine, and phosphoethanolamine observed in the frontal cerebral cortex might be a more serious sign of interaction than the elevated levels of glutamine and γ -aminobutyric acid observed in the posterior cerebellar vermis, since this brain area seems to be less sensitive towards dichloromethane exposure (23).

Acknowledgments

This work was supported by the Swedish Work Environmental Fund (grant 84-0217 and 85-0585).

Ms E Åwall is gratefully acknowledged for her skillful technical assistance. We thank Ms G Grönstedt for typing the manuscript.

References

1. Ahmed AE, Anders MW. Metabolism of dihalomethanes to formaldehyde and inorganic halide: I *In vitro* studies. *Drug Metab Dispos* 4 (1976) 357–361.
2. Anders MW, Kubic VL, Ahmed AE. Metabolism of halogenated methanes and macromolecular binding. *J Environ Pathol Toxicol* 1 (1977) 117–124.
3. Barrowcliff DF, Knell AJ. Cerebral damage due to endogenous chronic carbon monoxide poisoning caused by exposure to methylene chloride. *J Soc Occup Med* 29 (1979) 12–14.
4. Briving C, Jacobson I, Hamberger A, Kjellstrand P, Rosengren L, Haglid KG. Chronic effects of perchloroethylene and trichloroethylene on the gerbil brain amino acids and glutathione. *Neurotoxicology* (in press).
5. Casanova-Schmitz M, Starr TB, Heck Hd'A. Differentiation between metabolic incorporation and covalent binding in the labeling of macromolecules in the rat nasal mucosa and bone marrow by inhaled (^{14}C)- and (^3H)formaldehyde. *Toxicol Appl Pharmacol* 76 (1984) 26–44.
6. Collins HA, Rodkey FL, O'Neal JD. Determination of carbon monoxide in blood by gas chromatography. *Clin Chem* 14 (1968) 162–171.
7. Hagberg H, Lehmann A, Sandberg M, Nyström B, Jacobson I, Hamberger A. Ischemia-induced shift of inhibitory and excitatory amino acids from intra- to extracellular compartments. *J Cereb Blood Flow Metab* (in press).
8. Haglid KG, Briving C, Hansson H-A, Kjellstrand P, Rosengren LE, Stavrou D, Swedin U, Wronski A. Trichloroethylene: Long-lasting changes in the brain after rehabilitation. *Neurotoxicology* 4 (1981) 659–673.
9. Haglid KG, Kjellstrand P, Rosengren L, Wronski A, Briving C. Effects of trichloroethylene inhalation on proteins of the gerbil brain. *Arch Toxicol* 43 (1980) 187–199.
10. Hamberger A, Jacobson I, Lindroth P, Mopper K, Nyström B, Sandberg M, Molin S-O, Svanberg U. Neuron-glia interactions in the biocynthesis and release of transmitter amino acids. *Adv Biochem Psychopharmacol* 29 (1981) 509–518.
11. Honma T, Hasegawa H, Sato M, Sudo A. Changes of free amino acid content in rat brain after exposure to trichloroethylene and tetrachloroethylene. *Ind Health* 18 (1980) 1–7.
12. Kubic VL, Anders MW. Metabolism of dihalomethanes to carbon monoxide: II *In vitro* studies. *Drug Metab Dispos* 3 (1975) 104–112.
13. Lehmann A, Hagberg H, Jacobson I, Hamberger A.

- Effects of status epilepticus on extracellular amino acids in hippocampus. *Brain Res* (in press).
14. Lindroth P, Mopper K. High-performance liquid chromatographic determination of subpicomole amounts of amino acids by precolumn fluorescence derivatization by o-phthaldialdehyde. *Anal Chem* 51 (1979) 1667—1674.
 15. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem* 193 (1951) 265—275.
 16. Lundh B, Johansson MB, Mercke C, Cavallin-Ståhl E. Enhancement of heme catabolism by caloric restriction in man. *Scand J Clin Lab Invest* 30 (1972) 421—427.
 17. Marietta MP, Vesell ES, Hartman RD, Weisz J, Dvorchik BH. Characterization of cytochrome P-450-dependent aminopyrine N-demethylase in rat brain: Comparison with hepatic aminopyrine N-demethylation. *J Pharmacol Exp Ther* 208 (1979) 271—279.
 18. McKenna MJ, Zempel JA, Braun WH. The pharmacokinetics of inhaled methylene chloride in rats. *Toxicol Appl Pharmacol* 65 (1982) 1—10.
 19. Norenberg MD, Martinez-Hernandez A. Fine structural localizations of glutamine synthetase in astrocytes of rat brain. *Brain Res* 161 (1979) 303—310.
 20. Perry TL, Young VW, Kish SJ, Foulks JG, Godolphin WJ, Sweeney VP. Neurochemical abnormalities in brains of renal failure patients treated by repeated hemodialysis. *J Neurochem* 4 (1985) 1043—1048.
 21. Plaa GL, Larsson RE. Relative nephrotoxic properties of chlorinated methane, ethane and ethylene derivatives in mice. *Toxicol Appl Pharmacol* 7 (1965) 37—44.
 22. Precht W. Vestibular mechanisms. *Ann Rev Neurosci* 2 (1979) 265—289.
 23. Rosengren LE, Kjellstrand P, Aurell A, Haglid KG. Irreversible effects of dichloromethane on the brain after long-term exposure: A quantitative study of DNA and the glial cell marker proteins S-100 and GFA. *Br J Ind Med* (in press).
 24. Sandberg M, Nyström B, Hamberger A. Metabolically derived aspartate-elevated extracellular levels *in vivo* in iodoacetate poisoning. *J Neurosci Res* 13 (1985) 489—495.
 25. Sato A, Nakajima T. Partition coefficients of some aromatic hydrocarbons and ketones in water, blood and oil. *Br J Ind Med* 36 (1979) 231—234.
 26. Savolainen H, Kurppa K, Pfäffli P, Kivistö H. Dose-related effects of dichloromethane on rat brain in short-term inhalation exposure. *Chem Biol Interact* 34 (1981) 315—322.
 27. Tham R, Bunnfors I, Eriksson B, Larsby B, Lindgren S, Ödkvist LM. Vestibulo-ocular disturbances in rats exposed to organic solvents. *Acta Pharmacol Toxicol* 54 (1984) 58—63.
 28. Von Hanke C, Ruppe K, Otto J. Untersuchungsergebnisse zur toxischen Wirkung von Dichlormetan bei Fussbodenlegern. *Z Gesamte Hyg* 20 (1974) 81—84.
 29. Winek CL, Collom WD, Esposito F. Accidental methylene chloride fatality. *Forensic Sci Int* 18 (1981) 165—168.

Received for publication: 7 November 1985