Scand J Work Environ Health 2012;38(6):577-581

<https://doi.org/10.5271/sjweh.3273>

Published online: 16 Jan 2012, Issue date: 01 Nov 2012

Professional driving and prolapsed lumbar intervertebral disc diagnosed by magnetic resonance imaging: a case-control study

by [Palmer KT](#), [Griffin M](#), [Ntani G](#), [Shambrook J](#), [McNee P](#), [Sampson M](#), [Harris EC](#), [Coggon D](#)

Affiliation: MRC Lifecourse Epidemiology Unit, Southampton General Hospital, Southampton SO16 6YD, UK. ktp@mrc.soton.ac.uk

Refers to the following texts of the Journal: [1998;24\(5\):358-366](#)
[2008;34\(5\):364-373](#)

Key terms: [back pain](#); [case-control study](#); [disc pathology](#); [driving](#); [low-back pain](#); [magnetic resonance imaging](#); [MRI](#); [PID](#); [professional driving](#); [prolapsed lumbar intervertebral disc](#); [whole-body vibration](#)

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/22249859

This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Professional driving and prolapsed lumbar intervertebral disc diagnosed by magnetic resonance imaging: a case–control study

by Keith T Palmer, DM,¹ Michael Griffin, PhD,² Georgia Ntani, MSc,¹ James Shambrook, FRCR,³ Philip McNee,⁴ Madeleine Sampson, FRCR,³ E Clare Harris, PhD,¹ David Coggon, PhD¹

Palmer KT, Griffin M, Ntani G, Shambrook J, McNee P, Sampson M, Harris EC, Coggon D. Professional driving and prolapsed lumbar intervertebral disc diagnosed by magnetic resonance imaging: a case–control study. *Scand J Work Environ Health*. 2012;38(6):577–581. doi:10.5271/sjweh.3273

Objectives The aim of this study was to investigate whether whole-body vibration (WBV) is associated with prolapsed lumbar intervertebral disc (PID) and nerve root entrapment among patients with low-back pain (LBP) undergoing magnetic resonance imaging (MRI).

Methods A consecutive series of patients referred for lumbar MRI because of LBP were compared with controls X-rayed for other reasons. Subjects were questioned about occupational activities loading the spine, psychosocial factors, driving, personal characteristics, mental health, and certain beliefs about LBP. Exposure to WBV was assessed by six measures, including weekly duration of professional driving, hours driven at a spell, and current 8-hour daily equivalent root-mean-square acceleration A(8). Cases were sub-classified according to whether or not PID/nerve root entrapment was present. Associations with WBV were examined separately for cases with and without these MRI findings, with adjustment for age, sex, and other potential confounders.

Results Altogether 237 cases and 820 controls were studied, including 183 professional drivers and 176 cases with PID and/or nerve root entrapment. Risks associated with WBV tended to be lower for LBP with PID/nerve root entrapment but somewhat higher for risks of LBP without these abnormalities. However, associations with the six metrics of exposure were all weak and not statistically significant. Neither exposure–response relationships nor increased risk of PID/nerve root entrapment from professional driving or exposure at an A(8) above the European Union daily exposure action level were found.

Conclusions WBV may be a cause of LBP but it was not associated with PID or nerve root entrapment in this study.

Key terms back pain; disc pathology; low-back pain; MRI; PID; whole-body vibration.

Many studies have found an association of whole-body vibration (WBV) with low-back pain (LBP) and sciatica, and some indicate an increased risk of prolapsed intervertebral disc (PID) specifically (1–9). However, the latter have seldom confirmed diagnosis beyond self-report, so it remains uncertain whether WBV is a cause of PID. The increased use of magnetic resonance imaging (MRI) offers new opportunities to distinguish cases with PID and nerve root entrapment. In the present paper, we report a case–control study of MRI-investigated LBP,

which examines the association of PID and nerve root entrapment with occupational exposure to WBV.

Methods

The main methods have been described elsewhere (9). In brief, the study population comprised working-aged adults resident in the area served by a public hospital.

¹ MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK.

² Institute of Sound and Vibration Research, University of Southampton, Southampton, UK.

³ Southampton University Hospitals NHS Trust, Southampton, UK.

⁴ Centre for Defence Imaging, Portsmouth, UK.

Correspondence to: Professor Keith Palmer, MRC Lifecourse Epidemiology Unit, Southampton General Hospital, Southampton SO16 6YD, UK. [E-mail: ktp@mrc.soton.ac.uk]

Cases were a consecutive series referred to the radiology department at that hospital or to two local private hospitals for MRI of the lumbar spine during 2003–2006. Controls were investigated by radiography while attending the public hospital's emergency department over 2003–2006. They fulfilled the same residency requirement and were group-matched to cases by sex and age.

Participants completed a questionnaire on occupational history, work activities (digging, lifting, trunk bending/twisting), professional driving, and exposure to WBV (vehicle types, duration, intensity), demographic characteristics, somatizing tendency (10), low mood (11), consulting behavior (12), fear-avoidance beliefs (13), and beliefs regarding the work-relatedness of LBP. Cases were asked about their LBP symptoms and disability. Exposure to WBV in their latest job was assessed by six metrics: (i) professional driving (>1 hour/day); (ii) professional driving (>3 hours consecutively); (iii) weekly hours driven for the vehicle most used; (iv) weekly hours driven for all vehicles (none, <16 , ≥ 16); (v) maximum root-mean-square (rms) acceleration of any vehicle (0, -0.5 , ≥ 0.6 ms^{-2} rms) and (vi) A(8) rms ($<$ or ≥ 0.5 ms^{-2} rms [the action level in the European Union (EU) Physical Agents (vibration) Directive (14)]. Metrics (v) and (vi) were derived from driving times and imputed vibration magnitudes of vehicles (9, 15).

Blinded to work history, radiologists assessed MRI scans using a repeatable standardized protocol (16). Images were graded at three spinal levels (L3/L4, L4/L5, L5/S1) for: (i) disc herniation (protrusion, herniation, or disc sequestration); and/or (ii) nerve root entrapment (displacement or compression).

Analysis focused on cases whose latest LBP episode (that since last pain-free for ≥ 1 month) began in their current/most recent job and controls who gave a current or recent job history. Associations were summarized by odds ratios (OR) with 95% confidence intervals (95% CI). Separate logistic regression models were constructed for each metric of WBV. All were adjusted for age and sex as matching variables. Additional adjustment was made for well-recognized risk factors for LBP previously associated with case presentation in the sample (9) (see table 1).

Finally, as some cases came from the private hospitals, potential selection bias was explored by repeating analyses without the private cases.

Results

In all, 237 cases and 820 controls were analyzed. [Response rates, reasons for exclusion, control diagnoses, and patient characteristics were identical to those cited previously (9), save for 15 excluded cases with missing

scans.] The groups were well matched by age, sex, and interval from starting their latest job. Cases with relevant MRI findings ($N=174$) were similar to other cases in their duration of current LBP, disability in the past four weeks, and most personal and occupational risk factors.

The sample included 183 professional drivers (130 of whom drove cars). Table 1 shows associations with WBV adjusted for age and sex, and then additionally for other risk factors for LBP. Partially and fully adjusted models produced similar risk estimates. In cases without PID/nerve root entrapment, slightly elevated risks (OR 1.4) were found for driving for ≥ 3 hours continuously and driving in total >0 – <15 hours/week; while, in cases with PID/nerve root entrapment, reduced risks were sometimes found. However, no associations were significant at the 5% level and overall risks from being a professional driver or from an A(8) above the EU action level were close to unity, irrespective of MRI findings.

Table 2 presents findings after excluding private cases. Again, no associations were significant at the 5% level; but the inverse association of some WBV metrics with relevant MRI pathology was more marked, and risks were more than doubled among scan-negative cases who drove for >3 hours continuously and drivers whose A(8) exceeded the EU action level. To explore whether lower risks from driving among those with PID/nerve root entrapment could represent selection out of driving in previous employments, we assessed the proportion of subjects who had had a driving job for ≥ 1 year but had left such work when scanned: 55% of scan-positive cases had left such work versus 57% of scan-negative cases and 51% of controls.

Discussion

Previously in this sample, we found no important associations between WBV and LBP severe enough to occasion lumbar MRI imaging but strong associations with several physical and psychosocial risk factors for LBP (9). This analysis, stratified by MRI evidence of PID and/or nerve root entrapment, indicated no elevated risks from WBV for these pathologies but somewhat elevated risks for LBP in their absence. One possible interpretation is that WBV (or sitting while driving) increases the risk of LBP but not the risk of PID or nerve root entrapment.

Our study had certain limitations. Participation was incomplete, but this would bias analysis only if the relation between driving and MRI appearance differed by response status, which seems unlikely. Exposure ascertainment was retrospective and selection bias might arise if drivers with LBP avoided future WBV exposure. However, the proportion of cases holding a previous

Table 1. Relationship of driving and whole body vibration to magnetic resonance imaging (MRI)-diagnosed disc herniation and/or nerve root compression or displacement. Separate models were constructed for each vibration variable. [OR=odds ratio; 95% CI=95% confidence interval; BMI=body mass index; rms=root mean square.]

Vibration exposure	Controls (N=820)		Cases with disc herniation and/or nerve root compression/displacement on MRI (N=176)						Cases without disc herniation and/or nerve root compression/displacement on MRI (N=61)					
	N	%	N	%	OR	95% CI ^a	OR	95% CI ^b	N	%	OR	95% CI ^a	OR	95% CI ^b
Professional driving (≥1 hours/day)														
No	677	82.6	147	83.5	1	..	1	..	50	82.0	1	..	1	..
Yes	143	17.4	29	16.5	0.8	0.5–1.3	0.8	0.5–1.3	11	18.0	1.1	0.5–2.2	1.1	0.5–2.2
Professional driving (≥3 hours/time)														
No	765	93.3	163	92.6	1	..	1	..	56	91.8	1	..	1	..
Yes	55	6.7	13	7.4	1.0	0.5–1.8	0.9	0.5–1.8	5	8.2	1.3	0.5–3.5	1.4	0.5–4.1
Average hours driven/week for commonest source														
None	677	82.6	147	83.5	1	..	1	..	50	82.0	1	..	1	..
<16	67	8.2	19	10.8	1.2	0.7–2.0	1.2	0.7–2.2	5	8.2	1.0	0.4–2.7	1.1	0.4–3.1
≥16	76	9.3	10	5.7	0.5	0.2–1.0	0.5	0.2–1.0	6	9.8	1.1	0.4–2.9	1.0	0.4–2.7
Total hours driven/ week, all sources														
Not a regular driver	677	82.6	147	83.5	1	..	1	..	50	82.0	1	..	1	..
<15	54	6.6	17	9.7	1.3	0.7–2.3	1.4	0.7–2.5	5	8.2	1.3	0.5–3.3	1.4	0.5–3.9
≥15	89	10.9	12	6.8	0.5	0.3–1.0	0.5	0.2–1.0	6	9.8	1.0	0.4–2.4	0.8	0.3–2.2
Max rms of any machine (ms ⁻²)														
Not a regular driver	677	82.6	147	83.5	1	..	1	..	50	82.0	1	..	1	..
0.5	87	10.6	21	11.9	1.0	0.6–1.7	1.0	0.6–1.7	6	9.8	1.0	0.4–2.4	1.0	0.4–2.5
≥0.6	56	6.8	8	4.6	0.5	0.2–1.2	0.5	0.2–1.2	5	8.2	1.3	0.5–3.5	1.1	0.4–3.3
Current rms A(8) (ms ⁻²)														
<0.5	697	85.0	149	84.7	1	..	1	..	52	85.3	1	..	1	..
≥0.5	123	15.0	27	15.3	0.9	0.5–1.4	0.9	0.5–1.5	9	14.8	1.0	0.5–2.2	0.9	0.4–2.2

^a Adjusted for age (in 3 bands) and sex.^b Adjusted for age (in 3 bands), sex, BMI (in 3 bands), somatizing tendency (in 3 bands), SF-36 mental health score (in 3 bands), smoking status (current, ex-, never smoker), propensity to consult over back pain (in three bands), fear avoidance beliefs (in three bands), belief in work as a cause of back pain (in three bands), occupational digging and/or lifting (in two bands) and occupational bending and/or twisting (in two bands).

driving job for >1 year but no longer was similar to controls, and similar by scan result. Non-differential errors in exposure assessment could bias risk estimates to the null for complex metrics relying on recalled exposure times and imputed estimates of vibration magnitude. However, several exposure measures were straightforward (eg, driving for >1 hour/week); evidence exists that occupational drivers can assess driving times accurately (17); we focused on recent exposures for which reporting should be more dependable and provided an aide-memoire checklist of sources. In any event, such errors are unlikely to explain differences of association between MRI-defined subgroups.

Nor is it likely that confounding explains such differences. The assessed risk factors for LBP were similarly frequent in the two MRI-defined case subgroups and adjustment changed risk estimates only marginally.

More telling is the source of recruitment. Risks became doubled only in a sensitivity analysis that

excluded private hospital cases. This strengthened associations between WBV and LBP in the absence of PID/nerve root entrapment and, if anything, reduced estimates of risk for LBP associated with these pathologies. Analyses in table 1 had the advantage that they included both private and non-private cases from the study population, while controls should have provided representative exposure information on those at risk of becoming cases (emergencies were treated only at the public hospital). However, risks from WBV could be underestimated if private cases were investigated by MRI at a lower-severity threshold. Conversely, risk estimates in table 2 would be underestimates if privately treated patients were less exposed to WBV. Importantly, neither analysis provided support for a positive association of WBV with PID or nerve root entrapment.

Associations between WBV and LBP have been observed frequently. Radiating leg pain and PID have also been associated with WBV (1–9); but only a few

Table 2. Relationship of driving and whole body vibration to MRI-diagnosed disc herniation and/or nerve root compression or displacement, with analysis restricted to subjects from the public hospital. Separate models were constructed for each vibration variable. [OR=odds ratio; 95% CI=95% confidence interval; rms=root mean square.]

Vibration exposure	Cases with disc herniation and/or nerve root compression/displacement on MRI (N=91)				Cases without disc herniation and/or nerve root compression/displacement on MRI (N=36)			
	N	%	OR ^a	95% CI	N	%	OR ^a	95% CI
Professional driving (≥ 1 hrs/day)								
No	78	85.7	1	..	28	77.8	1	..
Yes	13	14.3	0.7	0.3–1.3	8	22.2	1.7	0.7–4.2
Professional driving (≥ 3 hrs/time)								
No	84	92.3	1	..	32	88.9	1	..
Yes	7	7.7	0.9	0.4–2.3	4	11.1	2.3	0.7–7.8
Average hrs driven/week for commonest source								
None	78	85.7	1	..	28	77.8	1	..
<16	9	9.9	1.1	0.5–2.4	3	8.33	1.3	0.4–4.7
≥ 16	4	4.4	0.3	0.1–1.0	5	13.9	2.1	0.6–6.6
Total hours driven/week, all sources								
Not a regular driver	78	85.7	1	..	28	77.8	1	..
<15	7	7.7	1.1	0.5–2.6	3	8.3	1.6	0.4–6.0
≥ 15	6	6.6	0.4	0.2–1.1	5	13.9	1.7	0.5–5.3
Max rms of any machine (ms ⁻²)								
Not a regular driver	78	85.7	1.0	1.0–1.0	28	77.8	1	..
0.5	7	7.7	0.7	0.3–1.6	5	13.9	1.9	0.7–5.5
≥ 0.6	6	6.6	0.7	0.3–1.7	3	8.3	1.3	0.3–5.2
Current rms A(8) (ms ⁻²)								
<0.5	80	87.9	1	..	28	77.8	1	..
≥ 0.5	11	12.1	0.6	0.3–1.3	8	22.2	2.1	0.8–5.5

^a Adjusted for age (in 3 bands), sex, BMI (in 3 bands), somatizing tendency (in 3 bands), SF36 mental health score (in 3 bands), smoking status (current, ex, never smoker), propensity to consult over back pain (in three bands), fear avoidance beliefs (in three bands), belief in work as a cause of back pain (in three bands), occupational digging and/or lifting (in two bands) and occupational bending and/or twisting (in two bands).

studies have focused on PID diagnosed objectively. In these, the findings have been less clear cut. One case–control study reported an elevated risk of PID among those who sat most of the time in a motor vehicle [relative risk (RR) 2.75] (7). However, surgically confirmed cases constituted only ten matched pairs and, in this subset, the RR for driving was lower (1.3). A larger study, in which PID was surgically corroborated in half of cases, found slightly elevated risks among drivers of certain cars, but not among truck or van drivers (8). A study comparing machine drivers and carpenters with office workers, found higher risks of MRI-diagnosed lumbar anterior disc bulges among the machine drivers (OR elevated 3–4 fold), but reduced risks of posterior bulging (OR 0.6–0.9) (18); disc bulges and machine driving were independently associated with sciatic pain (19). A study of monozygotic twins discordant for occupational driving history found no relationship between driving and MRI-diagnosed lumbar disc bulges or herniations (20). A case–control study of MRI- or computed tomography-confirmed acute PID showed a trend in risk with lumbar spinal force but no significant trend with lifetime hours of WBV (21). A survey of Indian tractor driver farmers found they were no more likely than non-farmers to have disc bulging on MRI, despite

having more work-related backache (22).

The hypothesis that WBV causes LBP (and sciatica) but is not an important cause of PID is thus compatible with the broad literature on drivers' symptoms and objectively diagnosed PID. However, it requires corroboration in further studies that verify diagnoses objectively over a more extreme range of exposure circumstances; this does not preclude the possibility that WBV causes other degenerative disc pathology.

Acknowledgments

The European Commission supported this research under the Quality of Life and Management of Living Resources program, project no. QLK4-2002-02650 (VIBRISKS).

References

1. Boshuizen HC, Bongers PM, Hulshof CTJ. Self-reported back pain in forklift truck and freight-container drivers exposed to

- whole-body vibration. *Spine*. 1992;17:59–65. <http://dx.doi.org/10.1097/00007632-199201000-00010>.
2. Boshuizen HC, Bongers PM, Hulshof CTJ. Long-term sick leave and disability pensioning due to back disorders of tractor drivers exposed to whole-body vibration. *Int Arch Occup Environ Health*. 1990;62:117–22. <http://dx.doi.org/10.1007/BF00383587>.
 3. Bovenzi M, Betta A. Low-back disorders in agricultural tractor drivers exposed to whole-body vibration and postural stress. *Appl Ergon*. 1994;25:231–41. [http://dx.doi.org/10.1016/0003-6870\(94\)90004-3](http://dx.doi.org/10.1016/0003-6870(94)90004-3).
 4. Bongers PM, Boshuizen HC, Hulshof CTJ. Self-reported back pain in drivers of wheel-loaders. In: *Back disorders and whole-body vibration at work*. Amsterdam: Academisch Proefschrift, Universiteit van Amsterdam; 1990. p207–20.
 5. Boshuizen HC, Bongers PM, Hulshof CTJ. Self-reported back pain and exposure to whole-body vibration in helicopter pilots. In: *Back disorders and whole-body vibration at work*. Amsterdam: Academisch Proefschrift, Universiteit van Amsterdam; 1990. p191–206.
 6. Hannerz H, Tuchsén. Hospital admissions among male drivers in Denmark. *Occup Environ Med*. 2001;58:253–60. <http://dx.doi.org/10.1136/oem.58.4.253>.
 7. Kelsey JL, Hardy RL. Driving of motor vehicles as a risk factor for acute herniated lumbar intervertebral disc. *Am J Epidemiol*. 1975;102:63–73.
 8. Kelsey JL, Githens PB, O'Conner T, Weil U, Calogero JA, Holford T, et al. Acute prolapsed lumbar intervertebral disc: an epidemiologic study with special reference to driving automobiles and cigarette smoking. *Spine*. 1984;6:608–13. <http://dx.doi.org/10.1097/00007632-198409000-00012>.
 9. Palmer KT, Harris EC, Griffin MJ, Bennett J, Reading I, Sampson M, Coggon D. Case-control study of low-back pain presenting for MRI, with special relation to whole-body vibration. *Scand J Work Environ Health*. 2008;34:364–73. <http://dx.doi.org/10.5271/sjweh.1282>.
 10. Derogatis LR, Melisaratos N. The brief symptom inventory: an introductory report. *Psychological Medicine*. 1983;13:595–605. <http://dx.doi.org/10.1017/S0033291700048017>.
 11. Ware JE. SF-36 Health Status Questionnaire. Boston, MA: Institute for the Improvement of Medical Care and Health, New England Medical Center Hospital, Quality Quest Inc; 1989.
 12. Palmer KT, Calnan M, Wainwright D, O'Neill C, Winterbottom A, Watkins C, et al. Upper limb pain in primary care: Health beliefs, somatic distress, consulting and patient satisfaction. *Family Practice*. 2006;23:609–17. <http://dx.doi.org/10.1093/fampra/cml047>.
 13. Waddell G, Newton M, Henderson I, Somerville D, Main CJ. A Fear-Avoidance Beliefs Questionnaire and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain*. 1993;52:157–68. [http://dx.doi.org/10.1016/0304-3959\(93\)90127-B](http://dx.doi.org/10.1016/0304-3959(93)90127-B).
 14. The European Parliament, The Council of the European Union. Directive 2002/44/EC of 25 June 2002 on the minimum health and safety requirements regarding the exposure of workers to the risks arising from physical agents (vibration). *Off J Eur Communities*. 2002;L177:13–9.
 15. Lundström R. Risks of occupational vibration exposures (VIBRISKS): FP5 Project No. QLK4-2002-02650 January 2003 to December 2006: annex to the final technical report —appendix 2 [Internet]. Brussels: European Commission, Quality of Life and Management of Living Resources Programme; 2007 [cited 3 July 2008]. Available from: http://www.vibrisks.soton.ac.uk/reports/Annex12%20UMUH%20WP4_1%20WBV%20Protocol%20310107.pdf.
 16. Shambrook J, McNee P, Harris EC, Kim M, Palmer KT, Coggon D. Clinical presentation of low-back pain and association with risk factors according to findings on Magnetic Resonance Imaging. *Pain*. 2011;152:1659–65. <http://dx.doi.org/10.1016/j.pain.2011.03.011>.
 17. Palmer KT, Haward B, Griffin MJ, Bendall H, Coggon D. Validity of self-reported occupational exposures to hand-transmitted and whole-body vibration. *Occup Environ Med*. 2000;57:237–41. <http://dx.doi.org/10.1136/oem.57.4.237>.
 18. Luoma K, Riihimäki H, Raininko R, Luukkonen R, Lamminen A, Viikari-Juntura E. Lumbar disc degeneration in relation to occupation. *Scand J Work Environ Health*. 1998; 24:358–66. <http://dx.doi.org/10.5271/sjweh.356>.
 19. Luoma K, Riihimäki H, Luukkonen R, Raininko R, Lamminen A, Viikari-Juntura E, et al. Low back pain in relation to lumbar disc degeneration. *Spine*. 2000;25:487–92. <http://dx.doi.org/10.1097/00007632-200002150-00016>.
 20. Battie MC, Videman T, Gibbons L, Manninen H, Gill K, Pope M, Kaprio J. Occupational driving and lumbar disc degeneration: a case-control study. *Lancet*. 2002;360: 1369–74. [http://dx.doi.org/10.1016/S0140-6736\(02\)11399-7](http://dx.doi.org/10.1016/S0140-6736(02)11399-7).
 21. Seidler A, Bolm-Audorff U, Siol T, Henkel N, Fuchs C, Schug H, et al. Occupational risk factors for symptomatic lumbar disc herniation; a case-control study. *Occup Environ Med*. 2003;60:821–30. <http://dx.doi.org/10.1136/oem.60.11.821>.
 22. Kumar A, Varghese M, Mohan D, Mahajan P, Gulati P, Kale S. Effect of whole-body vibration on the low back: a study of tractor-driving farmers in North India. *Spine*. 1999;24:2506–15. <http://dx.doi.org/10.1097/00007632-199912010-00013>.

Received for publication: 17 October 2011