Scand J Work Environ Health 2014;40(3):230-234

<https://doi.org/10.5271/sjweh.3412>

Published online: 19 Dec 2013, Issue date: 01 May 2014

A meta-analysis of health effects of randomized controlled worksite interventions: Does social stratification matter?

by [Montano D](#), [Hoven H](#), [Siegrist J](#)

This study emphasizes the role of social stratification of occupational groups targeted for worksite interventions. This meta-analysis also updates and enriches previous studies on worksite interventions by considering only randomized controlled interventions. Significant effects on musculoskeletal symptoms, fruit and vegetable consumption, perceived stress, and body mass index were confirmed and/or reported.

Affiliation: "Work Stress Research", Life-Science-Center, Faculty of Medicine, Düsseldorf University, Merowingerplatz 1a, D-40225 Düsseldorf, Germany. diego.montano@med.uni-duesseldorf.de

Refers to the following text of the Journal: [2009;35\(1\):7-18](#)

The following article refers to this text: [2019;45\(6\):546-559](#)

Key terms: [body mass index](#); [body mass index](#); [fruit and vegetables consumption](#); [intervention](#); [job stress](#); [meta-analysis](#); [musculoskeletal disease](#); [musculoskeletal disorder](#); [musculoskeletal symptom](#); [randomized controlled intervention](#); [randomized controlled worksite intervention](#); [review](#); [social stratification](#); [socioeconomic position](#); [socioeconomic status](#); [worksite intervention](#)

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/24788850

Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](#).

This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

A meta-analysis of health effects of randomized controlled worksite interventions: Does social stratification matter?

by Diego Montano, PhD,¹ Hanno Hoven, MA,^{1,2} Johannes Siegrist, PhD¹

Montano D, Hoven H, Siegrist J. A meta-analysis of health effects of randomized controlled worksite interventions: Does social stratification matter? *Scand J Work Environ Health*. 2014;40(3):230–234. doi:10.5271/sjweh.3412

Objectives The aim of this review was to assess what types of socioeconomic positions (SEP) are being considered in randomized controlled intervention studies and estimate the moderation of SEP in workplace intervention effects on body mass index (BMI), fruit and vegetable consumption, musculoskeletal symptoms, and job stress.

Methods A meta-analysis of randomized controlled workplace interventions was undertaken. Studies were classified by participants' SEP. The overall standardized mean difference (SMD) for each outcome was estimated with random-effects models. Additionally, a random-effects model with SEP as moderating variable was calculated in order to assess intervention effect modification (EM).

Results This review covers 36 studies. Altogether 40 reports of intervention effects were considered. The overall mean differences in the models, without SEP as moderating variable, were significant for all outcomes. BMI, self-reported musculoskeletal symptoms, and self-reported job stress decreased [SMD -0.16, 95% confidence interval (95% CI) -0.29– -0.02, SMD -0.32, 95% CI -0.51– -0.14, and SMD -0.37, 95% CI -0.71– -0.04, respectively], whereas daily consumption of fruit and vegetables increased (SMD 0.12, 95% CI 0.01–0.22). There were no statistically significant differences between occupational classes for the health outcomes considered (SMD -0.102, 95% CI -0.264–0.060, EM -0.141, 95% CI -0.406–0.125; SMD 0.117, 95% CI -0.049–0.282, EM 0.000, 95% CI -0.230–0.231; SMD -0.301, 95% CI -0.494– -0.107, EM -0.369, 95% CI -1.169–0.430; and SMD -0.200, 95% CI -0.524–0.124, EM -0.598, 95% CI -1.208–0.012, respectively).

Conclusions Workplace interventions can achieve small positive effects on major health outcomes. We could not confirm whether these effects are moderated by occupational class.

Key terms body mass index; fruit and vegetables consumption; job stress; musculoskeletal disease; musculoskeletal disorder; musculoskeletal symptom; review; socioeconomic position; socioeconomic status.

Intervention effectiveness research is considered a priority area in occupational health (1). The physical and psychosocial workplace environment offers a crucial setting for improving the health of employed populations (2, 3). Although beneficial for all employees, the need to improve health at work is unequally distributed among employed populations. This is due to the well-established fact of a social gradient of health across the whole of a society. With each step up in the social hierarchy, the better the chances of good health (4, 5). Occupational groups with lower qualifications and occupational positions generally suffer from poorer health than those in higher positions (6, 7). As the quality of work and employment follows the same pattern of social

distribution as health, leaving those in lower socioeconomic positions (SEP) in more stressful jobs (8, 9), there is a clear need to prioritize worksite interventions among occupational groups with lower socioeconomic standing and their specific occupational exposures (10).

However, when reviewing worksite intervention trials, this double burden of poorer quality of work and higher risk of ill-health among less-privileged occupational groups is rarely defined as a major challenge. Although several systematic reviews and meta-analyses have been conducted on the effects of workplace interventions (11–25), they fail to deal explicitly with the social gradient of health and poor quality of work.

In this review, we report the results of a meta-

¹ Senior professorship “Work Stress Research”, Faculty of Medicine, Düsseldorf University, Düsseldorf, Germany.

² Institute of Medical Sociology, Faculty of Medicine, Düsseldorf University, Düsseldorf, Germany.

Correspondence to: Diego Montano, Senior professorship “Work Stress Research”, Life-Science-Center, Faculty of Medicine, Düsseldorf University, Merowingerplatz 1a, D-40225, Düsseldorf, Germany. [E-mail: diego.montano@med.uni-duesseldorf.de]

analysis of randomized controlled worksite interventions and their effects on health and well-being with a special focus on the social stratification of occupational target groups. Two research questions are addressed: (i) What is the social distribution of occupations included in randomized controlled worksite interventions? (ii) Do effect sizes differ between higher and lower SEP groups? The first objective aims to assess whether lower socioeconomic groups are being prioritized in intervention studies. The second addresses whether intervention effects are being moderated by socioeconomic class.

Methods

The extensive study protocol, search strategy, sensitivity analysis, quality of evidence, and detailed description of studies included in this systematic review are available as supplementary material (http://www.sjweh.fi/data_repository.php). This section is, therefore, restricted to the general selection criteria of the intervention studies and the methodology used to answer the two research questions mentioned above.

In order to achieve robust estimates of effect sizes (26), we aimed to identify only those studies which can be regarded as homogenous with respect to design and outcomes. First, only randomized controlled interventions at the workplace were included to allow the calculation of standardized mean differences (SMD). In order to assess the effect of interventions on healthy working populations, only primary interventions were selected (16). Furthermore, to decrease the small sample bias, studies had to provide a population size >80 (ie, ≥ 40 employees in each group) in order to be included. Second, we selected those studies whose outcome variables were highly similar or identical. The outcome variables included: (i) body mass index (BMI); (ii) fruit and vegetable consumption (measured in almost all studies by administering a validated screening instrument for daily servings of fruit and vegetables); (iii) musculoskeletal symptoms (self-reported upper-extremity symptoms, shoulder and neck and/or back pain symptoms); and (iv) perceived stress (assessment of operationalizations of self-reported levels of stress).

In line with our first research question, we applied the Erikson-Goldthorp-Portocarrero (EGP) occupational class categories to identify the SEP of the employees in each sample (27). However, due to a lack of specific information regarding the sample characteristics in many studies, only two major groups of SEP categories were defined: (i) EGP classes I–III: higher and lower service occupations, routine clerical, and sales occupations and (ii) EGP classes VI–VII: skilled, and semi- or unskilled manual occupations.

In dealing with differences in the scales of the instruments used for the measurement of outcome variables, SMD were estimated. For publications reporting means and standard errors in the intervention and control groups (26 effect sizes) Hedge's g (ie, the SMD), was calculated and corrected for small sample size (28). For the remaining publications, appropriate transformation formulae of effect sizes were utilized (29–32).

Due to large heterogeneity, random-effects models were calculated by the restricted maximum-likelihood estimator (33). All models were calculated using the R-package metaphor (34).

Results

After checking for duplicates, the manual search and queries in the databases generated a total of 18 132 initial records. After applying the study selection criteria (briefly mentioned in the Methods section), 104 studies were selected for full analysis. Finally, 36 randomized controlled interventions and 40 reported measurements were considered for the meta-analyses.

Regarding the first research question, the frequencies of reported outcomes within the different EGP occupational classes are displayed in figure 1. The number of interventions among occupations belonging to EGP classes I–III is twice as high as the number of studies conducted among classes VI–VII. In particular, this difference is due to a larger number of studies dealing with musculoskeletal disorders and perceived stress within EGP classes I–III.

Concerning the second research question, we estimated eight random-effects models that are reproduced in table 1. The first four models did not take SEP into account, whereas the remaining models include SEP as a moderating variable. The intercept is the estimate of the overall effect size (Hedge's g), and the beta coefficient is an estimate of the effect differences between EGP classes. Workplace interventions in models 1–4 resulted in significantly decreased BMI values, increased intake of fruit and vegetables, decreased level of self-reported musculoskeletal symptoms, and decreased perceived stress. The inclusion of SEP as moderating variable in models 5–8 resulted in a slight decrease of the intercepts [ie, the overall effect sizes, except for musculoskeletal symptoms, even though the proportion of heterogeneity (column I2) remained large]. The results do not show a moderation effect exerted by SEP, as can be seen from the large confidence intervals of the beta coefficients.

Figure 1. Number of outcomes reported in the 36 intervention studies of the meta-analysis within each Erikson-Goldthorpe-Portocarrero (EGP) occupational class. [BMI=body mass index; MSS=musculoskeletal symptoms]

Table 1. Eight random-effects models of effect sizes (Hedges *g*) with and without socioeconomic position (SEP) as moderating variable. The intercept correspond to the overall effect size (Hedges *g*) for each outcome. The reference category in the moderation models 5–8 consists of Erikson-Goldthorpe-Portocarrero (EGP) classes I–III. The coefficient beta corresponds to the average intervention effect differences between EGP classes. [BMI=body mass index; MSS=musculoskeletal symptoms; 95% CI=95% confidence interval]

Overall intervention effects					Overall intervention effects with SEP moderation				
Model	Outcome	Coefficient	95% CI	I2	Model	Outcome	Coefficient	95% CI	I2
1	BMI				5	BMI			
	Intercept	-0.155	-0.286– -0.024	69		Intercept	-0.102	-0.264– -0.060	67
						Beta	-0.141	-0.406– -0.125	
2	Fruit/vegetables				6	Fruit/vegetables			
	Intercept	0.116	0.009– 0.224	79		Intercept	0.117	-0.049– 0.282	79
						Beta	0.000	-0.230– 0.231	
3	MSS				7	MSS			
	Intercept	-0.322	-0.509– -0.136	91		Intercept	-0.301	-0.494– -0.107	91
						Beta	-0.369	-1.169– -0.430	
4	Perceived stress				8	Perceived stress			
	Intercept	-0.373	-0.709– -0.038	89		Intercept	-0.200	-0.524– -0.124	84
						Beta	-0.598	-1.208– -0.012	

Discussion

We addressed two questions in this review: (i) What is the social distribution of occupations included in randomized controlled worksite interventions? and (ii) Do effect sizes differ between higher and lower SEP groups? Concerning the first question, the results indicate that more than twice as many intervention trials were conducted among higher than lower skilled occupational groups (EGP classes I–III versus VI–VII), especially for musculoskeletal disorders and perceived stress. Therefore, prioritization of intervention activities according to socially unequal needs has not been a major concern in this area of research. We were unable to find

a clear answer to the second research question given the lack of statistical power in our study, which prevented the estimation of occupational class differences.

However, even though the random-effects models controlling for the moderating effect of the EGP classes do not demonstrate statistically significant intervention effects, the corresponding regression coefficients in table 1 suggest lower intervention effects in EGP classes VI–VII in comparison to I–III. For instance, this would imply that for BMI there might be, on average, a 14.1% intervention effect reduction for occupational classes VI–VII.

Finally, some limitations of this analysis need to be mentioned. First, relevant studies may have been omitted as a consequence of the keywords selected in

the search strategy. Second, concerning the moderating effect of occupational class, all corresponding random-effects models had low degrees of freedom, especially for the model of perceived stress with only seven studies available, whereby a consistent estimation of mediation effects and corresponding confidence intervals cannot be satisfactorily attained. Third, it should be kept in mind that the sample description in terms of occupational class was very heterogeneous, thus limiting the validity of the applied EPG classification.

In conclusion, the meta-analysis of 36 randomized controlled worksite intervention studies revealed small, but significant effects on four relevant health outcomes: weight reduction, healthy nutrition, reduced musculoskeletal symptoms, and lower levels of perceived stress. As only few studies were conducted among employees with lower socioeconomic standing, the challenge of reducing work-related health inequalities by targeting health-promoting activities at occupational groups with high needs remains largely unmet. Due to lack of statistical power, our study could not confirm a moderation of intervention effects by occupational class. Nonetheless, future research should aim at bridging worksite intervention research with scientific inquiry into social determinants of health.

Acknowledgments

The research leading to these results was done within the framework of the DRIVERS project (www.health-gradient.eu) coordinated by EuroHealthNet, which received funding from the European Community (FP7 2007-2013) under grant agreement no 278350. J Siegrist and D Montano were additionally supported by a Senior Professorship Grant from the Faculty of Medicine, University of Duesseldorf, Germany.

References

- LaMontagne AD, Youngstrom RA, Lewiton M, et al. An Exposure Prevention Rating Method for Intervention Needs Assessment and Effectiveness Evaluation. *Appl Occup Environ Hyg*. 2003;18(7):523–34. <http://dx.doi.org/10.1080/10473220301458>.
- World Health Organization (WHO). The Bangkok Charter for Health Promotion in a Globalized World. Paris: WHO; 2005.
- Wanless D, editors. Securing Good Health for the Whole Population [report]. London: Department of Health (UK); 2004.
- Marmot M, Wilkinson R, editors. Social Determinants of Health. Oxford: Oxford University Press; 2006.
- Mackenbach J. Health Inequalities: Europe in Profile [report]. Available from: http://ec.europa.eu/health/ph_determinants/socio_economics/documents/ev_060302_rd06_en.pdf.
- Schnall PL, Dobson M, Rosskam E, editors. Unhealthy Work: Causes, Consequences, Cures. Amityville NY: Baywood Publishing Co; 2009.
- Siegrist J, Theorell T. Socioeconomic position and health: the role of work and employment. In: Siegrist J, Marmot M, editors. Social Inequalities in Health. New Evidence and Policy Implications. Oxford: Oxford University Press; 2006. Available from: <http://dx.doi.org/10.1093/acprof:oso/9780198568162.001.0001>.
- Johnson J. The Growing Imbalance: Class, Work, and Health in an Era of Increasing Inequality. In: Schnall PL, Dobson M, Rosskam E, editors. Unhealthy Work: Causes, Consequences, Cures. Amityville NY: Baywood Publishing Co; 2009.
- Lynch JW, Kaplan GA. Socioeconomic position. In: Berkman LF, Kawachi I, editors. Social Epidemiology. New York: Oxford Univ Press; 2000. p13–35.
- Marmot M, Allen J, Bell R, et al. WHO European review of social determinants of health and the health divide. *The Lancet*. 2012;380(9846):1011–29. [http://dx.doi.org/10.1016/S0140-6736\(12\)61228-8](http://dx.doi.org/10.1016/S0140-6736(12)61228-8).
- Bambra C, Whitehead M, Sowden A, et al. Shifting Schedules: The Health Effects of Reorganizing Shift Work. *Am J Prev Med*. 2008;34(5):427–34. <http://dx.doi.org/10.1016/j.amepre.2007.12.023>.
- Joyce K, Pabayo R, Critchley JA, et al. Flexible working conditions and their effects on employee health and wellbeing. *Cochrane Database Syst Rev*; 2010.
- Benedict MA, Arterburn D. Worksite-based weight loss programs: a systematic review of recent literature. *Am J Health Promot*. 2008;22(6):408–16. <http://dx.doi.org/10.4278/ajhp.22.6.408>.
- Ni Mhurchu C, Aston LM, Jebb SA. Effects of worksite health promotion interventions on employee diets: a systematic review. *BMC Public Health*. 2010;10:62. <http://dx.doi.org/10.1186/1471-2458-10-62>.
- Anderson LM, Quinn TA, Glanz K, et al. The Effectiveness of Worksite Nutrition and Physical Activity Interventions for Controlling Employee Overweight and Obesity. A Systematic Review. *Am J Prev Med*. 2009;37(4):340–57. <http://dx.doi.org/10.1016/j.amepre.2009.07.003>.
- LaMontagne AD, Keegel T, Louie A, et al. A Systematic Review of the Job-stress Intervention Evaluation Literature, 1990-2005. *Int J Occup Environ Health*. 2007;13(3):268–80.
- Odeen M, Magnussen LH, Maeland S, et al.: Systematic review of active workplace interventions to reduce sickness absence. *Occup Med (Lond)*. 2013;63(1):7–16. <http://dx.doi.org/10.1093/occmed/kqs198>.
- Corbiere M, Shen J, Rouleau M, et al. A systematic review of preventive interventions regarding mental health issues in organizations. *Work*. 2009;33(1):81–116.
- Kennedy CA, Amick III BC, Dennerlein JT, et al. Systematic Review of the Role of Occupational Health and Safety

- Interventions in the Prevention of Upper Extremity Musculoskeletal Symptoms, Signs, Disorders, Injuries, Claims and Lost Time. *J Occup Rehabil*. 2010;20(2):127–62. <http://dx.doi.org/10.1007/s10926-009-9211-2>.
20. Conn VS, Hafdahl AR, Cooper PS, et al. Meta-Analysis of Workplace Physical Activity Interventions. *Am J Prev Med*. 2009;37(4):330–9. <http://dx.doi.org/10.1016/j.amepre.2009.06.008>.
21. Hutchinson AD, Wilson C. Improving nutrition and physical activity in the workplace: A meta-analysis of intervention studies. *Health Promot Int*. 2012;27(2):238–49. <http://dx.doi.org/10.1093/heapro/dar035>.
22. Portnoy DB, Scott-Sheldon LAJ, Johnson BT, et al. Computer-delivered interventions for health promotion and behavioral risk reduction: A meta-analysis of 75 randomized controlled trials, 1988–2007. *Prev Med*. 2008;47(1):3–16. <http://dx.doi.org/10.1016/j.jypmed.2008.02.014>.
23. Martin A, Sanderson K, Cocker F. Meta-analysis of the effects of health promotion intervention in the workplace on depression and anxiety symptoms. *Scand J Work Environ Health*. 2009;35(1):7–18. <http://dx.doi.org/10.5271/sjweh.1295>.
24. Richardson KM, Rothstein HR. Effects of occupational stress management intervention programs: a meta-analysis. *J Occup Health Psychol*. 2008;13(1):69–93. <http://dx.doi.org/10.1037/1076-8998.13.1.69>.
25. David AR, Szamoskozi S. A meta-analytical study on the effects of cognitive behavioral techniques for reducing distress in organizations. *Journal of Cognitive and Behavioral Psychotherapies*. 2011;11(2):221–36.
26. Ioannidis JPA. Why most published research findings are false. *PLoS Med*. 2005;2(8):e124. <http://dx.doi.org/10.1371/journal.pmed.0020124>.
27. Ganzeboom HBG, Treiman DJ. Internationally Comparable Measures of Occupational Status for the 1988 International Standard Classification of Occupations. *Social Science Research*. 1996;25(3):201–39. <http://dx.doi.org/10.1006/ssre.1996.0010>.
28. Hedges LV. Distribution Theory for Glass's Estimator of Effect Size and Related Estimators. *J Educ Behav Stat*. 1981;6(2):107–28. <http://dx.doi.org/10.3102/10769986006002107>.
29. Chinn S. A simple method for converting an odds ratio to effect size for use in meta-analysis. *Stat Med*. 2000;19(22):3127–31. [http://dx.doi.org/10.1002/1097-0258\(20001130\)19:22<3127::AID-SIM784>3.0.CO;2-M](http://dx.doi.org/10.1002/1097-0258(20001130)19:22<3127::AID-SIM784>3.0.CO;2-M).
30. Rosnow RL, Rosenthal R. Computing Contrasts, Effect Sizes, and Countermills on Other People's Published Data: General Procedures for Research Consumers. *Psychol Meth*. 1996;1(4):331–40. <http://dx.doi.org/10.1037/1082-989X.1.4.331>.
31. Rosnow RL, Rosenthal R, Rubin DB. Contrasts and Correlations in Effect-Size Estimation. *Psych Sci*. 2000;11(6):446–53. <http://dx.doi.org/10.1111/1467-9280.00287>.
32. Hedges LV, Olkin I. Statistical methods for meta-analysis. San Diego, CA: Academic Press; 1985.
33. Viechtbauer W. Bias and Efficiency of Meta-Analytic Variance Estimators in the Random-Effects Model. *J Educ Behav Stat*. 2005;30(3):261–93. <http://dx.doi.org/10.3102/10769986030003261>.
34. Viechtbauer W. Conducting Meta-Analyses in R with the metafor Package. *J Stat Softw*. 2010;36(3):1–48.

Received for publication: 22 May 2013