Scand J Work Environ Health 2016;42(2):103-124

<https://doi.org/10.5271/sjweh.3533>

Published online: 24 Nov 2015, Issue date: 01 Mar 2016

Systematic review of biochemical biomarkers for neck and upper-extremity musculoskeletal disorders

by Gold JE, Hallman DM, Hellström F, Björklund M, Crenshaw AG, Djupsjöbacka M, Heiden M, Mathiassen SE, Pilgian G, Barbe MF

A statistically significant association between the musculoskeletal disorders (MSD) and biomarkers studied was demonstrated in 82% of sufficient quality articles. Evidence suggested that: collagen repair marker (TIMP-1) is decreased in fibroproliferative disorders; serotonin (5-HT) is increased in trapezius myalgia; triglycerides are increased in various MSD. Limitations in the articles include possible selection bias, confounding, spectrum effect, and insufficient attention to comorbid conditions.

Affiliation: Centre for Musculoskeletal Research (CBF), Department of Occupational and Public Health Sciences, University of Gävle, SE - 801 76 Gävle, Sweden. Judith.Gold@hig.se

Refers to the following texts of the Journal: 1999;25(6):564-568
2000;26(4):292-298 2008;34(2):133-141 1992;18 suppl 2:85-87

Key terms: biochemical biomarker; biomarker; collagen degradation; collagen repair; inflammation; lipids; muscle injury; muscle metabolite; musculoskeletal disorder; neck; pain; serotonin; systematic review; tendon injury; triglycerides; upper-extremity; upper-extremity musculoskeletal disorder

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/26599377

Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](#).

This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

Systematic review of biochemical biomarkers for neck and upper-extremity musculoskeletal disorders

by Judith E Gold, ScD,¹ David M Hallman, PhD,¹ Fredrik Hellström, PhD,¹ Martin Björklund, PhD,^{1,4} Albert G Crenshaw, PhD,¹ Mats Djupsjobacka, PhD,¹ Marina Heiden, PhD,¹ Svend Erik Mathiassen, PhD,¹ George Piligian, MD,² Mary F Barbe, PhD³

Gold JE, Hallman DM, Hellström F, Björklund M, Crenshaw AG, Djupsjobacka M, Heiden M, Mathiassen SE, Piligian G, Barbe MF. Systematic review of biochemical biomarkers for neck and upper-extremity musculoskeletal disorders. *Scand J Work Environ Health*. 2016;42(2):103–124. doi:10.5271/sjweh.3533

Objective This study systematically summarizes biochemical biomarker research in non-traumatic musculoskeletal disorders (MSD). Two research questions guided the review: (i) Are there biochemical markers associated with neck and upper-extremity MSD? and (ii) Are there biochemical markers associated with the severity of neck and upper-extremity MSD?

Methods A literature search was conducted in PubMed and SCOPUS, and 87 studies met primary inclusion criteria. Following a quality screen, data were extracted from 44 articles of sufficient quality.

Results Most of the 87 studies were cross-sectional and utilized convenience samples of patients as both cases and controls. A response rate was explicitly stated in only 11 (13%) studies. Less than half of the studies controlled for potential confounding through restriction or in the analysis. Most sufficient-quality studies were conducted in older populations (mean age in one or more analysis group >50 years). In sufficient-quality articles, 82% demonstrated at least one statistically significant association between the MSD and biomarker(s) studied. Evidence suggested that: (i) the collagen-repair marker TIMP-1 is decreased in fibroproliferative disorders, (ii) 5-HT (serotonin) is increased in trapezius myalgia, and (iii) triglycerides are increased in a variety of MSD. Only 5 studies showed an association between a biochemical marker and MSD severity.

Conclusion While some MSD biomarkers were identified, limitations in the articles examined included possible selection bias, confounding, spectrum effect (potentially heterogeneous biomarker associations in populations according to symptom severity or duration), and insufficient attention to comorbid conditions. A list of recommendations for future studies is provided.

Key terms collagen degradation; collagen repair; inflammation; lipid; muscle injury; muscle metabolites; pain; serotonin; tendon injury; triglyceride.

Upper-extremity musculoskeletal disorders (MSD) affecting the soft tissues of the neck, shoulders, arms, and hands are prevalent worldwide (1–4) and are common causes of absence from work and disability (5). MSD impose a large economic burden on society (1, 3, 4, 6–11).

Options for screening, surveillance, and diagnosis of upper-extremity MSD are limited and most depend on

symptoms, which are by definition self-reported, or on functional ability. A need for improved diagnostic and screening methods, especially objective techniques, has been identified (12, 13). According to the US Institute of Medicine (IOM), a biomarker is “a characteristic that is objectively measured and evaluated as an indicator of normal biologic processes, pathogenic processes, or pharmacologic responses to a therapeutic intervention”

¹ Centre for Musculoskeletal Research, Department of Occupational and Public Health Sciences, University of Gävle, Gävle, Sweden.

² Department of Occupational Medicine, Epidemiology and Prevention, North Shore-Long Island Jewish Health System, Great Neck, NY, USA.

³ Department of Anatomy and Cell Biology, Temple University Medical School, Philadelphia, PA, USA.

⁴ Department of Community Medicine and Rehabilitation, Physiotherapy, Umeå University, Umeå, Sweden.

Correspondence to: Judith E Gold, Centre for Musculoskeletal Research (CBF), Department of Occupational and Public Health Sciences, University of Gävle, SE – 801 76 Gävle, Sweden. [E-mail: Judith.Gold@hig.se]

(14). In addition to improving diagnosis and screening methods (15), valid and reliable MSD biomarkers could aid in evaluating treatments and workplace interventions. Finally, such biomarkers could aid in elucidating MSD pathomechanisms.

The US National Research Council (NRC) has distinguished between three classes of biomarkers: exposure, effect (disease), and susceptibility (16). A biomarker of effect may be defined as “any change that is qualitatively or quantitatively predictive of health impairment or potential impairment...” (16). In this review, we focus on biomarkers of effect, which – through their window on pathophysiological disease processes – could potentially be utilized to stage the severity of a MSD.

Previous biomarker reviews

The potential for MSD biomarkers to detect subclinical disease and monitor MSD severity was discussed as early as 1992 by Mastin and colleagues (17). They recommended the use of animal models for biomarker research with subsequent validation in human populations. Since then, only one review focusing exclusively on biomarkers of MSD in humans has been published (18). Saxton (2000) (18) recommended longitudinal studies in which initially asymptomatic persons were enrolled, and then inflammatory substances and proteins specific to skeletal muscle be examined in relation to low force, highly repetitive tasks. In the intervening years, literature on biochemical markers of MSD has grown, although very few longitudinal studies have been performed. Recent reviews have highlighted biochemical alterations associated with the pathogenesis of particular MSD in humans or animal models (19–22). However, a systemic summary of the current state of MSD biomarker research in humans is needed. Therefore, we sought to conduct a comprehensive assessment of the current state of neck and upper-extremity MSD biochemical biomarker research, using two research questions: (i) Are there biochemical markers associated with neck and upper-extremity MSD? (ii) Are there biochemical markers associated with the severity of neck and upper-extremity MSD?

Methods

Review team and overview

Our review team consisted of ten researchers with expertise in epidemiologic, intervention, and experimental studies within the field of work-related MSD research. In a first step, we formulated research questions and defined principal concepts of the review. Then we devel-

oped a search strategy and search terms and conducted a search using PubMed and Scopus databases, with results pooled with articles identified from our own records. Thereafter, papers were screened based on pre-defined criteria (supplementary tables S1–S3, www.sjweh.fi/data_repository.php) using a two-step procedure of primary (title and abstract) and secondary (quality) screens. Summary tables were created from papers of sufficient quality, and evidence was synthesized with respect to the two research questions.

A consensus process was used throughout. The review team agreed on the definitions for neck and upper-extremity MSD and biochemical markers, operationalized severity, and developed specific inclusion/exclusion criteria.

Neck and upper-extremity MSD were defined as musculoskeletal symptoms or clinical diagnoses in the neck, shoulder, arm, wrist, or hand. These included both specific and non-specific conditions related to tendons, muscles, ligaments, blood vessels or nerves (23). We focused on MSD that occur in a work-related context (24).

Biochemical marker was defined as a substance obtained through biological fluids or tissues that may be evaluated as an indicator of normal biological or pathogenic processes. Such markers could be derived from serum, blood, urine, synovial fluid, saliva, muscle interstitium or through tissue biopsy samples. Genetic markers such as RNA, DNA or their constituents were considered as beyond the scope of this review.

Severity was operationalized as encompassing cross-sectional and longitudinal differences in symptom severity, or differences in extent of disorder in same anatomical segment (eg, size of rotator cuff tear).

Inclusion criteria

This review was limited to studies on humans and non-traumatic MSD of the neck and upper extremities. As categorized by Boocock et al (23), potential biomarkers were examined for the following specific MSD : flexor-extensor peritendinitis or tenosynovitis of forearm-wrist region, de Quervain’s disease (tenosynovitis of sheath/tunnel surrounding tendons controlling thumb movement), carpal or radial tunnel syndrome, Guyon (ulnar) canal syndrome, cubital tunnel syndrome (ulnar nerve entrapment at the elbow), epicondylitis (lateral and medial), secondary Raynaud’s phenomenon associated with vibration exposure, thoracic outlet syndrome, rotator cuff syndrome/shoulder tendonitis, and shoulder capsulitis. Non-specific diffuse forearm pain and other non-specific localized neck and upper-extremity MSD were included, as were upper-extremity pain in specific regions, including “shoulder pain”, “neck pain”, “neck/shoulder pain”, “elbow pain”, “arm pain”, and “hand pain”. Other specific MSD included are listed in Boocock

et al (table 7) (23), although cervico-brachial fibromyalgia status post-whiplash and joint-related conditions were excluded. Data from immunohistochemistry studies in which antibody assays were used [similar to an enzyme-linked immunosorbent assay (ELISA)] were included, and if an *in-vitro* study using cells from muscle or tendon biopsies included data from unstimulated cells, then that latter data were included. We included articles that fulfilled our inclusion criteria, even if some parts of the study were consistent with the exclusion criteria; however, only results in compliance with our criteria were taken into account.

Exclusion criteria are summarized in supplementary table S1 (www.sjweh.fi/data_repository.php).

Literature search

The search was first conducted in PubMed and combined with relevant articles identified from personal files. A second search for papers that cited the sufficient-quality articles was conducted using Scopus. See supplementary table S2 (www.sjweh.fi/data_repository.php) for an overview of the search strategy; figure 1 illustrates the search strategy and selection procedure. PubMed search terms included both MESH terms and keywords selected for two categories: upper-extremity MSD and biomarkers. We only included search terms which contributed more than two papers to the search. This elimination of papers did not result in having a lack of reviewable studies in any of the diagnostic categories listed in Boocock et al (23).

We identified 15 relevant articles and used these to refine the search and test its sensitivity. The search was restricted to peer-reviewed studies on humans, adults (>18 years of age), and English language text published within the last 25 years (June 4, 1988–June 4, 2013). See appendix 1 (www.sjweh.fi/data_repository.php) for the search string. The final PubMed search resulted in 2241 articles; four additional eligible papers were added from personal files (for a total of 2245 examined). A primary and secondary screening process was implemented (see below and supplementary tables S1–S3). After identifying the sufficient-quality articles, a second search was conducted in Scopus to identify relevant papers citing these articles. The main reason for the additional search in Scopus was that recently published articles in PubMed may not yet have been assigned MESH terms. Also, as our search strategy in PubMed may have missed some studies of relevance from other databases, the Scopus search was intended to reduce such potential bias. Pilot testing of forms and evaluation criteria was conducted at each stage of the review process (primary screen, secondary screen, data extraction) to assure clarity in review instruments and to limit bias among the reviewers.

Primary screen-selection of relevant articles

After importing PubMed records into systematic review software (EPPI-reviewer4 v4.3.4, EPPI-Centre at the Social Science Research Unit at the Institute of Education, University of London, UK), two independent reviewers conducted a primary screen assessing each title and abstract for eligibility based on the criteria in supplementary table S1 (www.sjweh.fi/data_repository.php). The fulltext article was read if necessary. A “yes” answer on any of the questions resulted in exclusion of the article. Results were compared between reviewers, and a consensus agreement was reached (see below). The same procedure was repeated for articles from personal files and additional articles identified using Scopus.

Secondary screen-quality assessment & data extraction

All articles passing the primary screen were scored for quality. The articles were randomly allocated to five different clusters, and each of five reviewers was assigned randomly to two clusters. The quality screen procedure was as follows: each article was assessed by two independent reviewers, scores were compared between reviewers, and a consensus agreement was reached after discussing disagreements. If an agreement could not be reached, a third person from the review team was involved. The questions used for the quality assessment are listed in supplementary table S3 (www.sjweh.fi/data_repository.php). These questions were derived from reporting guidelines for research studies and checklists assessing study quality in health-related research (25–29). The scoring system contained 17 items, where each item could be scored as either “yes” (1 point), “unknown or not applicable” (0), or “no” (0). Scores were summed for each paper (ranging from 0–17). To determine a criterion for article data extraction, a sample of five papers were scored by two reviewers. Based on our judgment of quality, we determined that articles scoring $\geq 65\%$ of the maximum (11/17) should be included for data extraction. These articles were labeled as “sufficient quality”. Regardless of quality, articles scoring “no” on question 15 were excluded from data extraction. These included papers with less appropriate statistical analysis methods, such as multiple comparisons without adjustment, and modeling without accounting for repeated measures. See appendix 2 for data extraction items (www.sjweh.fi/data_repository.php).

Research synthesis

Next, we evaluated the number of sufficient-quality articles identifying a particular biomarker or class of biomarkers in its association with MSD in conjunction with a best evidence synthesis approach (30). Consider-

Figure 1. Flow chart demonstrating literature search.

ing the heterogeneity of biomarkers and MSD studied, it was not possible to conduct a meta-analysis.

We did not expect a universal biomarker or biomarkers for the variety of MSD examined. Results for biomarkers were categorized by MSD in the tables, and summarized by physiological process in the text (eg, collagen degradation and repair). This dichotomy in presentation was chosen as we anticipate that our target audience is more familiar with MSD diagnoses. It should be noted that MSD diagnoses do not necessarily reflect underlying pathophysiology, and common physiological processes may underlie multiple MSD.

Physiological processes for which there were five or more relevant papers are presented. A biomarker was considered to belong to a category based on known physiological processes or, if several categories were appropriate, on the processes presented in the original article. In experiments with biomarker values across several time points, we considered any statistical comparison between cases and controls (or tests of association with MSD severity) cases) to be relevant.

An association between the biomarker and MSD in three or more sufficient-quality studies (and at most one sufficient-quality study with a null finding) was regarded as evidence that a substance could serve as a MSD biomarker. On a per biomarker basis, the smallest percentage of papers that demonstrated an association was 75%.

Results

Of the 2241 articles identified in the primary PubMed review (figure 1), plus the 4 papers added from authors'

personal files, 89 met the inclusion criteria for secondary (quality) screening. We excluded 27 papers upon viewing the entire article and 31 of the remaining 62 articles met the sufficient-quality criteria score of ≥ 11 . Next, we searched Scopus for articles that cited these 31 papers. Of 817 citations initially found in Scopus, 31 were determined to be non-duplicates and of relevance through a title and abstract review; 7 were eliminated at the secondary screening stage upon viewing the entire article. Hence, 24 articles were scored with the secondary screen. Of these, 15 were scored at ≥ 11 , adding to the 31 similarly scored articles found in the initial PubMed search. These 46 studies were regarded as being of sufficient quality to merit data extraction, except for 2 that used less appropriate statistical analysis methods. Hence, data was extracted from 44 articles.

Secondary screen-quality assessment overview

Of all of the studies meeting the criteria for quality scoring ($N=86$; 62 from PubMed and 24 from Scopus), the majority had clearly defined aims, biomarkers, MSD, and results (supplementary table S4, www.sjweh.fi/data_repository.php). Most studies had a cross-sectional design and utilized convenience sampling of patients, for both cases and referents. The response rate was explicitly stated or easily inferred in only 11 studies. Of these, the response rate to an invitation to participate was $>65\%$ in 9 (31–39). Less than half of the 86 studies controlled for confounding factors through restriction of subjects to a particular age or gender, or through adjustment in the analysis. Thirteen studies (15%) explicitly stated that biomarker analyses were blinded to case conditions, and 27 (31%) used less appropriate statistical analysis methods.

Data extraction from sufficient-quality studies

The majority of included studies examined neck/shoulder disorders and symptoms, with shoulder disorders, Dupuytren's contracture, and hand-arm vibration syndrome (HAVS) also well represented (supplementary table S5, www.sjweh.fi/data_repository.php). Please reference appendix 3 for definitions and descriptions of abbreviations and biomarker analyses presented in the tables. Four studies examined upper-extremity MSD or symptoms without further specifying a particular body region (38, 40–42). Referent groups ranged from healthy controls to patient groups with conditions presumed to be sufficiently unrelated to the biomarker(s) under investigation. Only one was a longitudinal (non-treatment) study (37).

Many studies were conducted in older populations with at least one analysis group ≥ 50 years of age (33–35, 43–60). The range of symptom duration varied substantially within and across studies. Of the 15 studies that listed an average or range of symptom duration, nearly half involved participants with MSD symptoms of >70 months (59, 61–65). Three reported extremely wide-ranging MSD durations (49, 50, 66) (66: 7–120 months; 49: 7–360 months; 50: 9–288 months). Five studied patients with an average symptom duration of <1 year (33, 40, 41, 54, 56).

Are there biochemical biomarkers associated with neck and upper-extremity MSD?

The majority of studies demonstrated an association between at least one biomarker and the MSD examined (table 1). Only 8 of the 44 studies reported insignificant findings throughout (33, 38, 46, 48, 51, 58, 67, 68). Descriptions are provided below for the categories of biomarkers examined in ≥ 5 studies. See appendix 3 for definitions of abbreviations and biomarker analyses (www.sjweh.fi/data_repository.php).

Inflammatory. Fourteen studies examined biochemical mediators and markers related to inflammatory processes (33, 34, 40–42, 59, 61–63, 67, 69–72). Seven found significant associations between inflammatory markers and MSD (34, 40, 42, 59, 62, 63, 69), while seven did not (33, 41, 61, 67, 70–72). Where associations were found, all had increased levels of inflammatory substances in MSD cases or with increased MSD severity, except for two studies in which reduced levels of IL-8 and IL-18 were observed in cases (41, 59).

A great breadth of inflammatory substances was examined. Six studies examining inflammatory indices in serum or blood observed increases: (i) serum macrophage chemotactic factor-1 (CCL2/MCP-1) and macrophage inflammatory protein-1 alpha (CCL3/MIP-1 α) in cultured whole blood cells with chronic neck pain (69); (ii) serum

C-reactive protein (CRP) with bilateral shoulder pain in females (34); (iii) serum CRP, IL-1 β , IL-6, and tumor necrosis factor alpha (TNF- α) with proximal and distal upper-extremity MSD of <3 months duration (40); (iv) serum sIL-1RII with specific and non-specific upper-extremity MSD (42); (v) serum soluble intercellular adhesion molecule-1 (sICAM-1) with HAVS (59); and (vi) blood polymorphonuclear cells (PMN) cell activity with HAVS (63). Associations between serum inflammatory cytokines, mediators and biomarkers were not consistent across reviewed studies. No association was observed between serum CRP and rotator cuff tendinitis, adhesive capsulitis, or a mix of MSD of <1 month duration (33, 34, 41). No association was found between serum TNF- α levels (an acute inflammatory cytokine) and chronic neck pain (69), between serum IL-1 and IL-6 levels (a pleiotropic cytokine and myokine) and late stage carpal tunnel syndrome (70), or between four acute phase cytokines (IL-1 α , IL-1 β , IL-6, TNF- α) and a mix of MSD as each were below detectable levels (41). Results were less mixed when using local sampling methods, eg, assays of analytes from muscle extracellular fluid obtained using microdialysis. No association was observed between several inflammation-related cytokines in the muscle interstitium and trapezius myalgia (71, 72). No association was observed for muscle interstitial prostaglandin E2 (PGE2) and trapezius myalgia and chronic muscle pain in the neck/shoulder (61, 67), although it should be noted that PGE2's role is not as a principal driver of inflammation but more as a mediator involved in immune responses. In muscle biopsies, Hadrevi et al (62) found increased heat shock 70 kDa (HSP 70) protein and alpha-1-antitrypsin, proteins involved in the inflammatory response.

In summary, results were largely inconsistent for the inflammatory substances examined in more than one study. The exception was a null finding for PGE2, which was observed in trapezius myalgia, neck/shoulder pain, and carpal tunnel syndrome.

Collagen degradation and repair. Seven studies examined biomarkers of collagen degradation and repair, a category that included matrix metalloproteinases (MMP, collagen degradation markers), tissue inhibitors of matrix metalloproteinases (TIMP), cross-linked carboxyterminal telopeptide of type I collagen (ICTP, a collagen degradation marker), and procollagen I intact N-terminal propeptide (PINP, a collagen type I synthesis marker) (44, 49–51, 56, 72, 73) (table 1). All but one study (51) found a significant association between at least one biomarker of collagen degradation and repair and the MSD under investigation.

Three studies found greater concentrations of collagen degradation markers in MSD cases than controls, specifically increased MMP in synovial fluid and biopsies from subjects with torn rotator cuff tendons and palmar fascia

Table 1. Are there biochemical biomarkers associated with neck and upper-extremity musculoskeletal disorders (MSD)?

MSD classification and diagnosis	Author	Major results (case-control comparison)	Conclusion
Neck disorders and symptoms			
Chronic & recurring neck pain	Teodorczyk-Injeyan et al (69)	CCL2/MCP-1, CCL3/MIP-1 α , NO ₂ ⁻ , TNF- α : in serum and unstimulated whole blood cell cultures. CCL2/MCP-1: Serum: NS; cultured cells: cases > controls, P <0.01. CCL3/MIP-1 α : cultured cells: cases > controls, P <0.01. NO ₂ ⁻ : Serum: cases > controls, P <0.01. TNF- α : Serum & cultured cells: NS.	Yes ↑ NO ₂ ⁻ in serum of cases ↑ CCL2, CCL3 in unstimulated cultured whole blood cells of cases
Shoulder disorders & symptoms			
Rotator cuff tear	Abboud et al (43)	HDL, LDL, Total cholesterol (TC), triglycerides in serum. TC: cases > controls, P=0.02 (hypercholesterolemia in 47/84 cases with TC > 240 mg/dL). Triglycerides: cases > controls, P=0.02. LDL: cases > controls, P=0.03. HDL: NS in cases vs. controls, P=0.10.	Yes ↑ TC, Triglycerides and LDL in serum of cases
Rotator cuff tear	Castagna et al (51)	MMP-1,2,3,8,9,10,13, & TIMP-1,2 in biopsies from torn and intact supraspinatus tendon and intact subscapularis tendon from same patient. MMP 1,2,3,8,9,10,13: NS comparing torn tendon area & intact areas. TIMP 1,2: NS comparing torn tendon area & intact areas.	No Seemingly intact parts of injured tendon can present with altered cellular metabolism.
Rotator cuff tear	Lakemeier et al (45)	VEGF in biopsies of long head of biceps tendon. VEGF: Partial thickness rotator cuff tears > controls, P<0.05. VEGF: Full thickness rotator cuff tears > controls, P<0.05.	Yes ↑ VEGF in torn tendons (partial or full)
Rotator cuff tear	Lakemeier et al (44)	MMP-1,3,9 in biopsies of rotator cuff tendon. MMP-1 & 9 increased in each Patte grade vs. controls, P<0.05. MMP-3 decreased in each Patte grade vs. controls, P<0.05.	Yes ↑ MMP-1 & 9 in torn tendon ↓ MMP-3 in torn tendon
Rotator cuff tear	Longo et al (46)	Total cholesterol (TC), Triglycerides in serum. TC: NS in cases vs. controls, P=0.1. Triglycerides: NS in cases vs. controls, P=0.06.	No
Rotator cuff tear	Papalia et al (52)	Nonprotein nitrogen plasma (NPN), creatinine in plasma. NPN: cases > controls, P=0.03. Creatinine: NS in cases vs. controls, P=0.66.	Yes ↑ NPN in plasma of cases
Rotator cuff tear	Xu et al (54)	GAP43, PGP9.5 in biopsies of torn supraspinatus vs matched moderate tendinopathic subscapularis tendons from same patient, vs intact subscapularis tendon from controls. GAP43: moderate tendinopathic subscapularis > torn supraspinatus and control, P<0.0001 & P<0.05, respectively; control > torn supraspinatus, P<0.05. PGP9.5: moderate tendinopathic subscapularis > torn supraspinatus and control, P<0.0002 & P<0.05, respectively; control > torn supraspinatus, NS.	Yes ↑ GAP43 in moderate tendinopathic > control > torn tendon ↑ PGP9.5 in moderate tendinopathic > control & > torn tendon
Rotator cuff tendinitis	Rechardt et al (34)	CRP in serum. CRP (dichotomized at ≥ 3 mg/L): high levels associated with rotator cuff tendinitis in males, OR = 0.8 (95% CI: 0.4-1.6), and females, OR = 1.3 (95% CI 0.8-2.2).	No
Frozen shoulder (Adhesive capsulitis)	Bunker et al (35)	Total cholesterol (TC), triglyceride in serum. TC: cases > controls, P<0.01. Triglycerides: cases > controls, P<0.02.	Yes ↑ TC and triglycerides in serum of cases
Frozen shoulder (adhesive capsulitis)	Gumina et al (33)	CRP, glucose, HDL, LDL, rheumatoid factor, total cholesterol in serum. See table 2.	No
Frozen shoulder (adhesive capsulitis)	Lubis et al (56)	MMP-1,2, TIMP-1,2, MMP/TIMP ratio, TGF- β 1 in serum. MMP-1, 2: cases < controls, P<0.001. TIMP-1, 2: cases > controls, P<0.001. MMP/TIMP ratio: cases < controls, P<0.001. TGF- β 1: cases > controls, P<0.001.	Yes ↓ MMP-1,2 and MMP2,3/TIMP ratio in serum of cases ↑ TIMP-1,2, TGF- β 1 in serum of cases
Frozen shoulder (adhesive capsulitis)	Xu et al (55)	CD34, GAP43, PGP9.5, P75 in tendons biopsies of adhesive capsulitis versus rotator cuff tears. GAP43: adhesive capsulitis > rotator cuff tear, P<0.05. PGP9.5: adhesive capsulitis > rotator cuff tear, P<0.01.	Yes ↑ PGP9.5 and GAP43 in tendons with adhesive capsulitis vs rotator cuff tears
Shoulder pain (unilateral & bilateral)	Rechardt et al (34)	CRP in serum. CRP (dichotomized at ≥ 3 mg/l): high levels associated with bilateral shoulder pain in females, OR = 1.4 (1.0-2.1:95% CI); no association in males, OR = 1.1 (95% CI 0.7-1.8).	Yes ↑ CRP in serum of females with shoulder pain

Continued

Table 1. Continued

MSD classification and diagnosis	Author	Major results (case-control comparison)	Conclusion
Neck/shoulder disorders & symptoms			
Trapezius myalgia	Flodgren et al (61)	Glutamate, lactate, PGE2, pyruvate in muscle dialysate (MD); lactate in serum. Repetitive work: MD: glutamate, lactate and pyruvate increased vs baseline; all $P < 0.005$. PGE2 remained unchanged. Blood: lactate: decreased from baseline to after work, $P < 0.005$. Recovery: MD: PGE2 and glutamate decreased, lactate remained increased, pyruvate increased progressively compared to work; all $P < 0.005$. Blood: lactate returned to baseline values by end of recovery, $P < 0.005$.	Perhaps No control subjects were included in this study. Authors conclude normal response in these cases when comparing them to a previous similar study with normal subjects (see Flodgren (2005)).
Trapezius myalgia	Gerdle et al (74)	Bradykinin, kallidin in muscle dialysate (MD). Whole experiment (linear mixed model): Bradykinin, NS group effect; Kallidin: cases > controls, $P = 0.02$. NS interaction (group x time) for bradykinin and kallidin. Repetitive work (ANOVA: work minus baseline): Bradykinin: cases > controls, $P < 0.05$; Kallidin: NS in cases versus controls.	Yes ↑ MD kallidin during whole experiment in cases ↑ MD bradykinin during work in cases
Trapezius myalgia	Ghafouri et al (64)	Serotonin (5-HT) in muscle dialysate (MD). Rest (baseline): cases > controls, $P = 0.04$. Repetitive work: cases > controls, $P = 0.002$. Mental stress (TSST): cases > controls, $P = 0.003$. Recovery: cases > controls, $P < 0.001$.	Yes ↑ MD 5-HT at baseline, during repetitive work, during mental stress, and during recovery.
Trapezius myalgia	Hadrevi et al (62)	Proteins (proteomics study) in muscle biopsies. Partial least square discriminate analysis (PLS-DA) identified (SwissProt ID) 28 unique proteins separating case and control muscles (variable of importance >1). Glycogenolysis (P11217; P36871), cases > controls. Glycolysis (P04075; P04406; P13929; P14618) cases < controls. Tricarboxylic acid cycle (Q99798) cases < controls, (Q02218) cases > controls. Acute response (P08107; P01009) cases > control; (P07451) cases < controls Contractile (P60709; P13533; Q9UKX2; P123883) cases < controls, (P62736; Q00872; P12883; Q96A32) cases > control. Structural and other (P04264; O75112; P07437; P02768; P82650; P13645; P25705) cases < controls, (P40123; Q8TD99) cases > controls.	Yes ↑ Inflammatory mechanisms Altered anti-oxidant protection in cases ↑ Glycogen breakdown in cases ↓ Glycolysis in cases Possible alteration of contractile apparatus, TCA and muscle structure in cases
Trapezius myalgia	Larsson et al (71)	Lactate, pyruvate, glutamate, serotonin (5-HT), potassium, bradykinin, GM-CSF, IL-18, IL-6, TNF- α , IFN- γ , IL-2, IL-4, IL-5, IL-10, in muscle dialysate (MD); & Urine-cortisol. 120 min after catheter insertion: MD 5-HT (serotonin): cases > controls, $P = 0.02$. Work day (8 h): MD Glutamate and pyruvate: cases > controls, $P = 0.005$ & $P = 0.03$, respectively; lactate, potassium, bradykinin NS in cases vs. controls. Mixed Models Analyses (group, time, interaction): group effects during work: glutamate, $P = 0.006$; pyruvate, $P = 0.02$. NS group effect for: bradykinin, IL-18, IL-2, IL-4, IL-5, IL-6, IL-8, IL-10, IFN- γ , GM-CSF, TNF- α , cortisol. NS interaction effect (group x time) for: all biomarkers. Recovery (1 h): MD glutamate, lactate, pyruvate, potassium, bradykinin: NS in cases vs. controls. Immediately after recovery: MD 5-HT: cases > controls, $P = 0.001$. Relative recovery: MD lactate NS in cases vs. controls, $P = 0.11$.	Yes ↑ MD glutamate & pyruvate in cases during work; ↑ MD 5-HT, both before and after work in cases
Trapezius myalgia	Rosendal et al (75)	Glutamate, lactate, pyruvate, serotonin (5-HT) in muscle dialysate (MD); plasma creatine kinase and LDH. Rest: MD serotonin (5-HT): cases > controls, $P < 0.01$. Group effect (all time points): MD lactate, glutamate, pyruvate: cases > controls, $P < 0.001$, $P < 0.05$, and $P < 0.001$, respectively. With exercise: MD lactate increased in cases and returned to baseline during recovery 1, $P < 0.02$. MD lactate NS in controls. MD pyruvate increased in cases, $P < 0.005$; tended to be higher at recovery 1, $P = 0.08$; similar to baseline level at recovery 2. MD pyruvate NS in controls. At recovery: MD glutamate lower than baseline in cases; NS in controls. Plasma: creatine kinase and LDH, NS in cases vs controls.	Yes ↑ MD 5-HT at baseline in cases ↑ MD lactate, glutamate, pyruvate in cases, all time points
Trapezius myalgia	Rosendal et al (72)	IL-6, K $^{+}$, LDH, PINP/ICTP ratio in muscle dialysate (MD). Rest: PINP/ICTP ratio: Controls had 2-fold+ higher anabolic ratio than cases, NS, $P = 0.07$. Whole experiment - variance components model (group, time, and interaction): K $^{+}$: group effect, cases > controls, $P < 0.0001$. IL-6, LDH: NS, group and interaction effects.	Yes ↑ MD K $^{+}$ of cases

Continued

Table 1. Continued

MSD classification and diagnosis	Author	Major results (case-control comparison)	Conclusion
Trapezius myalgia	Sjogaard et al (31)	Lactate, pyruvate, glucose, potassium (K+) in muscle dialysate (MD) Rest: Lactate and pyruvate: cases > controls. Glucose, potassium: NS in cases vs. controls. Repetitive work: all biomarkers: NS in cases vs. controls. Mental stress: all biomarkers: NS in cases vs. controls.	Yes ↑ MD lactate and pyruvate at base-line in cases
Trapezius myalgia	Sjörs et al (65)	Cortisol in saliva. Circadian: NS group or interaction effect. Stress test (TSST): NS group or interaction effect. Peak cortisol after TSST and mean peak time point in response to TSST similar in both groups. Re-running analyses including only responders*: mean peak time point (after TSST): cases > controls, $P=0.03$. (*cortisol response > 2.5 nmol/l above pre-TSST level, an elevation thought to reflect a cortisol secretory episode.)	Yes, ↑ cortisol in saliva of responders
Myofascial pain syndrome (especially in periscapular region)	Ozgocmen et al (77)	HDL, LDL, triglyceride, total cholesterol, VLDL in blood. LDL, total cholesterol, triglyceride, VLDL: cases > controls, $P=0.02$, $P=0.003$, $P=0.01$, and $P=0.008$, respectively. HDL: NS in cases vs. controls.	Yes ↑ LDL, total cholesterol, triglyceride, VLDL in serum of cases
Chronic neck/shoulder pain	Flodgren et al (67)	Glutamate, PGE2 in muscle dialysate (MD). Glutamate and PGE2: NS, cases vs. controls in time-point of equilibrium.	No
Chronic neck/shoulder pain	Ghafouri et al (76)	PEA, SEA in muscle dialysate (MD). PEA: pre-exercise: cases > controls, $P<0.05$; post-exercise: cases > controls, $P<0.001$. SEA: pre-exercise: cases > controls, $P<0.001$; post-exercise: cases > controls, $P<0.01$.	Yes ↑ MD PEA and SEA of cases, both pre- and post-exercise
Shoulder & neck pain	Nilsen et al (68)	Noradrenaline PPP, noradrenaline PRP, adrenaline PPP, cortisol in plasma. Noradrenaline PPP, noradrenaline PRP, adrenaline PPP, cortisol: NS in cases vs. controls.	No
Shoulder & neck pain (SNP)	Riva et al (57)	Cortisol in saliva. Awakening, and 30 and 60 min later - significant group effect. Post-hoc tests: shoulder/neck pain (SNP) vs. controls, overall NS effect; awakening NS effect; 30 min: SNP > controls (Bonferroni: $P=0.077$, Dunnett's: $P=0.046$); 60 min: NS effect.	Perhaps Women with SNP had a tendency for higher saliva cortisol
Neck & shoulder disorders (see Kaergaard (2000) for disorder breakdown)	Kaergaard et al (37)	Testosterone in plasma. Baseline: neck & shoulder disorder associated with reduced testosterone after adjusting for age, BMI, workgroup, oral contraceptives, estrogen, smoking in multiple linear regression, $P=0.005$. 1-year follow-up: NS association between neck & shoulder disorders and testosterone when cross-sectional multiple linear regression was performed at follow up.	Perhaps Indication of ↓ testosterone in cases. But study failed to clarify relationship.
Hand/wrist disorders & symptoms			
Carpal tunnel syndrome (CTS)	Freeland et al (70)	IL-1, IL-6, MDA, PGE2 in serum. MDA: cases > controls, $P<0.001$; PGE2, IL-1, and IL-6: NS in cases vs. controls.	Yes ↑ MDA in serum of cases
Carpal tunnel syndrome (CTS)	Tsuji et al (58)	Tenascin-C in tenosynovium See table 2.	No
Hand-arm vibration syndrome (HAVS)	Bovenzi et al (39)	Endothelin (ET1-21) in saliva. Before cold challenge: ET1-21 forestry workers with HAVS > controls, $P=0.04$; forestry workers with HAVS > forestry workers without HAVS, $P=0.09$, NS. After cold challenge: ET1-21 forestry workers with HAVS > controls, $P=0.04$; forestry workers with HAVS vs forestry workers without HAVS, $P=0.19$, NS.	Yes ↑ ET ₁₋₂₁ in saliva of forestry workers with HAVS
HAVS	Kennedy et al (59)	sICAM-1, IL-8 in blood. sICAM-1: cases > controls, $P=0.02$; IL-8: cases < controls, $P<0.01$.	Yes ↑ sICAM-1 and ↓ IL-8 in blood of cases
HAVS	Lau et al (63)	MDA, PMN in blood. MDA: cases > controls, $P<0.0001$. PMN: increased PMN activity in cases > controls, $P=0.0004$.	Yes ↑ MDA and PMN activity in blood of cases
HAVS	Nakamura et al (60)	Endothelin-1 in plasma Endothelin-1: effects of cold water-immersion, $P<0.05$, of group, $P<0.01$, and interaction, $P<0.05$. Post-hoc Dunnett's test during immersion: cases < controls, $P<0.01$. NS group difference in endothelin-1 concentrations before or after immersion.	Yes ↓ plasma endothelin-1 in cases during cold water-immersion
Dupuytren's contracture	Augoff et al (73)	MMP-2 activation ratio (active MMP-2 / latent MMP-2) in palmar fascia MMP2-activation ratio: cases > controls, $P<0.001$. Each group (I-IV) had higher MMP2-activation ratio than control group (all, $P<0.001$).	Yes ↑ active MMP2 in fascia of cases

Continued

Table 1. Continued

MSD classification and diagnosis	Author	Major results (case-control comparison)	Conclusion
Dupuytren's contracture	Iqbal et al (47)	Mesenchymal stem cells (CD73, CD90, CD105) and hematopoietic (CD34) markers in tissues. Cells from cord, nodule, PNF, and SON of cases more positive for MSC markers CD73, CD90, and CD105, than controls, all $p < 0.05$, except for CD105 in PNF vs. control fat, NS $P = 0.09$. Cells from cord, nodule, and SON more positive for hematopoietic marker CD34 (cord, $P = 0.01$; nodule, $P = 0.02$; and SON, $P = 0.0009$); DPF vs. control fat, NS. Cells from cords, SON, PNF, and DPF had more CD34-73+ 90 +105+ cells than matching control tissue, all $P < 0.01$.	Yes ↑ mesenchymal stem cells in tissues from cases
Dupuytren's contracture	Hnanicek et al (48)	Ferritin, iron, total iron binding capacity, total cholesterol, transferrin saturation in serum. Each: NS, cases vs. controls	No
Dupuytren's contracture	Ulrich et al (49)	MMP-1, 2, 9; TIMP-1,2; MMP/TIMP ratio in serum. TIMP-1: cases: > controls, $P < 0.05$ TIMP-2: NS, cases vs. controls. MMP-1, 2, 9: NS, cases vs. controls. MMP/TIMP ratio (consists of all measured markers): cases < controls, $P < 0.05$.	Yes ↑ TIMP-1 in serum of cases ↓ MMP/TIMP ratio in serum of cases
Dupuytren's contracture	Ulrich et al (50)	MMP-2, 9; TIMP-1, 2 in palmar fascia. x-fold expression relative to normal palmar fascia MMP-2: case nodules: 4x, $P < 0.05$; case cords: 2.5x, $P < 0.05$. MMP-9: NS nodules and cords. TIMP-1: case nodules: 7.2x, $P < 0.05$; case cords: 2.5x, $P < 0.05$ TIMP-2: case nodules: 3.2x, $P < 0.05$; case cords: 2.1x, $P < 0.05$	Yes ↑ MMP-2, TIMP-1, TIMP-2 in tissues from cases
Non-specific pain in finger extensor musculature	Moreno-Torres et al (66)	Pi/PCr (phosphorous magnetic resonance spectroscopy) in hand extrinsic extensor muscles. P-MRS pattern B: appearance of a second acidic (low pH) inorganic phosphate (Pi) peak in cases > controls, $P = 0.0001$.	Yes ↑ acidic Pi peak using MRS in cases
Upper-extremity disorders and symptoms without specified location			
Musculoskeletal symptoms	Akesson et al (38)	Selenium (Se) in plasma; mercury (Hg) in whole blood, plasma and urine Se: NS association between Se concentration (in those not taking selenium tablets, $N = 315$) and musculoskeletal symptoms in past 7 days, according to particular region of pain. NS association between taking Se tablets and musculoskeletal symptoms. Hg: results in table 2.	No
Musculoskeletal disorders (MSDs) due to overuse (see Carp (2007) for disorder breakdown)	Carp et al (40)	IL-1 β , TNF- α , IL-6, CRP in serum. See table 2.	Yes
Specific & non-specific MSDs, predominantly lateral epicondylitis & rotator cuff tendinitis (see Rechartd (2011) for disorder breakdown)	Rechartd et al (42)	CRP, IL-1 α , IL-1 β , IL-1 RA, IL-6, IL-18, IL-33, sST2, sIL-1RII, TNF- α in serum Unconditional logistic regression models adjusting for age, gender, smoking, BMI, physical activity alcohol consumption and sleep activity. sIL-1RII: cases > controls, $P < 0.001$. sST2: cases > controls, $P = 0.014$; $P = 0.052$ after Bonferroni correction. IL-18: cases < controls, $P = 0.005$. IL-1Ra, IL-8, IL-33, CRP: NS in cases vs. controls. IL-1 α , IL-1 α , IL-6, TNF- α : below level of detection.	Yes ↑ sIL-1RII and ↓ IL-18 in serum of cases
MSDs (see Rechartd (2013) for disorder breakdown)	Rechartd et al (41)	Andiponectin, CRP, glucose, HDL, leptin, LDL, resistin, total cholesterol, triglyceride, visfatin in serum. See table 2.	Yes

biopsies from subjects with Dupuytren's contracture (44, 50, 73). A study of adhesive capsulitis found reduced MMP in patients (56). Two studies saw no association with this category of potential biomarkers (49, 51).

Three studies found increased TIMP (most consistently, TIMP-1) in serum from subjects with adhesive capsulitis, and in serum or palmar fascia from subjects with Dupuytren's contracture (49, 50, 56). Two of these showed a decreased MMP/TIMP ratio in serum of subjects with fibroproliferative MSD (49, 56). A study where the collagen synthesis-to-degradation ratio (PINP/ICTP) was examined in muscle interstitial dialysate found no association in subjects with and without trapezius myalgia (72). The one study that did not find a significant association with collagen-repair markers used adjacent tendon regions from subjects with torn supraspinatus tendons as internal control tissues, and found increased TIMP and MMP in both the torn and seemingly intact adjacent tendon regions (51). These authors postulated that tendon areas near injured tendons can present with altered cellular metabolism.

In summary, the most consistent result was increased TIMP-1 (a collagen-repair marker) in serum or tissue from subjects with Dupuytren's contracture and adhesive capsulitis. Additionally, collagen turnover (MMP/TIMP level) was decreased in these subjects.

Muscle metabolites/neurotransmitters. Eleven studies examined muscle metabolites/neurotransmitters using local sampling methods (31, 61, 62, 64, 66, 67, 71, 72, 74–76). Eight of these found at least one significant association between a biomarker and the MSD (31, 62, 66, 71, 72, 74–76), while three found no association (61, 67, 76). Except in one study (62), the concentration of the biomarker was greater among cases than controls. In one study (62), an increase was noted for 11 possible biomarkers and a decrease for 16. In all but one study (66), biomarkers were measured in the trapezius muscle interstitium or via muscle biopsy. In most of the muscle microdialysate experiments, a rest (baseline) period was followed by a repetitive physical or mental work challenge and then a subsequent recovery period.

Several studies reported higher levels among cases than controls of muscle metabolites that play a role in regulating pain. This was observed with lactate and pyruvate at baseline (31) and with glutamate (75), kallidin (74), lactate (75), and potassium (K⁺) (75) across an experiment. Three studies reported elevated bradykinin (74), glutamate (71), and pyruvate (71) among cases versus controls during repetitive work. Another study (61) also observed increased glutamate, lactate, and pyruvate in cases during work versus baseline levels. In contrast, Sjogaard et al (31) observed no significant changes in these muscle metabolites with work or mental stress, despite increased lactate and pyruvate among cases at

baseline compared to controls (31). With regard to the recovery period, two studies reported a rapid return of these metabolites to baseline levels (71, 75), while one (61) observed maintenance of work-induced increases in lactate and progressively increasing pyruvate levels during the recovery period. In muscle biopsies from subjects with trapezius myalgia, one study showed increased enzymes in glycogenolysis and the initial steps of glycolysis but a decrease of enzymes in the final steps of glycolysis (62).

In the four studies that examined glutamate, one observed an elevated level in cases during the entire experiment (75), while another found an elevation among cases versus controls during work but not during recovery (71). These results differ considerably from those of Flodgren et al (67) who observed no differences in biomarker concentration between cases and controls at rest. A repeat of this study (61) showed an elevation in cases of glutamate during repetitive work versus baseline (there was no control group), although they concluded that this was a normal increase due to muscle metabolism.

Of the four studies investigating lactate, one observed an elevation in cases versus controls during the entire experiment (75) while another (71) found no overall difference between cases and controls. Sjogaard et al (31) observed a comparative elevation in lactate during baseline, but not in response to repetitive physical or mental work, or during recovery. In contrast, Flodgren and colleagues (61) found increased lactate during repetitive work.

Of the four studies that examined pyruvate, one observed an elevation during the entire experiment (75), one observed an elevation at baseline only, but not in response to work or during recovery (31), and one observed an increase in response to work, but not during recovery (71). One study (61) found increased pyruvate during repetitive work.

Three studies examined 5-HT in muscle dialysates, a monoamine neurotransmitter known to sensitize nociceptors, and found increases in cases with trapezius myalgia and chronic neck/shoulder pain at baseline versus controls (64, 71, 75). One study (64) found increased 5-HT levels in cases during both physical and mental work, while two found increased 5-HT in cases during recovery (64, 71). Another study examining two other muscle metabolites that regulate pain responses, N-palmitoylethanolamine (PEA) and stearoylethanolamide (SEA), found a significant association between levels of these analytes in muscle dialysate in subjects with chronic neck pain, compared to controls (76).

Lastly, one study examined local muscle changes in the hand using phosphorous magnetic resonance spectroscopy. Moreno-Torres et al (66) observed an increased peak of acidic inorganic phosphate (Pi) in hand muscles in subjects with non-specific pain in finger musculature.

This factor is thought to induce fatigue by interfering with biochemical steps in the cross-bridge cycle.

In summary, results with respect to muscle metabolites/neurotransmitters were mixed with the exception of Pi, PEA and SEA, which were examined in only one study each, and 5-HT in which increases were generally observed in trapezius myalgia cases.

Lipids/insulin resistance. Seven studies investigated lipids and cholesterol using systemic sampling methods, ie, blood samples (33, 35, 41, 43, 46, 48, 77). Four observed statistically significant associations with MSD (35, 41, 43, 77); three did not (33, 46, 48). Total cholesterol (TC) was examined in all seven studies. Two studies investigated rotator cuff tears (43, 46), showing increased (43) or no difference in TC (46) among cases versus controls. Two studies investigated frozen shoulder (adhesive capsulitis), also with divergent results. One study found greater levels of TC in cases with frozen shoulder (35), whereas one did not (33). No association was found between TC and Dupuytren's contracture (48). Ozgocmen et al (77) found an association between increased TC in subjects with myofascial pain versus controls. When investigated, no correlation between TC and severity level was found (33, 41).

The association between serum triglycerides and MSD was studied in five papers. Of those studies investigating rotator cuff tears (43, 46), one showed increased triglycerides in cases (43) and one found no difference between cases and controls (46). Increased triglycerides were observed in adhesive capsulitis (35) and myofascial pain (77), and among MSD cases reporting greater pain intensity (41).

Four studies investigated low- and/or high-density lipoprotein (LDL and/or HDL) in cases with MSD. In cases with rotator cuff tears, an increased LDL was shown (43) and, in adhesive capsulitis, both LDL and HDL were increased (33). In a myofascial pain study, LDL was increased but HDL was not (77). A decreased concentration of HDL was found among MSD cases reporting greater pain intensity (41).

In summary, results with respect to lipids including cholesterol were mixed, with the exception of triglycerides where increased levels were generally observed among cases.

Are there biochemical biomarkers associated with the severity of upper-extremity MSD?

As shown in table 2, five studies demonstrated a relationship between MSD severity and biomarker concentration (40, 41, 49, 50, 76). Six studies investigated biomarker concentration in relation to a severity score or disease stage determined during a physical examination (33, 40, 45, 49, 50, 73). Increased collagen-repair

marker, TIMP-1 (49, 50), and a decreased MMP/TIMP ratio (49) were observed at an earlier (more active) Dupuytren's contracture stage. Increased MMP-2, a collagen degradation marker, was found in an earlier stage of Dupuytren's contracture in one study (50), although not in another (73). Carp et al (40) found positive correlations between inflammatory markers IL-1 β , IL-6, and TNF- α and MSD severity in a group of early stage (mean 1.7 months) patients.

Six studies solicited pain ratings from participants (37, 41, 61, 72, 74, 76). Ghafouri et al (76) found increased concentrations of PEA and SEA when patients reported smaller differences in pain between exercise and baseline, or between exercise and recovery periods. In Rechart et al's study of subjects with MSD of >1 month (41), triglyceride and visfatin were increased and HDL decreased among those reporting greater pain.

In summary, only a few studies found associations between biomarkers and severity of upper-extremity MSD. Results with respect to inflammatory markers were mixed. While Carp et al (40) observed increased inflammatory marker concentrations with increased MSD severity in a mix of MSD, Ghafouri et al (76) found increased inflammatory marker concentrations in chronic neck/shoulder pain patients reporting a lower increase in pain in response to repetitive low force work. Lastly, increased collagen-repair markers were found in earlier stages of Dupuytren's contracture (50).

Discussion

We conducted a comprehensive systematic review to summarize the current state of biochemical marker research in neck and upper-extremity MSD. We used a critical approach to synthesize results for the research questions: (i) are there biochemical markers associated with neck and upper-extremity MSD? and (ii) are there biochemical markers associated with the severity of neck and upper-extremity MSD? We found associations between biochemical biomarkers and neck and upper-extremity MSD in the sufficient-quality studies. A wide range of biomarkers were examined. Nonetheless, with several exceptions listed below, there was little consistency of results. However, we were able to identify a few commonalities.

An increased collagen-repair marker, TIMP-1, and decreased collagen turnover (MMP/TIMP ratio) was observed in serum among cases with Dupuytren's contracture (49) and early stage frozen shoulder (56). TIMP-1 was also increased in tissue from Dupuytren's contracture cases relative to normal palmar fascia (50). Elevated serotonin (5-HT) was found among muscle interstitium in trapezius myalgia cases (64, 71, 75, 76). Increased triglycerides were seen in serum of subjects

Table 2. Are there biochemical biomarkers associated with the severity of neck and upper-extremity musculoskeletal disorders (MSD)?

MSD classification or diagnosis	Author	Biomarker	Severity measure (mean & range if known)	Symptom duration (mean & SD if known)	Major results (association between biomarker & disease)	Conclusion
Neck disorders and symptoms						
Rotator cuff tear	Lakemeier et al (45)	VEGF	Bateman classification, 1963		No association between VEGF & Bateman's classification.	No
Rotator cuff tear	Longo et al (46)	Total cholesterol, triglyceride	Size of rotator cuff tear		No association between biomarkers & size of tear	No
Rotator cuff tear	Papalia et al (52)	NPN, creatinine	Size of rotator cuff tear	49 months (range: 28-60) of pre-operative follow-up	No association between biomarkers & size of tear: NPN: P=0.76. Creatinine: P=0.09.	No
Frozen shoulder (adhesive capsulitis)	Gumina et al (33)	CRP, glucose, HDL, LDL, rheumatoid factor, total cholesterol	Constant Score (shoulder function) (Constant & Murley 1987) (mean 39.0 (SD 6.7) range: 18-49)	≤6 weeks from initiation symptoms ^a	No correlation between any biomarkers & Constant Score.	No
Neck/shoulder disorders & symptoms						
Trapezius myalgia	Flodgren et al (61)	Lactate, pyruvate, glutamate, PGE2	VAS pain in right shoulder median (range) [47 (12-79) for the last 12 months, 34 (14-63) for the last 7 days, & 27 (4-78) when arriving at the laboratory]	Mean 70.5 (SD 74.7) months	Pain intensity reported during the experiment not correlated with MD [glutamate] (P=0.276) or [PGE2] (P=0.492).	No
Trapezius myalgia	Gerdle et al (74)	Bradykinin, kallidin	VAS pain during experiment		No multivariable correlation between the increase in pain intensity during exercise (reported through VAS) & bradykinin or kallidin.	No
Trapezius myalgia	Rosendal et al (72)	IL-6, LDH, K+, collagen turnover (PINP/ICTP ratio)	VAS pain during experiment Baseline VAS cases mean 53.3 (SE 8.9) * controls: 0 (0) * estimated from figure	Unknown average 55 months (range 7-216) on disability	All subjects: mean K+ correlated w. VAS (r=0.74, P<0.001); stratify on group, only sig. in controls (r=0.52, P=0.03).	No
Chronic neck/shoulder pain	Ghafari et al (76)	PEA, SEA	Pain intensity ratings during experiment (0-10 scale)	>6 months ^a	Reported pain intensity in cases, partial least squares regression modeling During exercise minus baseline: PEA at 140 min: VIP = -1.51 SEA at 180 min: VIP = -1.25 PEA at 180 min: VIP = -0.93 Post-exercise minus during exercise: PEA at 140 min: VIP = -1.91 PEA difference 180 min minus 140 min: VIP = 1.36 * VIP - variable influence on projection, VIP ≥ 1 regarded as significant, 0.8 < VIP < 1 regarded as baseline significant.	Yes
Neck & shoulder disorders (see paper for complete list)	Kaergaard et al (37)	testosterone	Modified pain severity grading method (Von Korff, et al 1992) (scale 0-108) [baseline mean: 20.2 (SD 21.2) 1-year follow-up: mean 22.0 (SD 20.5)]		No correlation between change in testosterone level & change in pain level borderline to follow-up.	No
Hand/wrist disorders and symptoms						
Carpal tunnel syndrome	Tsuji et al (58)	Tenascin-C	Modified pain severity grading method (Von Korff, et al 1992) (scale 0-108) [baseline mean: 20.2 (SD 21.2) 1-year follow-up: mean 22.0 (SD 20.5)]		No difference in tenascin-C concentration between duration groups.	No
Hand-arm vibration syndrome (HAVS)	Lau et al (63)	PMN, MDA	Number & duration of Raynaud's attacks during a 2-week period	Median 7 (range 1-15) years	No correlation between PMN or MDA & number or duration of Raynaud's attacks.	No
Dupuytren's contracture	Augoff et al (73)	MMP-2	Iselin's classification of disease stage (Eckes, et al 2000)		No difference between MMP-2 activation ratios by clinical stage.	No

Continued

Table 2. Continued

MSD Classification or diagnosis	Author	Biomarker	Severity measure (mean & range if known)	Symptom duration (mean & SD if known)	Major results (association between biomarker & disease)	Conclusion
Dupuytren's contracture	Ulrich et al (49)	MMP-1, 2, 9, TIMP-1,2, MMP-to-TIMP ratio	Stage of Dupuytren's (Luck 1959)	Range: 7 months - 30 years	TIMP-1: early stage > late stage, $P < 0.05$. TIMP-2, MMP-1, MMP-9: NS, early vs. late stage. MMP-to-TIMP ratio: early stage < late stage, $P < 0.05$.	Yes
Dupuytren's contracture	Ulrich et al (50)	MMP-2, 9, TIMP-1, 2	Cords present in late stage and nodules present in active phase (earlier than cords)	Range: 9 months–24 years	MMP-2: nodules > cords, $P < 0.05$. TIMP-1: nodules > cords, $P < 0.05$. MMP-9, TIMP-2: NS, nodules vs. cords, $P > 0.05$.	Yes
Upper extremity disorders & symptoms without specified location Musculoskeletal symptoms	Akesson et al (38)	Selenium, mercury	Number of areas with symptoms during the past 12 months by occupational group Median (4 - dental hygienists 3 - dentists 2 - female referents 1 - medical nurses 0.5 - male referents)		No association of biomarkers with number of symptomatic anatomical regions	No
Musculoskeletal disorders (MSDs) due to overuse (see paper for complete list)	Carp et al (40)	IL-1 β , TNF- α , IL-6, CRP	UBMA (upper-body musculoskeletal assessment) (Kramer et al 2000) UBMA score [mean 81.6 (SD 20.5) symptomatic, mean 39.0 (SD 5.0) asymptomatic]	Mean 51.7 (SD 18.9) days range 17–76 days	Spearman's rank correlation CRP: $r = 0.81$, $P < 0.0001$. IL-1 β : $r = 0.70$, $P < 0.0001$. TNF- α : $r = 0.66$, $P < 0.0001$. IL-6: $r = 0.52$, $P = 0.003$. Ordinal logistic regression analysis using the cutoff points for UBMA score indicated (0-49,50-74, 75-99,100-150), CRP & TNF- α in same model. CRP: OR >999.9 (95% CI: 111.8–>999.9); $P = 0.0003$ TNF- α : OR = 2.2 (95% CI: 1.4–3.4); $P = 0.0007$	Yes
MSDs (see paper for complete list)	Rechardt et al (41)	cholesterol, triglyceride, HDL, LDL, glucose, CRP, leptin, adiponectin, resistin, visfatin	VAS (0-100) mean 48 (SD 22) pain intensity in the last week	< 1 month ^a	Triglyceride: increases with pain intensity, tertiles (mmol/L) <0.72, OR=1 (ref) (95% CI 0.72-1.08), OR=1.4 (95% CI 0.5-3.8) >1.08, OR=2.6 (95% CI 1.0-6.8) HDL: decreases with pain intensity, tertiles, (mmol/L) >1.83, OR=1 (ref) (95% CI 1.48-1.83), OR=0.7 (95% CI 0.2-2.0) <1.48, OR=3.9 (95% CI 1.4-10.1) Visfatin: increases with pain intensity (continuous, increment of 1 SD) OR=1.4 (95% CI 1.0-2.1)	Yes

^a Inclusion criteria.

with rotator cuff tears (43), frozen shoulder (35), myofascial pain (77), and subjects reporting greater pain intensity with mostly tendon-related MSD (41). In contrast to the first research question, associations between biomarkers and severity of neck and upper-extremity MSD were observed in only a few studies. Greater collagen degradation and repair is likely occurring in earlier stages of Dupuytren's contracture (49, 50).

Both Dupuytren's contracture and frozen shoulder are fibroproliferative disorders (78). Biomarker papers

examined in this review suggest that an imbalance in extracellular matrix remodeling may be occurring, similar to other fibroproliferative conditions (79). MMP, which degrade collagen, are active in wound repair, whereas TIMP are inhibitors of MMP. An increased concentration of TIMP may result in fibrosis. Two studies demonstrated increased TIMP-1 in serum of subjects with fibroproliferative disorders and an inverse association of the MMP/TIMP ratio in MSD cases (49, 56), consistent with the above interpretation.

In a previous review of microdialysis studies in human myalgia and tendon-related conditions (20) and in a review of microdialysis studies specifically focused on chronic muscle pain conditions (22), the authors concluded that 5-HT may potentially be a biomarker of chronic myalgia; we concur with their conclusions. They note that 5-HT can activate nociceptors which transmit pain signals to the central nervous system, and that it may have an analgesic or hyperalgesia effect depending on the particular cell and receptor targeted (20). Potential involvement of this mechanism in trapezius myalgia should be further examined in longitudinal studies.

A number of studies found associations between triglyceride concentration and MSD. However, there is insufficient evidence to suggest that triglycerides are a necessary condition for MSD occurrence. Increased serum triglycerides are often found in obesity, metabolic syndrome, and diabetes. Thus, it is likely that the observed increases in triglycerides with MSD are due to comorbidity of these conditions.

Nonetheless, several mechanisms have been hypothesized by which hyperlipidemia may increase tendinopathy risk. First, lipids (xanthoma) may be deposited on tendons which may alter their biomechanical properties (43). Xanthoma may also initiate a low-grade inflammation in tendons, initiating chronic tendon degeneration (80, 81). Lastly, lipids may alter the extracellular matrix and reduce circulation, both impairing tissue healing (43). These possible lipid effects on tendons combined with metabolic and biomechanical alterations due to tendon repetitive loading (82, 83) may increase MSD risk.

Methodological limitations of the articles in the review

Selection bias. It was only possible to ascertain the response rate to an invitation to participate in 9/44 sufficient-quality studies. In many articles, it was stated that patients were enrolled consecutively, but the control group response rate was not stated. Without response rates, it is not possible to adequately assess selection bias, ie, whether cases and controls fairly represent the underlying population, or to what extent cases and controls may be comparable. Future studies should delineate the response rates for both cases and controls.

In addition, there is potential for bias due to selection of referents. For our first research question, we evaluated studies that compared subjects with the MSD or symptom(s) of interest to one or more referent groups. The referent group sometimes consisted of asymptomatic participants. However, several studies utilized convenient patient populations as controls (supplementary table S5, www.sjweh.fi/data_repository.php). These may not necessarily be appropriate controls for the biomarkers under investigation. Some studies of shoulder disorders used referents with shoulder disorders

presumably affecting different anatomical systems than the condition under study. For instance, in Lakemeier et al's studies of rotator cuff tears (44, 45), patients with traumatic humeral head fractures were utilized as controls. However, the biomarkers studied could be present in both cases and controls. Vascular endothelial growth factor (VEGF) promotes both angiogenesis and bone formation (84), and genes encoding MMP were found to be up-regulated in fracture repair (85). Concentrations of the biomarkers examined were greater among cases than controls in Lakemeier et al's studies (44, 45). Nonetheless, when individual studies each examine a particular biomarker for a specified disease, yet select unhealthy controls with differing disorders, mixed results may occur. Selection of controls from the same hospital as cases can be advantageous (in that confounding factors are more apt to have similar distributions within groups); nevertheless, disease processes among unhealthy controls could potentially affect the presence or absence of the biomarkers being studied. Likewise, collecting tissues during surgery from nearby seemingly "intact" tissues to use as internal negative controls may lead to confusing results as both may have altered cellular metabolism, as was found in the Castagna et al study in which MMP and TIMP were elevated in both torn and adjacent "intact" tendon regions (51).

Spectrum effect (variation of biomarkers across subgroups). The term spectrum effect has been used to describe heterogeneous diagnostic test results in different patient subgroups (86). We apply it here to refer to the potentially heterogeneous biomarker associations in populations according to symptom severity or duration, age, gender, and the like. Many of the sufficient-quality studies in this review have been conducted among older patients at the time of surgery (44, 45, 47, 48, 50, 54, 58). Comparatively fewer authors have researched MSD biomarkers in younger subjects with incipient disorders (40, 41). Animal models would suggest that different biomarkers are present at different MSD stages and may differ by age (87–89). Various theories of MSD pathophysiology also suggest that differing biochemical biomarkers would occur at different MSD stages (21, 82, 90). The literature reviewed in the present paper does not allow for any empirical evaluation of the above assertion because there were an insufficient number of studies that detailed symptom duration. We recommend that the duration and/or symptom range of cases in any given biochemical biomarker study be reported. We further suggest that the duration and/or symptom range be restricted or results be analyzed according to duration and/or symptom indicator. Certainly studies enrolling patients with less severe disease than surgical cases, and studies with MSD cases <50 years of age are warranted for future research.

Confounding. Approximately half of the sufficient-quality studies reviewed controlled for potential confounders, eg, gender or age, either through restriction of study subjects or through adjustment in the analysis. It is possible that there was a randomization of potential confounders among the analysis groups in the studies where confounding was not controlled for, which would not invalidate the research. However, potential confounding factors in the relationship between biomarkers and MSD should at least be mentioned in future studies. More ideally, information on these factors should be collected from study subjects and controlled for, either through selection or in the analysis.

Comorbid conditions. The majority of biomarkers in this review are certainly not specific to the MSD studied. Thus, careful consideration of subject comorbidities is warranted in order to avoid spurious associations. For instance, one inflammatory marker, IL-1 β , may be up-regulated in rheumatoid arthritis, gout, osteoarthritis, type 2 diabetes, multiple myeloma, and post-myocardial infarction heart failure (91). In another example, CRP is a sensitive biomarker of inflammation and has been used to identify in individuals with unstable angina pectoris (92), and underlies coronary artery disease and risk of future stroke (93, 94). We recommend careful attention to comorbidities in future research since the specificity of potential biomarkers needs to improve to be useful in clinical practice.

Heterogeneity – other considerations. The collection method (eg, blood, microdialysis of muscle interstitial fluid, tissue biopsy), and detection method used to assay an analyte (eg, single plex versus multiplex ELISA, radioimmunoassay (RIA), immunohistochemistry) may account for some varying results. A recent study compared single-plex versus multiplex ELISA kits for detection of inflammatory cytokines and chemokines in plasma (95). Compared to single-plex ELISA, the multiplex ELISA system proportionally and significantly overestimated IL-1 β and interferon (IFN) gamma, but underestimated IL-6. The authors noted that such differences may distort epidemiologic relationships if data from both methods are merged (95). Tubes selected to collect blood samples can also affect results based on the analyte being assayed or planned detection system (eg, ELISA versus Western Blot analysis) as different anti-coagulants in the tubes can be problematic with some detection systems (96). Other considerations for best sample collection, storage and implementation of internal controls for clinical trial studies are reviewed by de Jager et al (96).

Additional gaps in the literature

The majority of the sufficient-quality studies focused on

neck/shoulder or shoulder disorders, with nine conducting research on trapezius myalgia, and seven researching rotator cuff tears. There is a paucity of biomarker studies of arm and hand MSD. Although Dupuytren's contracture studies were represented here, we found no sufficient-quality studies researching forearm-wrist tendinitis, epicondylitis, or de Quervain's tenosynovitis. Biomarker studies of MSD in the lower extremity could inform upper-extremity MSD research. For instance, there is a rather sizable body of literature on Achilles tendinopathy [eg, see (97–99)] that may be applicable to upper-extremity tendinopathies. Such articles were beyond the scope of the present review.

For most MSD, pathophysiological mechanisms are not well understood. Many researchers chose biomarkers based on theoretical models of MSD pathophysiology. However, some were less grounded in previous work. To some extent it appears plausible that some biochemical biomarkers were selected primarily on the basis of assay with easy availability, and/or low cost. While exploratory studies are necessary, a key concept in improving biomarker discovery would be to choose potential markers with reference to MSD animal models and/or proposed pathomechanisms. An alternative approach to individual testing of a few potential biomarkers would be to use different pattern analysis methods on larger sets of predefined potential markers (selected on either a theoretical or empirical basis). The biomarker could be a specific pattern of individual markers rather than a single substance (100).

Strengths and limitations of the review

This review makes a contribution by systematically synthesizing the literature on biomarkers and MSD. This is the first comprehensive review of biochemical biomarkers in human neck and upper-extremity MSD since 2000 (18). Additionally, it is the only such review that we are aware of to address the question of MSD severity. Since different MSD affect a variety of anatomical systems with differing localization of symptoms, the pathophysiological mechanisms within each anatomical system may differ from each other and thus express separate, relevant biomarkers. However, MSD may also affect multiple anatomical regions (101–103), thus yielding either expression of systemic markers or a pattern of more specific markers related to a specific anatomical segment. We attempted to analyze the results on an MSD basis (separate analysis for each MSD). However, there was little consistency of biomarkers studied within one particular MSD. Hence, we also looked more broadly at classes of biomarkers or at individual biomarkers for multiple upper-extremity MSD.

As with all reviews, publication bias (in which studies with statistically significant results are more apt to be published than studies with null findings) may

certainly be a factor in this review. Although many studies with null findings were reviewed, it is possible that even more such studies have been carried out and were not published. Another concern is selective reporting of positive findings. Several articles in this review have yielded mixed results for the biomarkers studied, although it is possible that some authors only reported their statistically significant findings. Since rheumatic or widespread pain conditions were excluded from this review, it is conceivable that a bias was introduced which may have selected out more severe MSD that affect numerous anatomic sites concurrently – perhaps those that triggered an autoimmune response affecting widespread anatomical sites.

Recommendations

Our review shows that biochemical biomarker research for neck and upper-extremity MSD is at a nascent stage. Briefly, these are our recommendations for future biochemical biomarker research:

- Report the response rate for all analysis groups (nominally, for cases and referents).
- Use healthy control subjects where possible.
- Restrict subjects by duration and/or symptom range, or adjust for this in analyses.
- Select biomarkers identified in MSD animal models and/or proposed pathomechanisms or use previously investigated biomarkers in order to verify findings in new populations.
- Carefully consider and report potential confounders, gather information on these factors from study subjects, and potentially control for them through restriction or through adjustment in the analysis.
- Consider analysis by disorder severity.
- Prioritize biomarker studies that are longitudinal, include patients with early or intermediate stage disorders, and address patients with elbow, arm, or hand symptoms.

Concluding remarks

This comprehensive review used systematic methodology and demonstrated that although there are studies that show an association between biochemical biomarkers and upper-extremity MSD, the evidence is inconsistent. The most consistent studies suggest that 5-HT may be used as a biomarker for trapezius myalgia, and that TIMP and MMP/TIMP ratio may be used as biomarkers for selected fibroproliferative disorders. Further research is needed to confirm these conjectures. We have identified gaps in the literature and provided guidelines for future studies. Prospective high quality studies are needed to propel this field forward. Systematic biomarker selection and deliberate testing of biomarkers in

different MSD is recommended. This testing should be done with regard to duration and severity of the MSD. Selection criteria and response rates of all analysis groups should be detailed, and confounding factors described. It would be advisable for researchers to select biomarkers in future studies based on MSD animal models or theories of MSD pathophysiology.

Acknowledgement

This study was funded by a grant from the Swedish Research Council for Health, Working Life and Welfare [Forte Dnr. 2009–1761].

References

1. Hassard J, Teoh K, Cox T, Dewe P, Cosmar M, Grundler R, et al. Calculating the cost of work-related stress and psychosocial risks - a literature review. Luxembourg: Publications Office of the European Union: European Agency for Safety and Health at Work; 2014.
2. Huisstede BM, Bierma-Zeinstra SM, Koes BW, Verhaar JA. Incidence and prevalence of upper-extremity musculoskeletal disorders. A systematic appraisal of the literature. *BMC Musculoskelet Disord.* 2006;7:7. <http://dx.doi.org/10.1186/1471-2474-7-7>.
3. Health and Safety Executive (V2. Handling injuries). Crown Copyright. Available from: www.hse.gov.uk [accessed January 28, 2013]; 2012.
4. World Health Organization. Physical Inactivity: A Global Public Health Problem. Geneva: WHO; 2014. Available from: http://www.who.int/dietphysicalactivity/factsheet_inactivity/en/. [cited 2014 October 8, 2014].
5. Schneider E, Irastorza X. OSH in figures: Work-related musculoskeletal disorders in the EU — Facts and figures. Luxembourg: Publications Office of the European Union: European Agency for Safety and Health at Work; 2010.
6. Buckle PW, Devereux JJ. The nature of work-related neck and upper limb musculoskeletal disorders. *Appl Ergo.* 2002;33:207–17. [http://dx.doi.org/10.1016/S0003-6870\(02\)00014-5](http://dx.doi.org/10.1016/S0003-6870(02)00014-5).
7. National Academy of Sciences. Panel on Musculoskeletal Disorders and the Workplace, Commission on Behavioral and Social Sciences and Education, National Research Council. Musculoskeletal Disorders and the Workplace: Low Back and Upper Extremities. Washington, D.C.: National Academy Press; 2001.
8. Virta L, Joranger P, Brox JJ, Eriksson R. Costs of shoulder pain and resource use in primary health care: a cost-of-illness study in Sweden. *BMC Musculoskelet Disord.* 2012;13:17. <http://dx.doi.org/10.1186/1471-2474-13-17>.

9. Coyte PC, Asche CV, Croxford R, Chan B. The economic cost of musculoskeletal disorders in Canada. *Arthritis Care Res.* 1998;11(5):315–25. <http://dx.doi.org/10.1002/art.1790110503>.
10. Bureau of Labor Statistics. Prevention of work-related musculoskeletal disorders. Washington, DC: Bureau of Labor Statistics; 2014. Available from: <http://www.bls.gov/news.release/pdf/osh2.pdf>.
11. OSHA. Prevention of work-related musculoskeletal disorders: U.S. Department of Labor, Occupational Safety & Health Administration; 2014 [cited 2014 July 29]. Available from: https://www.osha.gov/pls/oshaweb/owadisp.show_document?p_table=UNIFIED_AGENDA&p_id=4481.
12. Viikari-Juntura E. New avenues in research on musculoskeletal disorders. *Scand J Work Environ Health.* 1999;26(6):564–68.
13. Punnett L, Gold JE. Work-related upper-extremity disorders: epidemiologic findings and unresolved questions. In: Johansson H, Djupsjöbacka M, Windhorst U, Passatore M, Ljubisavljevic M, Blair SJ, editors. *Chronic Work-Related Myalgia: Neuromuscular Mechanisms behind Work-Related Chronic Muscle Pain Syndromes*. Umea, Sweden: National Institute for Working Life; 2003.
14. Institute of Medicine. Evaluation of Biomarkers and Surrogate Endpoints in Chronic Disease. Micheel CM, Ball JR, editors. Washington, DC: The National Academies Press; 2010.
15. Mayeux R. Biomarkers: potential uses and limitations. *NeuroRx.* 2004;1(2):182–8. <http://dx.doi.org/10.1602/neurorx.1.2.182>.
16. Committee on Biological Markers of the National Research Council. Biological markers in environmental health research. *Environ Health Persp.* 1987;74:3–9.
17. Mastin JP, Henningsen GM, Fine LJ. Use of biomarkers of occupational musculoskeletal disorders in epidemiology and laboratory animal model development. *Scand J Work Environ Health.* 1992;18 Suppl 2:85–7.
18. Saxton JM. A review of current literature on physiological tests and soft tissue biomarkers applicable to work-related upper limb disorders. *Occup Med.* 2000;50(2):121–30. <http://dx.doi.org/10.1093/occmed/50.2.121>.
19. Visser B, van Dieën JH. Pathophysiology of upper-extremity muscle disorders. *J Electromyogr Kinesiol.* 2006;16(1):1–16. <http://dx.doi.org/10.1016/j.jelekin.2005.06.005>.
20. Gerdle B, Larsson B. Potential Muscle Biomarkers of Chronic Myalgia in Humans - A Systematic Review of Microdialysis Studies, Biomarker. Khan T, editor. Rijeka: Biomarker: InTech; 2012.
21. Barbe MF, Barr AE. Inflammation and the pathophysiology of work-related musculoskeletal disorders. *Brain Behav Immun.* 2006;20(5):423–9. <http://dx.doi.org/10.1016/j.bbi.2006.03.001>.
22. Gerdle B, Ghafouri B, Ernberg M, Larsson B. Chronic musculoskeletal pain: review of mechanisms and biochemical biomarkers as assessed by the microdialysis technique. *J Pain Res.* 2014;7:313–26. <http://dx.doi.org/10.2147/JPR.S59144>.
23. Boocock MG, Collier JM, McNair PJ, Simmonds M, Larmer PJ, Armstrong B. A framework for the classification and diagnosis of work-related upper-extremity conditions: systematic review. *Semin Arthritis Rheum.* 2009;38(4):296–311. <http://dx.doi.org/10.1016/j.semarthrit.2007.10.006>.
24. World Health Organization. Identification and control of work-related diseases. Geneva: World Health Organization, 1985 Contract No.: 714.
25. Bruns DE, Huth EJ, Magid E, Young DS. Toward a checklist for reporting of studies of diagnostic accuracy of medical tests. *Clin Chem.* 2000;46(7):893–5.
26. Whiting P, Rutjes AW, Reitsma JB, Bossuyt PM, Kleijnen J. The development of QUADAS: a tool for the quality assessment of studies of diagnostic accuracy included in systematic reviews. *BMC Med Res Methodol.* 2003;3:25. <http://dx.doi.org/10.1186/1471-2288-3-25>.
27. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ.* 2007;335(7624):806–8. <http://dx.doi.org/10.1136/bmj.39335.541782.AD>.
28. Altman DG. Statistical reviewing for medical journals. *Stat Med.* 1998;17(23):2661–74. [http://dx.doi.org/10.1002/\(SICI\)1097-0258\(19981215\)17:23<2661::AID-SIM33>3.0.CO;2-B](http://dx.doi.org/10.1002/(SICI)1097-0258(19981215)17:23<2661::AID-SIM33>3.0.CO;2-B).
29. Downs SH, Black N. The feasibility of creating a checklist for the assessment of the methodological quality both of randomised and non-randomised studies of health care interventions. *J Epidemiol Community Health.* 1998;52(6):377–84. <http://dx.doi.org/10.1136/jech.52.6.377>.
30. Slavin RE. Best evidence synthesis: an intelligent alternative to meta-analysis. *J Clin Epidemiol.* 1995;48(1):9–18. [http://dx.doi.org/10.1016/0895-4356\(94\)00097-A](http://dx.doi.org/10.1016/0895-4356(94)00097-A).
31. Sjogaard G, Rosendal L, Kristiansen J, Blangsted AK, Skotte J, Larsson B, et al. Muscle oxygenation and glycolysis in females with trapezius myalgia during stress and repetitive work using microdialysis and NIRS. *Eur J Appl Physiol.* 2010;108(4):657–69. <http://dx.doi.org/10.1007/s00421-009-1268-2>.
32. Andersson HI, Ejlertsson G, Leden I, Rosenberg C. Characteristics of subjects with chronic pain, in relation to local and widespread pain report. A prospective study of symptoms, clinical findings and blood tests in subgroups of a geographically defined population. *Scand J Rheumatol.* 1996;25(3):146–54. <http://dx.doi.org/10.3109/03009749609080005>.
33. Gumina S, Carbone S, Perugia D, Vestri AR, Postacchini F. Shoulder adhesive capsulitis in the early freezing phase: correlations between blood exams and Constant Score. *Musculoskelet Surg.* 2011;95 Suppl 1:S37–42. <http://dx.doi.org/10.1007/s12306-011-0121-z>.
34. Rechardt M, Shiri R, Karppinen J, Jula A, Heliovaara M, Viikari-Juntura E. Lifestyle and metabolic factors in relation to shoulder pain and rotator cuff tendinitis: a population-based study. *BMC Musculoskelet Disord.* 2010;11:165. <http://dx.doi.org/10.1186/1471-2474-11-165>.

35. Bunker TD, Esler CN. Frozen shoulder and lipids. *J Bone Joint Surg Br.* 1995;77(5):684–6.
36. Ohlsson K, Schutz A, Attewell R, Skerfving S. Selenium status in females with occupational cervico-brachial complaints. *Int Arch Occup Environ Health.* 1988;61(3):167–9. <http://dx.doi.org/10.1007/BF00381014>.
37. Kaergaard A, Hansen AM, Rasmussen K, Andersen JH. Association between plasma testosterone and work-related neck and shoulder disorders among female workers. *Scand J Work Environ Health.* 2000;26(4):292–8. <http://dx.doi.org/10.5271/sjweh.545>.
38. Akesson I, Schutz A, Horstmann V, Skerfving S, Moritz U. Musculoskeletal symptoms among dental personnel; - lack of association with mercury and selenium status, overweight and smoking. *Swed Dent J.* 2000;24(1-2):23–38.
39. Bovenzi M, D'Agostin F, Rui F, Ambrosi L, Zefferino R. Salivary endothelin and vascular disorders in vibration-exposed workers. *Scand J Work Environ Health.* 2008;34(2):133–41. <http://dx.doi.org/10.5271/sjweh.1224>.
40. Carp SJ, Barbe MF, Winter KA, Amin M, Barr AE. Inflammatory biomarkers increase with severity of upper-extremity overuse disorders. *Clin Sci (London).* 2007;112(5):305–14. <http://dx.doi.org/10.1042/CS20060050>.
41. Rechardt M, Shiri R, Lindholm H, Karppinen J, Viikari-Juntura E. Associations of metabolic factors and adipokines with pain in incipient upper-extremity soft tissue disorders: a cross-sectional study. *BMJ Open.* 2013;3(8):e003036. <http://dx.doi.org/10.1136/bmjopen-2013-003036>.
42. Rechardt M, Shiri R, Matikainen S, Viikari-Juntura E, Karppinen J, Alenius H. Soluble IL-1RII and IL-18 are associated with incipient upper-extremity soft tissue disorders. *Cytokine.* 2011;54:149–53. <http://dx.doi.org/10.1016/j.cyt.2011.02.003>.
43. Abboud JA, Kim JS. The effect of hypercholesterolemia on rotator cuff disease. *Clin Orthop Relat Res.* 2010;468(6):1493–7. <http://dx.doi.org/10.1007/s11999-009-1151-9>.
44. Lakemeier S, Braun J, Efe T, Foelsch C, Archontidou-Aprin E, Fuchs-Winkelmann S, et al. Expression of matrix metalloproteinases 1, 3, and 9 in differing extents of tendon retraction in the torn rotator cuff. *Knee Surg Sports Traumatol Arthrosc.* 2011;19(10):1760–5. <http://dx.doi.org/10.1007/s00167-010-1367-y>.
45. Lakemeier S, Reichelt JJ, Timmesfeld N, Fuchs-Winkelmann S, Paletta JR, Schofer MD. The relevance of long head biceps degeneration in the presence of rotator cuff tears. *BMC Musculoskelet Disord.* 2010;11:191. <http://dx.doi.org/10.1186/1471-2474-11-191>.
46. Longo UG, Franceschi F, Spiezia F, Forriol F, Maffulli N, Denaro V. Triglycerides and total serum cholesterol in rotator cuff tears: do they matter? *Br J Sports Med.* 2010;44(13):948–51. <http://dx.doi.org/10.1136/bjsm.2008.056440>.
47. Iqbal SA, Manning C, Syed F, Kolluru V, Hayton M, Watson S, et al. Identification of mesenchymal stem cells in perinodular fat and skin in Dupuytren's disease: a potential source of myofibroblasts with implications for pathogenesis and therapy. *Stem Cells Dev.* 2012;21(4):609–22. <http://dx.doi.org/10.1089/scd.2011.0140>.
48. Hnanicek J, Cimburova M, Putova I, Svoboda S, Stritesky J, Kratka K, et al. Lack of association of iron metabolism and Dupuytren's disease. *J Eur Acad Dermatol Venereol.* 2008;22(4):476–80. <http://dx.doi.org/10.1111/j.1468-3083.2007.02506.x>.
49. Ulrich D, Hrynyszyn K, Pallua N. Matrix metalloproteinases and tissue inhibitors of metalloproteinases in sera and tissue of patients with Dupuytren's disease. *Plast Reconstr Surg.* 2003;112(5):1279–86. <http://dx.doi.org/10.1097/01.PRS.0000081462.40448.49>.
50. Ulrich D, Ulrich F, Piatkowski A, Pallua N. Expression of matrix metalloproteinases and their inhibitors in cords and nodules of patients with Dupuytren's disease. *Arch Orthop Trauma Surg.* 2009;129(11):1453–9. <http://dx.doi.org/10.1007/s00402-008-0726-3>.
51. Castagna A, Cesari E, Gigante A, Conti M, Garofalo R. Metalloproteases and their inhibitors are altered in both torn and intact rotator cuff tendons. *Musculoskelet Surg.* 2013;97 Suppl 1:39–47. <http://dx.doi.org/10.1007/s12306-013-0264-1>.
52. Papalia R, Del Buono A, Leonardi F, Osti L, Maffulli N, Denaro V. Creatinine and nonprotein nitrogen plasma levels: possible etiopathogenetic factors in rotator cuff tears. *Phys Sportsmed.* 2011;39(2):127–32. <http://dx.doi.org/10.3810/psm.2011.05.1903>.
53. Dale AM, Novak CB, Mackinnon SE. Utility of vibration thresholds in patients with brachial plexus nerve compression. *Ann Plast Surg.* 1999;42:613–8. <http://dx.doi.org/10.1097/0000637-199906000-00005>.
54. Xu Y, Bonar F, Murrell GA. Neoinnervation in rotator cuff tendinopathy. *Sports Med Arthrosc.* 2011;19(4):354–9. <http://dx.doi.org/10.1097/JSA.0b013e318229d7e3>.
55. Xu Y, Bonar F, Murrell GA. Enhanced expression of neuronal proteins in idiopathic frozen shoulder. *J Shoulder Elbow Surg.* 2012;21(10):1391–7. <http://dx.doi.org/10.1016/j.jse.2011.08.046>.
56. Lubis AM, Lubis VK. Matrix metalloproteinase, tissue inhibitor of metalloproteinase and transforming growth factor-beta 1 in frozen shoulder, and their changes as response to intensive stretching and supervised neglect exercise. *J Orthop Sci.* 2013;18(4):519–27. <http://dx.doi.org/10.1007/s00776-013-0387-0>.
57. Riva R, Mork PJ, Westgaard RH, Lundberg U. Comparison of the cortisol awakening response in women with shoulder and neck pain and women with fibromyalgia. *Psychoneuroendocrinology.* 2012;37(2):299–306. <http://dx.doi.org/10.1016/j.psyneuen.2011.06.014>.
58. Tsujii M, Hirata H, Yoshida T, Imanaka-Yoshida K, Morita A, Uchida A. Involvement of tenascin-C and PG-M/versican in flexor tenosynovial pathology of idiopathic carpal tunnel syndrome. *Histol Histopathol.* 2006;21(5):511–8.
59. Kennedy G, Khan F, McLaren M, Belch JJ. Endothelial activation and response in patients with hand arm vibration syndrome. *Eur J Clin Invest.* 1999;29(7):577–81. <http://dx.doi.org/10.1046/j.1365-2362.1999.00502.x>.

60. Nakamura H, Matsuzaki I, Hatta K, Nagase H, Nobokuni Y, Kambayashi Y, et al. Blood endothelin-1 and cold-induced vasodilation in patients with primary Raynaud's phenomenon and workers with vibration-induced white finger. *Int Angiol.* 2003;22(3):243–9.
61. Flodgren GM, Crenshaw AG, Hellstrom F, Fahlstrom M. Combining microdialysis and near-infrared spectroscopy for studying effects of low-load repetitive work on the intramuscular chemistry in trapezius myalgia. *J Biomed Biotechnol.* 2010;2010:513803. <http://dx.doi.org/10.1155/2010/513803>.
62. Hadrevi J, Ghafouri B, Larsson B, Gerdle B, Hellstrom F. Multivariate modeling of proteins related to trapezius myalgia, a comparative study of female cleaners with or without pain. *PLoS One.* 2013;8(9):e73285. <http://dx.doi.org/10.1371/journal.pone.0073285>.
63. Lau CS, Bridges AB, Muir A, Scott N, Bancroft A, Belch JJ. Further evidence of increased polymorphonuclear cell activity in patients with Raynaud's phenomenon. *Br J Rheumatol.* 1992;31(6):375–80. <http://dx.doi.org/10.1093/rheumatology/31.6.375>.
64. Ghafouri B, Larsson BK, Sjors A, Leandersson P, Gerdle BU. Interstitial concentration of serotonin is increased in myalgic human trapezius muscle during rest, repetitive work and mental stress - an in vivo microdialysis study. *Scand J Clin Lab Invest.* 2010;70(7):478–86. <http://dx.doi.org/10.3109/00365513.2010.511257>.
65. Sjors A, Larsson B, Karlson B, Osterberg K, Dahlman J, Gerdle B. Salivary cortisol response to acute stress and its relation to psychological factors in women with chronic trapezius myalgia - a pilot study. *Psychoneuroendocrinology.* 2010;35(5):674–85. <http://dx.doi.org/10.1016/j.psyneuen.2009.10.007>.
66. Moreno-Torres A, Rosset-Llobet J, Pujol J, Fabregas S, Gonzalez-de-Suso JM. Work-related pain in extrinsic finger extensor musculature of instrumentalists is associated with intracellular pH compartmentation during exercise. *PLoS One.* 2010;5(2):e9091. <http://dx.doi.org/10.1371/journal.pone.0009091>.
67. Flodgren GM, Crenshaw AG, Alfredson H, Fahlstrom M, Hellstrom FB, Bronemo L, et al. Glutamate and prostaglandin E2 in the trapezius muscle of female subjects with chronic muscle pain and controls determined by microdialysis. *Eur J Pain.* 2005;9(5):511–5. <http://dx.doi.org/10.1016/j.ejpain.2004.11.004>.
68. Nilsen KB, Sand T, Westgaard RH, Stovner LJ, White LR, Bang Leistad R, et al. Autonomic activation and pain in response to low-grade mental stress in fibromyalgia and shoulder/neck pain patients. *Eur J Pain.* 2007;11(7):743–55. <http://dx.doi.org/10.1016/j.ejpain.2006.11.004>.
69. Teodorczyk-Injeyan JA, Triano JJ, McGregor M, Woodhouse L, Injeyan HS. Elevated production of inflammatory mediators including nociceptive chemokines in patients with neck pain: a cross-sectional evaluation. *J Manipulative Physiol Ther.* 2011;34(8):498–505. <http://dx.doi.org/10.1016/j.jmpt.2011.08.010>.
70. Freeland AE, Tucci MA, Barbieri RA, Angel MF, Nick TG. Biochemical evaluation of serum and flexor tenosynovium in carpal tunnel syndrome. *Microsurgery.* 2002;22(8):378–85. <http://dx.doi.org/10.1002/micr.10065>.
71. Larsson B, Rosendal L, Kristiansen J, Sjogaard G, Sogaard K, Ghafouri B, et al. Responses of algescic and metabolic substances to 8 h of repetitive manual work in myalgic human trapezius muscle. *Pain.* 2008;140(3):479–90. <http://dx.doi.org/10.1016/j.pain.2008.10.001>.
72. Rosendal L, Kristiansen J, Gerdle B, Sogaard K, Peolsson M, Kjaer M, et al. Increased levels of interstitial potassium but normal levels of muscle IL-6 and LDH in patients with trapezius myalgia. *Pain.* 2005;119(1-3):201–9. <http://dx.doi.org/10.1016/j.pain.2005.09.026>.
73. Augoff K, Ratajczak K, Gosk J, Tabola R, Rutowski R. Gelatinase A activity in Dupuytren's disease. *J Hand Surg Am.* 2006;31(10):1635–9. <http://dx.doi.org/10.1016/j.jhsa.2006.08.007>.
74. Gerdle B, Hilgenfeldt U, Larsson B, Kristiansen J, Sogaard K, Rosendal L. Bradykinin and kallidin levels in the trapezius muscle in patients with work-related trapezius myalgia, in patients with whiplash associated pain, and in healthy controls - A microdialysis study of women. *Pain.* 2008;139(3):578–87. <http://dx.doi.org/10.1016/j.pain.2008.06.012>.
75. Rosendal L, Larsson B, Kristiansen J, Peolsson M, Sogaard K, Kjaer M, et al. Increase in muscle nociceptive substances and anaerobic metabolism in patients with trapezius myalgia: microdialysis in rest and during exercise. *Pain.* 2004;112(3):324–34. <http://dx.doi.org/10.1016/j.pain.2004.09.017>.
76. Ghafouri N, Ghafouri B, Larsson B, Stensson N, Fowler CJ, Gerdle B. Palmitoylethanolamide and stearoylethanolamide levels in the interstitium of the trapezius muscle of women with chronic widespread pain and chronic neck-shoulder pain correlate with pain intensity and sensitivity. *Pain.* 2013;154(9):1649–58. <http://dx.doi.org/10.1016/j.pain.2013.05.002>.
77. Ozgocmen S, Ardicoglu O. Lipid profile in patients with fibromyalgia and myofascial pain syndromes. *Yonsei Med J.* 2000;41(5):541–5. <http://dx.doi.org/10.3349/ymj.2000.41.5.541>.
78. Bunker TD, Anthony PP. The pathology of frozen shoulder. A Dupuytren-like disease. *J Bone Joint Surg Br.* 1995;77(5):677–83.
79. Wynn TA. Common and unique mechanisms regulate fibrosis in various fibroproliferative diseases. *J Clin Invest.* 2007;117(3):524–9. <http://dx.doi.org/10.1172/JCI31487>.
80. Abate M, Schiavone C, Salini V, Andia I. Occurrence of tendon pathologies in metabolic disorders. *Rheumatology (Oxford).* 2013;52(4):599–608. <http://dx.doi.org/10.1093/rheumatology/kes395>.
81. Beason DP, Abboud JA, Kuntz AF, Bassora R, Soslowsky LJ. Cumulative effects of hypercholesterolemia on tendon biomechanics in a mouse model. *J Orthop Res.* 2011;29(3):380–3. <http://dx.doi.org/10.1002/jor.21255>.
82. Hart DA, Frank CB, Bray RC. Inflammatory processes in

- repetitive motion and overuse syndromes: potential role of neurogenic mechanisms in tendons and ligaments. In: Gordon SL, Blair SJ, Fine LJ, editors. Repetitive Motion Disorders of the Upper-extremity. Rosemont, IL: American Academy of Orthopaedic Surgeons; 1994. p. 247–62.
83. Vogel K. Fibrocartilage in tendon: a response to compressive load. In: Gordon SL, Blair SJ, Fine LJ, editors. Repetitive Motion Disorders of the Upper-extremity. Rosemont, IL: American Academy of Orthopaedic Surgeons; 1994.
84. Beamer B, Hettrich C, Lane J. Vascular endothelial growth factor: an essential component of angiogenesis and fracture healing. *HSS J*. 2010;6(1):85–94. <http://dx.doi.org/10.1007/s11420-009-9129-4>.
85. Hadjiargyrou M, Lombardo F, Zhao S, Ahrens W, Joo J, Ahn H, et al. Transcriptional profiling of bone regeneration. Insight into the molecular complexity of wound repair. *J Biol Chem*. 2002;277(33):30177–82. <http://dx.doi.org/10.1074/jbc.M203171200>.
86. Mulherin SA, Miller WC. Spectrum bias or spectrum effect? Subgroup variation in diagnostic test evaluation. *Ann Intern Med*. 2002;137(7):598–602. <http://dx.doi.org/10.7326/0003-4819-137-7-200210010-00011>.
87. Gao HG, Fisher PW, Lambi AG, Wade CK, Barr-Gillespie AE, Popoff SN, et al. Increased serum and musculotendinous fibrogenic proteins following persistent low-grade inflammation in a rat model of long-term upper-extremity overuse. *PLoS One*. 2013;8(8):e71875. <http://dx.doi.org/10.1371/journal.pone.0071875>.
88. Fedorczyk JM, Barr AE, Rani S, Gao HG, Amin M, Amin S, et al. Exposure-dependent increases in IL-1 β , substance P, CTGF, and tendinosis in flexor digitorum tendons with upper-extremity repetitive strain injury. *J Orthop Res*. 2010;28(3):298–307.
89. Kietrys DM, Barr-Gillespie AE, Amin M, Wade CK, Popoff SN, Barbe MF. Aging contributes to inflammation in upper-extremity tendons and declines in forelimb agility in a rat model of upper-extremity overuse. *PLoS One*. 2012;7(10):e46954. <http://dx.doi.org/10.1371/journal.pone.0046954>.
90. Barr AE, Barbe MF. Inflammation reduces physiological tissue tolerance in the development of work-related musculoskeletal disorders. *J Electromyogr Kinesiol*. 2004;14(1):77–85. <http://dx.doi.org/10.1016/j.jelekin.2003.09.008>.
91. Dinarello CA. Interleukin-1 in the pathogenesis and treatment of inflammatory diseases. *Blood*. 2011;117(14):3720–32. <http://dx.doi.org/10.1182/blood-2010-07-273417>.
92. Liuzzo G, Biasucci LM, Gallimore JR, Grillo RL, Rebuzzi AG, Pepys MB, et al. The prognostic value of C-reactive protein and serum amyloid a protein in severe unstable angina. *N Engl J Med*. 1994;331(7):417–24. <http://dx.doi.org/10.1056/NEJM199408183310701>.
93. Malik S, Wong ND, Franklin S, Pio J, Fairchild C, Chen R. Cardiovascular disease in U.S. patients with metabolic syndrome, diabetes, and elevated C-reactive protein. *Diabetes Care*. 2005;28(3):690–3. <http://dx.doi.org/10.2337/diacare.28.3.690>.
94. Beckman JA, Preis O, Ridker PM, Gerhard-Herman M. Comparison of usefulness of inflammatory markers in patients with versus without peripheral arterial disease in predicting adverse cardiovascular outcomes (myocardial infarction, stroke, and death). *Am J Cardiol*. 2005;96(10):1374–8. <http://dx.doi.org/10.1016/j.amjcard.2005.07.041>.
95. de Koning L, Liptak C, Shkreta A, Bradwin G, Hu FB, Pradhan AD, et al. A multiplex immunoassay gives different results than singleplex immunoassays which may bias epidemiologic associations. *Clin Biochem*. 2012;45(10-11):848–51. <http://dx.doi.org/10.1016/j.clinbiochem.2012.04.006>.
96. de Jager W, Bourcier K, Rijkers GT, Prakken BJ, Seyfert-Margolis V. Prerequisites for cytokine measurements in clinical trials with multiplex immunoassays. *BMC Immunol*. 2009;10:52. <http://dx.doi.org/10.1186/1471-2172-10-52>.
97. Astrom M, Rausing A. Chronic Achilles tendinopathy: A survey of surgical and histopathologic findings. *Clin Orthop*. 1995;316:151–64.
98. Andersson G, Danielson P, Alfredson H, Forsgren S. Presence of substance P and the neurokinin-1 receptor in tenocytes of the human Achilles tendon. *Regul Pept*. 2008;150(1-3):81–7. <http://dx.doi.org/10.1016/j.regpep.2008.02.005>.
99. Petersen W, Pufe T, Pfrommer S, Tillmann B. [Overload damage to the Achilles tendon: the importance of vascularization and angiogenesis]. *Orthopade*. 2005;34(6):533–42. <http://dx.doi.org/10.1007/s00132-005-0808-7>.
100. Saccenti E, Hoeflsloot HC, Smilde AK, Westerhuis JA, Hendriks MM. Reflections on univariate and multivariate analysis of metabolomics data. *Metabolomics*. 2014;10(3):361–74. <http://dx.doi.org/10.1007/s11306-013-0598-6>.
101. Brunnekreef JJ, Thijssen DH, Oosterhof J, Hopman MT. Bilateral changes in forearm oxygen consumption at rest and after exercise in patients with unilateral repetitive strain injury: a case-control study. *J Orthop Sports Phys Ther*. 2012;42(4):371–8. <http://dx.doi.org/10.2519/jospt.2012.3751>.
102. Pilgian GJ, Gold JE. The repetitive strain injury process: alterations in physiology vis-a-vis symptoms. *J Orthop Sports Phys Ther*. 2012;42(7):660; author reply 1.
103. Novak CB, Mackinnon SE. Multiple nerve entrapment syndromes in office workers. *Occup Med*. 1999;14(1):39–59.
104. Riva R, Mork Paul Jarle J, Westgaard Rolf Harald H, Lundberg U. Comparison of the cortisol awakening response in women with shoulder and neck pain and women with fibromyalgia. *Psychoneuroendocrinology*. 2011;37(2):299–306. <http://dx.doi.org/10.1016/j.psyneuen.2011.06.014>.
105. Sjörs A, Larsson B, Karlson B, Osterberg K, Dahlman J, Gerdle B. Salivary cortisol response to acute stress and its relation to psychological factors in women with chronic trapezius myalgia—a pilot study. *Psychoneuroendocrinology*. 2009;35(5):674–85. <http://dx.doi.org/10.1016/j.psyneuen.2009.10.007>.
106. Sjøgaard G, Zebis MK, Küllerich K, Saltin B, Pilegaard H. Exercise training and work task induced metabolic and stress-related mRNA and protein responses in myalgic muscles. *BioMed Res Inter*. 2013;2013:984523. <http://dx.doi.org/10.1155/2013/984523>.

107. Demirkol A, Uludag M, Soran N, Aksoy N, Gun K, Incebiyik S, et al. Total oxidative stress and antioxidant status in patients with carpal tunnel syndrome. *Redox Report*. 2012;17(6):234–8. <http://dx.doi.org/10.1179/1351000212Y.00000000027>.
108. Ghafouri N, Ghafouri B, Larsson B, Turkina Maria VV, Karlsson L, Fowler Christopher JJ, et al. High levels of N-palmitoylethanolamide and N-stearoylethanolamide in microdialysate samples from myalgic trapezius muscle in women. *PloS One*. 2011;6(11):e27257. <http://dx.doi.org/10.1371/journal.pone.0027257>.
109. Hallgren Hanna CBC, Eliasson P, Aspenberg P, Adolfsson Lars EE. Elevated plasma levels of TIMP-1 in patients with rotator cuff tear. *Acta Orthop*. 2012;83(5):523–8. <http://dx.doi.org/10.3109/17453674.2012.736174>.
110. Han SH, An HJ, Song JY, Shin DE, Kwon YD, Shim JS, et al. Effects of corticosteroid on the expressions of neuropeptide and cytokine mRNA and on tenocyte viability in lateral epicondylitis. *J Inflamm (United Kingdom)*. 2012 Oct 30;9(1):40. <http://dx.doi.org/10.1186/1476-9255-9-40>.
111. Hasselhorn HMM, Theorell T, Vingård E, Musculoskeletal Intervention Center N. Endocrine and immunologic parameters indicative of 6-month prognosis after the onset of low back pain or neck/shoulder pain. *Spine*. 2001;26(3):E24–9. <http://dx.doi.org/10.1097/00007632-200102010-00005>.
112. Hindocha S, Iqbal SAA, Farhatullah S, Paus R, Bayat A. Characterization of stem cells in Dupuytren's disease. *British J Surg*. 2011;98(2):308–15. <http://dx.doi.org/10.1002/bjs.7307>.
113. Hirata H, Nagakura T, Tsujii M, Morita A, Fujisawa K, Uchida A. The relationship of VEGF and PGE2 expression to extracellular matrix remodelling of the tenosynovium in the carpal tunnel syndrome. *J Surg*. 2004;204(5):605–12. <http://dx.doi.org/10.1002/path.1673>.
114. Hirata H, Tsujii M, Yoshida T, Yoshida KI, Morita A, Okuyama N, et al. MMP-2 expression is associated with rapidly proliferative arteriosclerosis in the flexor tenosynovium and pain severity in carpal tunnel syndrome. *J Pathol*. 2005;205:443–50. <http://dx.doi.org/10.1002/path.1709>.
115. Kim Yang-Soo S, Kim Jung-Man M, Lee Yun-Gyoung G, Hong Oak-Kee K, Kwon Hyuk-Sang S, Ji Jong-Hoon H. Intercellular adhesion molecule-1 (ICAM-1, CD54) is increased in adhesive capsulitis. *J Bone Joint Surg (Amer)*. 2013;95(4):e181–8. <http://dx.doi.org/10.2106/JBJS.K.00525>.
116. Lau CSS, O'Dowd A, Belch JJJ. White blood cell activation in Raynaud's phenomenon of systemic sclerosis and vibration induced white finger syndrome. *Ann Rheum Dis*. 1992;51(2):249–52. <http://dx.doi.org/10.1136/ard.51.2.249>.
117. Nakama K, Gotoh M, Yamada T, Mitsui Y, Yasukawa H, Imaizumi T, et al. Interleukin-6-induced activation of signal transducer and activator of transcription-3 in ruptured rotator cuff tendon. *J Inter Med Res*. 2006;34:624–31. <http://dx.doi.org/10.1177/147323000603400607>.
118. Nilsson PM, Kandell-Collén A, Andersson HI. Blood pressure and metabolic factors in relation to chronic pain. *Blood Press*. 1997;6:294–8. <http://dx.doi.org/10.3109/08037059709062085>.
119. Nowak J, Barregård L, Benthin G, Granung G, Wennmalm A. Thromboxane metabolite excretion in patients with hand-arm vibration syndrome. *Clin Physiol (Oxford, England)*. 1996;16(4):361–7. <http://dx.doi.org/10.1111/j.1475-097X.1996.tb00725.x>.
120. Sanderson PLL, Morris MAA, Stanley JKK, Fahmy NRR. Lipids and Dupuytren's disease. *J Bone Joint Surg (British)*. 1992;74(6):923–7.
121. Tajana MS, Murena L, Valli F, Passi A, Grassi FA. Correlations between biochemical markers in the synovial fluid and severity of rotator cuff disease. *Chir Organi Mov*. 2009;93 Suppl 1:S41–8. <http://dx.doi.org/10.1007/s12306-009-0004-8>.
122. Zeisig E, Ljung BOO, Alfredson H, Danielson P. Immunohistochemical evidence of local production of catecholamines in cells of the muscle origins at the lateral and medial humeral epicondyles: of importance for the development of tennis and golfer's elbow? *Br J Sports Med*. 2009;43(4):269–75. <http://dx.doi.org/10.1136/bjsm.2008.054619>.
123. Bergenudd H, Lindgärde F, Nilsson BEE. Prevalence of Dupuytren's contracture and its correlation with degenerative changes of the hands and feet and with criteria of general health. *J Hand Surg (Edinburgh, Scotland)*. 1993;18(2):254–7. [http://dx.doi.org/10.1016/0266-7681\(93\)90123-W](http://dx.doi.org/10.1016/0266-7681(93)90123-W).
124. Hyvönen P, Melkko J, Lehto VPP, Jalovaara P. Involvement of the subacromial bursa in impingement syndrome of the shoulder as judged by expression of tenascin-C and histopathology. *J Bone Joint Surg (British)*. 2003;85(2):299–305. <http://dx.doi.org/10.1302/0301-620X.85B2.13124>.
125. Keniston RCC, Nathan PAA, Leklem JEE, Lockwood RSS. Vitamin B6, vitamin C, and carpal tunnel syndrome. A cross-sectional study of 441 adults. *J Occup Environ Med*. 1997;39(10):949–59. <http://dx.doi.org/10.1097/00043764-199710000-00007>.
126. Laskar Shawkatuzzaman MM, Iwamoto M, Nakamoto M, Koshiyama H, Harada N. Heart rate variation and urinary catecholamine excretion in response to acute psychological stress in hand-arm vibration syndrome patients. *J Occup Health*. 2004;46(2):125–31. <http://dx.doi.org/10.1539/joh.46.125>.
127. Okumus M, Ceceli E, Tuncay F, Kocaoglu S, Palulu N, Yorgancioglu ZR. The relationship between serum trace elements, vitamin B12, folic acid and clinical parameters in patients with myofascial pain syndrome. *J Back Musc Rehab*. 2010;23:187–91.
128. Olausson P, Gerdle B, Ghafouri N, Larsson B, Ghafouri B. Identification of Proteins from Interstitium of Trapezius Muscle in Women with Chronic Myalgia Using Microdialysis in Combination with Proteomics. *PloS One*. 2012;7. <http://dx.doi.org/10.1371/journal.pone.0052560>.
129. Ratajczak-Wielgomas K, Gosk J, Rabczyński J, Augoff K, Podhorska-Okołów M, Gamian A, et al. Expression of MMP-2, TIMP-2, TGF-β1, and decorin in Dupuytren's contracture. *Conn Tissue Res*. 2012;53:469–77. <http://dx.doi.org/10.3109/03008207.2012.686542>.
130. Toesca A, Pagnotta A, Zumbo A, Sadun R. Estrogen and

- progesterone receptors in carpal tunnel syndrome. *Cell Biol Inter.* 2008;32:75–9. <http://dx.doi.org/10.1016/j.cellbi.2007.08.014>.
131. Kang Jong-Won W, Kim H, Cho Soo-Hun H, Lee Myung-Koo K, Kim Yong-Dae D, Nan Hong-Mei M, et al. The association of subjective stress, urinary catecholamine concentrations and PC game room use and musculoskeletal disorders of the upper limbs in young male Koreans. *J Korean Med Sci.* 2003;18(3):419–24. <http://dx.doi.org/10.3346/jkms.2003.18.3.419>.
132. Kent PJJ, Williams GAA, Kester RCC. Platelet activation during hand vibration. *Brit J Surg.* 1994;81(6):815–8. <http://dx.doi.org/10.1002/bjs.1800810608>.
133. Meek RMM, McLellan S, Crossan JFF. Dupuytren's disease. A model for the mechanism of fibrosis and its modulation by steroids. *J Bone Joint Surg (Brit).* 1999;81(4):732–8. <http://dx.doi.org/10.1302/0301-620X.81B4.9163>.
134. Riondino S, La Farina F, Martini F, Guadagni F, Ferroni P. Functional impairment in video terminal operators is related to low-grade inflammation. *Int Arch Occup Environ Health.* 2010;84(7):745–51. <http://dx.doi.org/10.1007/s00420-010-0605-4>.
135. Chamblor A, Rawlinson S, Emery R, Pitsillides A. Quantitative cytochemical evidence for local increases in bone turnover at the acromial enthesis of the human coracoacromial ligament. *J Rheum.* 2004;31(11):2216–25.
136. Greenstein D, Kester RCC. The role of leukocytes in the pathogenesis of vibration-induced white finger. *Angiol.* 1998;49(11):915–22. <http://dx.doi.org/10.1177/000331979804901107>.
137. Ljung Björn-Ove O, Alfredson H, Forsgren S. Neurokinin 1-receptors and sensory neuropeptides in tendon insertions at the medial and lateral epicondyles of the humerus. Studies on tennis elbow and medial epicondylalgia. *J Orthop Res.* 2004;22(2):321–7. [http://dx.doi.org/10.1016/S0736-0266\(03\)00183-9](http://dx.doi.org/10.1016/S0736-0266(03)00183-9).
138. Osawa T, Shinozaki T, Takagishi K. Multivariate analysis of biochemical markers in synovial fluid from the shoulder joint for diagnosis of rotator cuff tears. *Rheumatol Inter.* 2005;25(6):436–41. <http://dx.doi.org/10.1007/s00296-004-0509-2>.
139. Tucci MAA, Barbieri RAA, Freeland AEE. Biochemical and histological analysis of the flexor tenosynovium in patients with carpal tunnel syndrome. *Biomed Sci Instru.* 1997;33:246–51.
140. Gudmundsson KGG, Arngrimsson R, Arinbjarnarson S, Olafsson A, Jonsson T. T-and B-lymphocyte subsets in patients with Dupuytren's disease. Correlations with disease severity. *J Hand Surg (Edinburgh, Scotland).* 1998;23(6):724–7. [http://dx.doi.org/10.1016/S0266-7681\(98\)80083-1](http://dx.doi.org/10.1016/S0266-7681(98)80083-1).
141. Harada N, Kondo H, Kimura K. Assessment of autonomic nervous function in patients with vibration syndrome using heart rate variation and plasma cyclic nucleotides. *Brit J Indus Med.* 1990;47(4):263–8. <http://dx.doi.org/10.1136/oem.47.4.263>.
142. Ljung BOO, Forsgren S, Fridén J. Substance P and calcitonin gene-related peptide expression at the extensor carpi radialis brevis muscle origin: implications for the etiology of tennis elbow. *J Orthop Res.* 1999;17(4):554–9. <http://dx.doi.org/10.1002/jor.1100170414>.
143. Yoshihara Y, Hamada K, Nakajima T, Fujikawa K, Fukuda H. Biochemical markers in the synovial fluid of glenohumeral joints from patients with rotator cuff tear. *J Orthop Res.* 2001;19(4):573–9. [http://dx.doi.org/10.1016/S0736-0266\(00\)00063-2](http://dx.doi.org/10.1016/S0736-0266(00)00063-2).
144. Donato G, Galasso O, Valentino P, Conforti F, Zuccalà V, Russo E, et al. Pathological findings in subsynovial connective tissue in idiopathic carpal tunnel syndrome. *Clin Neuropath.* 2009;28(2):129–35. <http://dx.doi.org/10.5414/NPP28129>.
145. Kohout J, Topolcan O, Bejcková H. The serum level of endothelin in patients with hand-arm vibration syndrome. *Cent Eur J Public Health.* 1995;3 Suppl:43–4.
146. Santavirta S, Kontinen YTT, Antti-Poika I, Nordström D. Inflammation of the subacromial bursa in chronic shoulder pain. *Arch Orthop Trauma Surg.* 1992;111(6):336–40. <http://dx.doi.org/10.1007/BF00420062>.

Received for publication: 4 May 2015