Scand J Work Environ Health 2016;42(1):86-90

<https://doi.org/10.5271/sjweh.3541>

Published online: 10 Dec 2015, Issue date: 01 Jan 2016

The 18th Collegium Ramazzini statement: The global health dimensions of asbestos and asbestos-related diseases

by [Collegium Ramazzini](#)

Affiliation: Collegium Ramazzini, via Saliceto 3 40010 Bentivoglio (BO), Italy. collegium@ramazzini.it

Key terms: [asbestos](#); [asbestos-containing product](#); [asbestos-related disease](#); [cancer](#); [Collegium Ramazzini](#); [health](#)

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/26686520

This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

The 18th Collegium Ramazzini statement: The global health dimensions of asbestos and asbestos-related diseases

The Collegium Ramazzini reaffirms its long-standing position that responsible public health action is to ban all extraction and use of asbestos, including chrysotile. This current statement updates the Collegium's earlier statements with a focus on global health dimensions of asbestos and asbestos-related diseases (ARD). The ARD epidemic will likely not peak for at least a decade in most industrialized countries and for several decades in industrializing countries. Asbestos and ARD will continue to present challenges in the arena of occupational medicine and public health, as well as in clinical research and practice, and have thus emerged as a global health issue. Industrialized countries that have already gone through the transition to an asbestos ban have learned lessons and acquired know-how and capacity that could be of great value if deployed in industrializing countries embarking on the transition. The accumulated wealth of experience and technologies in industrialized countries should thus be shared internationally through global campaigns to eliminate ARD.

Background

Every mined asbestos fiber is indestructible. As a result, many individuals are repeatedly exposed during the silicate mineral's lifecycle, from mining and extraction of asbestos-containing rocks to manufacturing of asbestos-containing products (ACP), and further during use, repair, demolition and abatement of ACP. Since 1993, the Collegium has repeatedly called for a global ban on all mining, manufacture, and use of asbestos (1–4). The Collegium has taken this position based on well-validated scientific evidence showing that all types of asbestos, including chrysotile the most widely used form, cause cancers (such as mesothelioma and lung cancer) and additionally that there is no safe level of exposure. The Collegium has continued to criticize as fallacious and unachievable the so-called “controlled use” of chrysotile advocated by the asbestos industry. Unfortunately, despite these concerns and abundant scientific evidence, global usage of chrysotile has remained at around two million metric tons per year in recent years. Most of this current use is concentrated in low- and middle-income countries (5).

The Collegium reaffirms its position that, given the well-documented availability of safer, cost-effective

alternative materials, the responsible public health action is to ban all extraction and use of asbestos. State-of-the-art technologies must be employed in asbestos removal and disposal. This current statement updates earlier statements with a focus on the global health dimensions of asbestos and asbestos-related diseases (ARD).

UN Organizations

In 2006, the World Health Organization (WHO) called for the elimination of ARD (6), taking the position that the most efficient way to eliminate such diseases is to cease using all types of asbestos. The 2014 update of this statement, which was attached to the WHO document “Chrysotile Asbestos” (7), published in response to the continuing widespread production and use of chrysotile, emphasized that all forms of asbestos, including chrysotile, are causally associated with an increased risk of cancer of the lung, larynx and ovary, mesothelioma and asbestosis. These observations are in line with the recent evaluation by the International Agency for Research on Cancer (IARC) (8). In its 2014 update, the WHO reiterated its call for global campaigns to eliminate ARD. These efforts have been joined by other UN agencies including the International Labour Organization (ILO) and the United Nations Environment Programme (UNEP). The Chemical Review Committee of the Rotterdam Convention has repeatedly recommended that chrysotile asbestos be put on the Convention's list of hazardous substances, thus requiring exporting countries to obtain prior informed consent (PIC) from the importing countries. A handful of countries have opposed that recommendation, thus preventing this basic safety protection from coming into effect. The Collegium calls on all Parties to the Rotterdam Convention to support the listing of chrysotile asbestos as a hazardous substance.

Global burden of asbestos-related diseases

Occupational exposure to asbestos causes an estimated 107 000 deaths each year worldwide. These deaths result from asbestos-related lung cancer (ARLC), mesothelioma and asbestosis (6, 7). When the global burden of each type

of ARD was considered separately, the estimated number of deaths per year was 41 000 for ARLC (9), 43 000 (10) – 59 000 (7, 9, 11) for mesothelioma, and 7000 (12) – 24 000 (13) for asbestosis. No estimate is available for the annual numbers of deaths due to asbestos-related cancers of the larynx or ovary. Because asbestos is more likely to cause lung cancer than mesothelioma, the total burden of ARD will differ substantially with the estimated magnitude of ARLC. The WHO recently advanced a risk ratio of 6:1 for contracting lung cancer versus mesothelioma following chrysotile exposure (7). As these estimates are derived by different methods, inconsistencies might be eliminated through a cross-verification of the various estimation methods used. Regardless, the ARD burden is more likely to be under- than overestimated because ARD are well known to be under-diagnosed and reported.

National bans

Since Iceland first introduced a ban on all types of asbestos in 1983, more than 50 countries have implemented similar bans (14). However, the pace of countries adopting bans has slowed in the past decade. Indeed, the governments of several industrializing countries have withdrawn bans while others have prescribed long periods over which to move towards a ban. Such actions are likely a consequence of the corrupting influence of pro-chrysotile lobbies, whether foreign or domestic. Asbestos industry lobbyists employ “product defense” science to foment uncertainty to sway the opinions of industrializing countries, a delaying tactic which, unfortunately, often succeeds. Nine of the ten most populous countries in the world, all of which use or have used substantial amounts of asbestos, have yet to adopt bans. Coverage of the world population by bans thus remains low and is biased towards industrialized countries.

Alternatives to asbestos

In countries where asbestos has been banned, safer, cost-effective substitute materials have been successfully introduced. Polyvinyl alcohol fibers and cellulose fibers can be used instead of asbestos in building products such as flat and corrugated fiber-cement sheets, which are used in roofing, interior walls, and ceilings. Polypropylene and cellulose fibers have been used instead of asbestos to make fiber-cement products in Brazil. Virtually all of the polymeric and cellulose fibers used instead of asbestos in fiber-cement sheets are >10 microns in diameter and hence non-respirable. For roofing in remote locations, lightweight concrete tiles can be

fabricated using cement, sand and gravel and, optionally, locally available plant fibers such as jute, hemp, sisal, palm nut, coconut coir, kenaf, and wood pulp. Galvanized iron roofing and clay tiles are other alternative materials. Substitutes for asbestos-cement pipe include ductile iron pipe, high-density polyethylene pipe, and metal-wire-reinforced concrete pipes (15, 16). While these materials are considered safer than asbestos, good work practices should be observed for the protection of those working with these materials.

Patterns of an asbestos-related diseases epidemic

Countries continuing to use asbestos will shoulder the burden of ARD in proportion to their prior levels of asbestos use (17). Countries where asbestos has been banned or greatly limited invariably exhibit a sustained epidemic of ARD. Age-adjusted mortality rates of mesothelioma are increasing in most industrialized countries (18) but the rate of increase has slowed in only a few industrialized countries, which started to reduce asbestos use decades ago. With the known synergy of asbestos and smoking, it can be expected that the many industrializing countries with high smoking prevalence and continued use of asbestos will shoulder a substantial burden of asbestos-related lung cancer. The ARD epidemic will likely not peak for at least a decade in most industrialized countries and for several decades in industrializing countries. Asbestos and ARD will therefore continue to present challenges in the arena of occupational medicine and public health as well as in clinical research and practice. Hence, asbestos and ARD are global health issues.

Industrializing countries

Many industrializing countries have been slow to reduce, let alone ban, the use of asbestos. The multiple factors at play include the low price and easy accessibility of asbestos, demand from the construction sector in emerging economies, scarcity of medicosocial resources, and fierce propaganda by the asbestos industry and other parties with conflicting interests. These factors are interrelated and converge uniquely in each country, presenting significant challenges to concerned parties. For example, a number of rapidly growing industrializing countries in Asia and former Soviet Union countries currently sustain a high level of asbestos use and/or production and they fail to provide even minimal protection to workers; they have a serious lack of expertise and resources required to diagnose and report ARD. Furthermore, several industrializing coun-

tries that were importers (but not exporters) of asbestos were among the countries that opposed the inclusion of chrysotile into the aforementioned PIC procedure of the Rotterdam Convention. This is a blatant reflection of the corrupt influence of the asbestos industry and crude trade pressures of asbestos-exporting countries. Advocates for banning asbestos must continue to strive to overcome the reluctance, denial, and antagonism of their opponents.

Industrialized countries

The highest priority in reducing ARD is primary prevention, that is banning asbestos use in countries where it remains legal and preventing exposure to *in situ* sources in all countries with historical asbestos use. In industrialized countries, large quantities of asbestos remain as a legacy from past construction practices in many thousands of schools, homes, and commercial buildings. Significant quantities of asbestos also remain in various industrial applications. It is of importance to document and mark existing asbestos in buildings and industrial applications to avoid exposure during maintenance, repair, and demolition. As the materials weather, erode, break, or are cut by power tools, asbestos fibers are released into the air, soil and water, where they become a source of community-wide exposure. Policies, regulations and practices should safeguard workers engaged in the removal of asbestos-containing structures and the handling of the resulting waste material, via schemes for specialized training and licensing (19).

Secondary and tertiary prevention are also assuming vital importance in industrialized countries. In particular, workers exposed to asbestos in current or past occupations should be identified, registered, and followed-up for health monitoring and surveillance (19). The unfolding ARD epidemic in these countries poses costly challenges in the arenas of basic and clinical medicine. In medical practice, such challenges include the development of biomarkers for the early detection of mesothelioma, as well as effective modalities for its treatment. It is imperative to design and implement fair compensation schemes for people with ARD and their families. Industrialized countries should provide assistance to industrializing countries on issues related to asbestos and ARD.

In countries having banned asbestos, as well as in countries still using asbestos, a large number of workers remain at high risk of developing ARD from past exposure, in particular lung cancers and mesotheliomas. Most of these previously exposed people remain in the general population without any ongoing health monitoring. The Collegium recommends that countries develop

strategies for identifying their previously and currently asbestos-exposed workers to quantify their exposure and register them, subsequently developing methods for continuous health surveillance and secondary prevention (20). In addition to workers, there should be monitoring of household members of workers if they bring asbestos into their homes.

International cooperation

The accumulated wealth of experience and technologies in industrialized countries should be shared internationally through global campaigns to eliminate ARD. Industrialized countries have experience in primary, secondary, and tertiary prevention, with the strengths of any given country depending on its particular stage in their epidemic of ARD. The knowledge and technological developments that have emerged from these experiences could be of great benefit to countries in which asbestos continues to be used. The Statement on asbestos by the International Commission on Occupational Health (ICOH) (21) describes a broad range of activities at each of the three levels of prevention. For optimum effect, the resources of industrialized countries should be combined and distributed in a manner tailored to the needs of the beneficiaries. Scientific expertise, including capacity building and surveillance of ARD, is an important resource to be shared. Given the wide range of problems encountered at the global level, the development of regional initiatives should be particularly valuable (22).

Industrialized countries that have already transitioned to an asbestos ban have learned lessons and acquired know-how and capacity (ie, “soft” technology) that could be of great value if deployed in industrializing countries embarking on the transition. Collaboration between industrialized and industrializing countries can be led by international organizations, the scientific community and/or grassroots NGO, and should involve practitioners, researchers, administrators, and civil society. For example, through fora such as international workshops or conferences, countries with bans in place can outline how they implemented a ban and provide practical guidance on how countries currently using asbestos can move towards a ban.

Conclusion - the need for a global health approach

Asbestos and ARD have emerged as global health issues. All countries with a history of asbestos use are experiencing an epidemic of ARD, with the stage of the

epidemic being a function of the country's past asbestos use, whether and when it implemented a ban, and, if no ban is in place, at what levels it continues to use the material. Gaps in human capital and technology available to countries warrant international cooperation. The expansion of national bans in industrializing countries and reducing the burden of ARD in industrialized countries are the short-term targets. Given that ARD are 100% preventable, zero new cases of ARD should be the ultimate goal for both industrializing and industrialized countries. The pandemic of ARD is an urgent international priority for action by public health workers.

References

1. Collegium Ramazzini. Third Collegium Ramazzini Statement (1993). Chrysotile as a Carcinogen. [http://collegiumramazzini.org/download/3_ThirdCRStatement\(1993\).pdf](http://collegiumramazzini.org/download/3_ThirdCRStatement(1993).pdf)
2. Collegium Ramazzini. Sixth Collegium Ramazzini Statement (1999). Call for an International Ban on Asbestos. [http://collegiumramazzini.org/download/6_SixthCRStatement\(1999\).pdf](http://collegiumramazzini.org/download/6_SixthCRStatement(1999).pdf)
3. Collegium Ramazzini. Eleventh Collegium Ramazzini Statement (2004). Call for an International Ban on Asbestos: Statement Update. [http://collegiumramazzini.org/download/11_EleventhCRStatement\(2004\).pdf](http://collegiumramazzini.org/download/11_EleventhCRStatement(2004).pdf)
4. Collegium Ramazzini. Collegium Ramazzini Statement (2010, 15th). Asbestos is Still with Us: Repeat Call for a Universal Ban. [http://collegiumramazzini.org/download/15_FifteenthCRStatement\(2010\).pdf](http://collegiumramazzini.org/download/15_FifteenthCRStatement(2010).pdf)
5. United States Geological Survey. 2013 Minerals Yearbook: Asbestos [Advance Release]. <http://minerals.usgs.gov/minerals/pubs/commodity/asbestos/myb1-2013-asbes.pdf>
6. World Health Organization. Elimination of Asbestos-related Diseases. Geneva: WHO; 2006. http://whqlibdoc.who.int/hq/2006/WHO_SDE_OEH_06.03_eng.pdf?ua=1
7. World Health Organization. Chrysotile Asbestos. Geneva: WHO; 2014. http://www.who.int/ipcs/assessment/public_health/chrysotile_asbestos_summary.pdf
8. International Agency for Research on Cancer. IARC Monographs Volume 100C: Arsenic, Metals, Fibres and Dusts; A Review of Human Carcinogens. 2012. <http://monographs.iarc.fr/ENG/Monographs/vol100C/mono100C.pdf>
9. Prüss-Ustün A, Vickers C, Haeffliger P, Bertollini R. Knowns and unknowns on burden of disease due to chemicals: a systematic review. *Environ Health*. 2011;10:9. <http://dx.doi.org/10.1186/1476-069X-10-9>
10. Driscoll T, Nelson DI, Steenland K, Leigh J, Concha-Barrientos M, Fingerhut M, Prüss-Ustün A. The global burden of disease due to occupational carcinogens. *Am J Ind Med*. 2005;48(6):419-31. <http://dx.doi.org/10.1002/ajim.20209>
11. Global health risks: mortality and burden of disease attributable to selected major risks. Geneva: World Health Organization, 2009. http://www.who.int/healthinfo/global_burden_disease/GlobalHealthRisks_report_full.pdf
12. Driscoll T, Nelson DI, Steenland K, Leigh J, Concha-Barrientos M, Fingerhut M, Prüss-Ustün A. The global burden of non-malignant respiratory disease due to occupational airborne exposures. *Am J Ind Med*. 2005;48(6):432-45. <http://dx.doi.org/10.1002/ajim.20210>
13. GBD 2013 Mortality and Causes of Death Collaborators. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2015;385:117-71. [http://dx.doi.org/10.1016/S0140-6736\(14\)61682-2](http://dx.doi.org/10.1016/S0140-6736(14)61682-2)
14. International Ban Asbestos Secretariat. Chronology of National Asbestos Bans. Compiled by Laurie Kazan-Allen. http://www.ibasecretariat.org/chron_ban_list.php
15. World Bank Group. Good Practice Note: Asbestos: Occupational and Community Health Issues. Washington DC: WBG; May 2009. <http://siteresources.worldbank.org/EXTPOPS/Resources/AsbestosGuidanceNoteFinal.pdf>
16. World Health Organization Regional Office for Europe. National Programmes for Elimination of Asbestos-Related Diseases: Review and Assessment 07-08 June 2011, Bonn. Annex 4: Review of substitutes for asbestos construction products by a WHO-temporary advisor, pp 22-29. http://www.euro.who.int/_data/assets/pdf_file/0005/176261/National-Programmes-For-Elimination-Of-Asbestos-related-Diseases-Review-And-Assessment.pdf
17. Lin RT, Takahashi K, Karjalainen A, Hoshuyama T, Wilson D, Kameda T, Chan CC, Wen CP, Furuya S, Higashi T, Chien LC, Ohtaki M. Ecological association between asbestos-related diseases and historical asbestos consumption: an international analysis. *Lancet*. 2007;369(9564): 844-9. [http://dx.doi.org/10.1016/S0140-6736\(07\)60412-7](http://dx.doi.org/10.1016/S0140-6736(07)60412-7)
18. Delgermaa V, Takahashi K, Park E-K, Le GV, Hara T, Sorahan T. Global mesothelioma deaths reported to the World Health Organization between 1994 and 2008. *Bull World Health Organ*. 2011; 89: 716-724C. doi:10.2471/BLT.11.086678. <http://dx.doi.org/10.2471/BLT.11.086678>
19. Finnish Institute of Occupational Health and International Commission on Occupational Health. The Helsinki Declaration on Management and Elimination of Asbestos-Related Diseases. Adopted by the International Conference on Monitoring and Surveillance of Asbestos-Related Diseases, 10-13 February 2014, Espoo, Finland. http://www.ttl.fi/en/international/conferences/helsinki_asbestos_2014/Documents/20%20March%202014%20Final%20Signed%20Declaration%20for%20website.pdf
20. Langård S. Identification and prevention of work- and environment-related individual *à priori* disease risks. In: The Identification and Control of Environmental and Occupational Diseases. Part II. (Mehlman MA, Upton A, eds). *Adv Mod Environ Toxicol*. 1994;23:21-32.
21. International Commission on Occupational Health. ICOH Statement: Global Asbestos Ban and the Elimination of Asbestos-Related Diseases. 2013. http://www.icohweb.org/site_new/multimedia/news/pdf/ICOH%20Statement%20on%20global%20asbestos%20ban.pdf

22. Marsili D, Comba P, Pasetto R, Terracini B. International scientific cooperation on asbestos-related disease prevention in Latin America. *Ann Glob Health* 80(4): 247-50, 2014. <http://dx.doi.org/10.1016/j.aogh.2014.09.002>

The Collegium Ramazzini is an international scientific society that examines critical issues in occupational and environmental medicine with a view towards action to prevent disease and promote health. The Collegium derives its name from Bernardino Ramazzini, the father of occupational medicine, a professor of medicine of the Universities of Modena and Padua in the late 1600s and the early 1700s. The Collegium is comprised of 180 physicians and scientists from 35 countries, each of whom is elected to membership. The Collegium is independent of commercial interests.

Correspondence to: General Secretary, Dr. Morando Soffritti, Collegium Ramazzini, via Saliceto 3 40010 Bentivoglio (BO), Italy. [E-mail: collegium@ramazzini.it]