



Original article

Scand J Work Environ Health 2017;43(5):405-414

doi:10.5271/sjweh.3656

Work-related biomechanical exposure and job strain in midlife separately and jointly predict disability after 28 years: a Finnish longitudinal study

by [Prakash KC](#), [Neupane S](#), [Leino-Arjas P](#), [von Bonsdorff MB](#), [Rantanen T](#), [von Bonsdorff ME](#), [Seitsamo J](#), [Ilmarinen J](#), [Nygård C-H](#)

Higher level of work-related biomechanical exposure and job strain in midlife separately and jointly carried a higher risk for increase in disability after 28 years. Mitigation of both of these co-occurring exposures at work in midlife could reduce the risk of disability in later life. Thus the workplace should be promoted as an arena for preventive interventions regarding disability in old age.

Affiliation: School of Health Sciences, University of Tampere, FI-33014, Tampere, Finland. Prakash.kc@uta.fi

Refers to the following texts of the Journal: [2009;35\(2\):96-112](#)
[2013;39\(5\):468-476](#) [2014;40\(4\):335-342](#)

Key terms: [ADL](#); [ageing](#); [biomechanical exposure](#); [disability](#); [exposure](#); [Finland](#); [Finland](#); [IADL](#); [interaction](#); [job strain](#); [longitudinal study](#); [occupation](#); [occupational exposure](#); [occupational exposure](#); [old age](#)

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/28653077

Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](#).



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

Work-related biomechanical exposure and job strain in midlife separately and jointly predict disability after 28 years: a Finnish longitudinal study

by Prakash KC, MSc (Public Health),^{1,2} Subas Neupane, PhD,^{1,2} Päivi Leino-Arjas, PhD,³ Mikaela B von Bonsdorff, PhD,^{4,5} Taina Rantanen, PhD,⁴ Monika E von Bonsdorff, PhD,^{4,5} Jorma Seitsamo, PhD,³ Juhani Ilmarinen, PhD,⁶ Clas-Håkan Nygård, PhD^{1,2}

KC P, Neupane S, Leino-Arjas P, von Bonsdorff MB, Rantanen T, von Bonsdorff ME, Seitsamo J, Ilmarinen J, Nygård C-H. Work-related biomechanical exposure and job strain in midlife separately and jointly predict disability after 28 years: a Finnish longitudinal study. *Scand J Work Environ Health*. 2017;43(5):405–414. doi:10.5271/sjweh.3656

Objectives We investigated whether the extent of biomechanical exposures and job strain in midlife separately and jointly predict disability in old age.

Methods Participants of the Finnish Longitudinal Study on Aging Municipal Employees (FLAME) in 1981 (aged 44–58 years) responded to disability questionnaires in 2009 (1850 women and 1082 men). Difficulties in performing five activities of daily living (ADL) and seven instrumental ADL (IADL) were used to assess severity of disability (score range: 0–12, 0=no disability). Information on biomechanical exposures and job strain was collected by questionnaire at baseline. Adjusted prevalence proportion ratios (PR) and 95% confidence intervals (95% CI) were modelled using mixed negative binomial regression with robust variance. The joint effect of two exposures was quantified using the concept of relative excessive risk due to interaction (RERI).

Results The overall prevalence of disability (score: 1–12) was 46.7% (women: 41%; men: 57%). Compared to low-level exposures in an adjusted model, the PR of high baseline biomechanical exposures for each one unit increase in the disability score was 1.31 (95% CI 1.10–1.55) and PR of high job strain was 1.71 (95% CI 1.26–2.32). Associations were rather similar in gender-stratified analyses. Furthermore, the joint effect (high strain/high biomechanical) was multiplicative (women: PR 1.32, 95% CI 1.21–1.45; men: PR 1.27, 95% CI 1.13–1.44), but no additive effect was observed when fully adjusted.

Conclusion High biomechanical exposure and job strain in midlife were strongly associated with the severity of disability in later life. The workplace could serve as arena for preventive interventions regarding disability in old age.

Key terms ADL; ageing; Finland; IADL; interaction; occupation; occupational exposure; old age.

Adequate performance of activities of daily living (ADL) is essential for independent living (1) and difficulty in performing ADL and instrumental ADL (IADL) tasks is commonly used in assessing old age disability (1–3). Physical frailties are known predictors of disability among older people (4, 5). The occupational class in midlife also plays a vital role in the onset of later life disability, with unskilled blue-collar workers being the high-risk group compared to white-collar workers (6).

However, work-related biomechanical and psychosocial exposures are undeniably related to work disability among all types of occupations (7), exposures such as monotonous work, whole-body vibration, heavy physical work, prolonged standing, and low level of support being the substantially associated ones.

High occupational physical loading activity in midlife could likewise result in limitations in physical functioning in later life (5, 8–11) which could, eventu-

¹ School of Health Sciences, University of Tampere, Tampere, Finland.

² Gerontology Research Center, University of Tampere, Tampere, Finland.

³ Finnish Institute of Occupational Health, Helsinki, Finland.

⁴ Gerontology Research Center and Department of Health Sciences, University of Jyväskylä, Jyväskylä, Finland.

⁵ Folkhälsan Research Center, Helsinki, Finland.

⁶ Juhani Ilmarinen Consulting Ltd, Helsinki, Finland.

ally progress to disability. The work-related psychosocial exposure during midlife has persistent effects into old age and substantially predicts disability pension in later life (12). There is variation in the type of occupations of women and men, so the extent and nature of work-related exposures and effects may vary by gender (13). The joint effect of these exposures is equally crucial to detect the probable health impact as they are co-occurring exposures of midlife. A study of the GAZEL cohort found an additive effect of the joint biomechanical and psychosocial occupational exposures on post-retirement disability (14) through the interaction effect of the exposures. The same study reported gender-specific variations of the joint effects of the exposures.

The longitudinal evidence on individual associations of biomechanical and psychosocial occupational hazards with disabilities in later life, along with the evidence on the prediction of disability by joint biomechanical and psychosocial exposures is scarce. For the additional examination of the rare joint effect along with the separate effect, we aimed to investigate whether work-related biomechanical exposure and job strain (psychosocial exposure) in midlife separately and jointly predict disability in later life in a cohort of Finnish public sector employees followed up for 28 years.

Methods

Participants and design

This study is based on the Finnish Longitudinal Study on Aging Municipal Employees (FLAME), a prospective follow-up study which was conducted by the Finnish Institute of Occupational Health. A baseline survey was conducted in 1981 with the latest follow-up in 2009. The baseline cohort consisted of 6257 of 7344 (85.2% response rate) public sector employees aged 44–58 years and was representative of the largest municipal occupational groups in Finland (15–17). The study population of the present study comprised individuals who replied at baseline and in 2009 on information about disability, gathered by postal questionnaires (N=2932 complete data on ADL and IADL tasks). According to the mortality data (January 1981 to July 2009) obtained from the Finnish National Population Register, 33.2% of the baseline respondents died during the 28-year follow-up period. The ethics committee of the Finnish Institute of Occupational Health, Helsinki, Finland approved the study.

Disability

The assessment of ADL and IADL tasks was done by questionnaires distributed among the participants

in 2009. Difficulty in performing ADL and IADL is widely used as a measure of disability among the older people (2, 18–20). Five ADL tasks (eating, going to bed, dressing, bathing, and toileting) and seven IADL tasks (preparing meals, doing laundry, shopping, doing light domestic work, dosing/taking medicine, using the phone, and management of personal finances) partly adapted from Katz ADL (21) were used. The listed tasks were assessed on a scale 0–4, (4=manage without difficulties, 3=manage with little difficulties, 2=manage with lots of difficulties, 1=cannot manage without help from others, and 0=cannot manage even with help of others). In this analysis, responses were dichotomized (supplementary table A, www.sjweh.fi/index.php?page=data-repository). Then the dichotomized responses were summed up to get a continuous score 0–12. Score 0 represented those who could perform all the tasks without any difficulty (classified as non-disabled in this study) and those who had at least some difficulty in performing one or more of the 12 tasks scored 1–12 depending on the number of tasks entailing difficulties (classified as disabled in this study). The higher the score the more severe the disability (3).

Biomechanical exposure

The assessment of work-related biomechanical exposure was done using the self-reported responses to seven questions related to exposures to biomechanical hazards. The seven self-reported items were (i) continuous walking or movement, (ii) standing in one place, (iii) bent or twisted postures, (iv) similar repeated movements, (v) carrying objects by hand, (vi) sudden strenuous efforts and (vii) other poor postures. All the items were answered on a scale 0=not at all, 1=a little, 2=somewhat, 3=often, and 4=quite often, except item (vii) other poor posture which was answered on 0=not at all, 1=a little, 2=somewhat or 3=often. The summary score (Cronbach's $\alpha=0.84$) ranging from 0–26 was dichotomized into low (summary score=0–12) and high (summary score=13–26) biomechanical exposure using the median value 12 (22).

Job strain

Job control. Self-reported responses to ten questions related to respondents' possibilities for control, guidance, and influence on the job were used to assess job control: including (i) receiving guidance on the job, (ii) participating in planning the work, (iii) gaining promotion, (iv) having chances for further training in professional skills, (v) gaining recognition and respect, (vi) influencing the work environment, (vii) having chances to use own abilities and talents, (viii) having the aptitude to learn new things, (ix) communicating and working with co-workers,

and (x) seeing the meaning of the work were answered on a scale from 0=not at all, 1=a little, 2=somewhat, or 3=a lot. The summary score (Cronbach's $\alpha=0.86$) ranging from 0–30 was dichotomized into low (summary score=0–16) and high (summary score=17–30) job control using the median value 16 (17, 22).

Job demands. The assessment of job demands was based on the responses of the respondents to eight questions about pressure and demands related to the job. The eight self-reported items were namely: (i) tight time schedule, (ii) hectic pace of work, (iii) taking responsibility, (iv) pressure and interference from supervisor, (v) conflicting demands regarding work tasks and responsibility, (vi) pressure of failure or making mistakes, (vii) isolation or loneliness, and (viii) monotonous work, which were answered on a scale from 0=not at all, 1= a little, 2=somewhat, or 3=a lot. The summary score (Cronbach's $\alpha=0.77$) ranging from 0–24 was dichotomized into low (summary score=0–6) and high (summary score=7–24) job demands using the median value 6 (17, 22).

Job strain. Job demand–control concept from the Karasek model was used to create job strain levels (23). The dichotomized values of job demands and control were used to construct the four levels of job strain: namely low strain (high control and low demands), passive job (low control and low demands), active job (high control and high demands), and high strain (low control and high demands).

Covariates

The covariates used in this study were occupational class, age, gender, alcohol intake, smoking status, body mass index (BMI; kg/m²), physical activity, major chronic diseases and tenure period were assessed in baseline. Two occupational groups namely white- and blue-collar workers were created being based on 13 job profiles, which were a cluster of 133 different job titles (17). The tenure period used in this study was occupationally active years between the baseline of this study and retirement. Those reporting smoking at least one cigarette per day were classified as current smokers. Information on alcohol intake was classified into 0, ≤ 2 drinks a month, and ≥ 1 drink a week being based on seven original categories. Physical activity during leisure time in the previous year was classified in this study as active, moderately active, and inactive. Information on major chronic diseases was related to the diagnosis of chronic diseases of different body systems.

Statistical analysis

The descriptive characteristics of the subjects were first cross tabulated as frequencies and percentages accord-

ing to biomechanical exposure and job strain, with significance level of $P<0.05$. An interaction term between gender and exposures was tested and found statistically significant ($P<0.001$) with respect to disability and therefore the analyses were stratified by gender. Disability was over-dispersed so mixed negative binomial regression with robust variances was used to calculate prevalence proportion ratios (PR) and 95% confidence intervals (95% CI). Negative binomial regression was used in order to avoid errors related to overestimations. The estimates for job strain are presented as pairwise comparisons (Bonferroni adjusted). Tenure period from the baseline survey until retirement was also used in the model as an exposure item. Three models were fitted namely; model I was adjusted for age and tenure period, model II was further adjusted for occupational groups, and model III was further adjusted for smoking status, alcohol intake, physical activity, BMI, and chronic diseases.

The potential interaction between biomechanical exposure and job strain was quantified to check the joint prediction of the exposures. PR and 95% CI of multiplicative interaction and relative excessive PR and 95% CI for additive interactions were estimated using the advanced prefix known as "icp" from Stata, which calculates the estimates of log relative risk to use interaction contrast from the regression settings. The setting uses the concepts of relative excessive risk due to interaction ($RERI = RR_{1,1} - RR_{1,0} - RR_{0,1} + 1$) to calculate the estimates (24). In the given formula, $RR_{1,1}$ signifies the presence of high biomechanical and high job strain (joint exposure), $RR_{1,0}$ signifies presence of high biomechanical and absence of high job strain and $RR_{0,1}$ signifies absence of high biomechanical and presence of high job strain. Additive interaction is meant to be present if $RERI > 0$ with statistical significance. Two models were fitted, the first model was adjusted for age and tenure period and the second model was further adjusted for occupational groups, BMI, smoking status, chronic diseases, alcohol intake and physical activity. The overall test of the hypothesis and fit of the models are presented in supplement table D (for pairwise comparisons of job strain) and supplement table E (RERI), www.sjweh.fi/index.php?page=data-repository.

In order to account for potential bias of the study due to dropout, we did a sensitivity analysis for which we conducted a regression imputation. Information on disability was imputed for those who were alive and who failed to respond to the disability-related questionnaires (N=1100) and for that we used the most recent responses (follow-up of 1992 or 1997) from these respondents on chronic conditions, sum score of mobility limitation [squatting, bending, maintaining body position for 2 hours, lifting objects >10 kg, precise hand use, running 100 meters, walking 2 km and climbing three floors

based on the International Classification of Functioning (ICF)], workability, occupational class and age. We were able to impute the information on disability for 521 out of 1100 missing responses. The analysis for separate and combined ADL and IADL disability were almost identical, therefore only combined results are presented. Likewise, there was no difference in the pattern of associations in the results produced by the sensitivity analyses with imputed data and those derived using the original data. Therefore, the results are presented for the original respondents only. In order to examine the effect of mortality on the results, we made a secondary analysis where we included the deceased respondents in the highest disability group. The results were comparable to the original results (supplementary table F, www.sjweh.fi/index.php?page=data-repository). Proportion/percentage calculation and cross-tabulation were conducted in SPSS version 23.0 (IBM corporation, Armonk, New York, USA) and all the other analyses and plotting were done in STATA 14.0 (StataCorp LP, College Station, Texas 77845, USA).

Results

Tables 1 and 2 present the distribution of baseline characteristics according to biomechanical exposure and job strain among 1850 women and 1082 men, respectively. The overall prevalence of disability was 46.7% (severity score: 1–12), which was 41% and 57% among women and men, respectively. The baseline characteristics according to biomechanical exposure and job strain among non-respondent women (N=1542) and men (N=1630) are presented in supplement tables B and C, www.sjweh.fi/index.php?page=data-repository. PR and 95% CI for biomechanical exposure and job strain (pairwise comparisons) predicting disability are presented separately for all, men and women in table 3 in three different models. In the age-adjusted model, high baseline biomechanical exposure and high job strain was significantly associated (model I) with a unit increase in the severity of disability in later life compared to low-level exposures. The estimates were attenuated when

Table 1a. Baseline characteristics of women according to job strain and work-related biomechanical exposures, Finnish Longitudinal study on Aging Municipal Employees (FLAME), 1981–2009.

Baseline characteristics	N=1850		Biomechanical exposure				Job strain							
			Low (N=849)		High (N=1001)		Low (N=615)		Passive (N=443)		Active (N=349)		High (N=443)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Age (years)	49.6	3.4	49.7	3.4	49.5	3.3	49.7	3.5	50.0	3.4	49.4	3.2	49.5	3.4
Body mass index	25.0	3.4	24.6	3.4	25.3	3.3	24.7	3.4	25.1	3.5	25.0	3.4	25.4	3.2
Tenure period	9.3	3.6	9.8	3.6	9.0	3.6	9.3	3.8	9.2	3.6	9.5	3.2	9.3	3.7

Table 1b. Baseline characteristics of women (continued).

Baseline characteristics	N=1850		Biomechanical exposure				Job strain						
			Low (N=849)		High (N=1001)		Low (N=615)		Passive (N=443)		Active (N=349)		High (N=443)
	N	%	%	%	%	%	%	%	%	%	%	%	
Occupational class													
White collar	1256	68	87	52	84	53	77	53					
Blue collar	594	32	13	48	16	47	23	47					
Chronic diseases													
No	637	35	43	27	46	31	29	26					
Yes	1212	65	57	73	54	69	71	74					
Physical activity													
Active	943	52	53	52	56	47	57	48					
Moderately active	750	41	40	42	37	47	37	45					
Inactive	117	7	7	6	7	6	6	7					
Smoking status													
Never	1473	80	76	83	79	80	79	80					
Former	229	12	15	10	13	12	13	11					
Current	148	8	9	7	8	8	8	9					
Alcohol intake													
0	1596	87	85	87	90	90	82	83					
≤2 drinks/month	190	10	11	10	8	8	14	13					
≥1drink/week	55	3	4	3	2	2	4	4					

further adjusted for occupational groups (model II). When adjusted further for BMI, alcohol intake, chronic diseases, smoking status, and physical activity during previous years (model III), the estimates were mitigated, but the direction of association remained unchanged. High versus low biomechanical exposure had on average 49% and 37% higher likelihood, respectively for women and men for a unit increase in severity of disability. Likewise, high versus low strain carried significantly higher likelihood for a unit increase in severity of disability among women (PR 1.47, 95% CI 1.08–2.01) and men (PR 2.15, 95% CI 1.42–3.26), respectively.

The joint prediction of disability in old age by job strain and biomechanical exposure in midlife are presented in figure 1 (all respondents) and figure 2 (gender stratified) respectively in two different models. In the model adjusted for age, the passive/high, active/high, high/low, low/high and high/high combinations of the categories of jobs strain and biomechanical exposures were on average highly associated with a unit increase in severity of disability in old age compared to low/low

combination (model I) among all subjects and among men. Among women only passive/high, active/high, low/high, and high/high were highly associated (model I). Further adjustment for chronic diseases, occupational class, and other covariates in model II attenuated the estimates, but did not change the direction of associations among all subjects, women and men. The estimates for joint (high/high) job strain and biomechanical exposures described higher multiplicative interactions among all (PR 1.28, 95% CI 1.19–1.38), women (1.32, 1.21–1.45), and men (1.27, 1.13–1.44) in the final model compared to the low/low combination.

RERI estimates for the additive interaction of job strain and biomechanical exposure are shown in table 4 in two models. The joint prediction of high strain and high biomechanical exposure was additive among women in model I (RERI 0.27, 95% CI 0.04–0.49) but was attenuated and the direction was changed after further adjustment in model II.

Table 2a. Baseline characteristics of men according to job strain and work-related biomechanical exposures, Finnish Longitudinal study on Aging Municipal Employees (FLAME), 1981–2009.

Baseline characteristics	N=1082		Biomechanical exposure				Job strain							
			Low (N=743)		High (N=348)		Low (N=355)		Passive (N=275)		Active (N=156)		High (N=296)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Age (years)	49.6	3.4	49.6	3.3	49.7	3.4	49.7	3.5	50.1	3.4	49.3	3.3	49.3	3.1
Body mass index	25.9	2.8	25.9	2.8	25.8	2.9	25.7	2.8	25.7	2.9	26.0	2.9	26.2	2.7
Tenure period	9.5	3.8	9.8	3.8	8.9	3.7	10.2	3.9	9.1	3.6	9.7	3.9	8.8	3.6

Table 2b. Baseline characteristics of men (continued).

Baseline characteristics	N=1082		Biomechanical exposure				Job strain							
			Low (N=743)		High (N=348)		Low (N=355)		Passive (N=275)		Active (N=156)		High (N=296)	
	N	%	%	%	%	%	%	%	%	%	%	%	%	
Occupational class														
White collar	422	39	53	10	56	21	58	25						
Blue collar	660	61	47	90	44	79	42	75						
Chronic diseases														
No	391	36	42	24	46	33	36	28						
Yes	691	64	58	76	54	67	64	72						
Physical activity														
Active	570	53	56	48	58	48	53	52						
Moderately active	442	42	40	44	36	45	45	42						
Inactive	53	5	4	8	5	6	2	6						
Smoking status														
Never	412	38	39	35	38	38	38	39						
Former	489	45	45	46	47	47	35	46						
Current	181	17	16	19	15	15	27	15						
Alcohol intake														
0	554	52	52	50	52	57	45	50						
≤2 drinks/month	356	33	32	35	31	30	36	36						
≥1drink/week	164	15	16	15	17	13	19	14						

Table 3. Prevalence proportion ratios (PR) and 95% confidence intervals (95% CI) for the association between work-related exposures (biomechanical and job strain) and later life disability, Finnish Longitudinal study on Aging Municipal Employees (FLAME), 1981–2009.

Work-related exposures	All		Women		Men	
	PR	95 % CI	PR	95 % CI	PR	95 % CI
Biomechanical ^a						
High vs low	1.62	1.38–1.91	1.98	1.59–2.47	1.75	1.36–2.24
Job strain ^{a, b}						
Passive vs low	1.72	1.27–2.32	1.40	0.95–2.06	2.16	1.40–3.35
Active vs low	1.50	1.09–2.07	1.25	0.82–1.92	2.12	1.31–3.43
High vs low	2.31	1.70–3.14	2.15	1.40–3.32	2.55	1.67–3.90
Active vs passive	0.87	0.63–1.21	0.90	0.58–1.38	0.98	0.61–1.58
High vs passive	1.34	0.99–1.82	1.54	1.02–2.34	1.18	0.77–1.80
High vs active	1.54	1.10–2.16	1.72	1.08–2.75	1.21	0.75–1.93
Biomechanical ^c						
High vs low	1.35	1.14–1.60	1.69	1.33–2.14	1.47	1.13–1.92
Job strain ^{b, c}						
Passive vs low	1.44	1.07–1.95	1.22	0.82–1.80	1.88	1.20–2.94
Active vs low	1.49	1.08–2.05	1.22	0.80–1.86	2.15	1.34–3.45
High vs low	1.94	1.42–2.65	1.86	1.20–2.87	2.24	1.44–3.48
Active vs passive	1.03	0.74–1.43	0.99	0.65–1.54	1.14	0.69–1.88
High vs passive	1.35	0.99–1.83	1.52	1.00–2.30	1.19	0.78–1.82
High vs active	1.30	0.93–1.83	1.53	0.95–2.44	1.04	0.63–1.71
Biomechanical ^d						
High vs low	1.31 ^e	1.10–1.55	1.49	1.17–1.89	1.37	1.06–1.78
Job strain ^{b, d}						
Passive vs low	1.43 ^e	1.06–1.93	1.13	0.76–1.69	1.99	1.26–3.13
Active vs low	1.37 ^e	0.99–1.90	1.08	0.70–1.67	2.08	1.30–3.34
High vs low	1.71 ^e	1.26–2.32	1.48	0.97–2.24	2.15	1.39–3.34
Active vs passive	0.96 ^e	0.68–1.35	0.95	0.61–1.49	1.05	0.62–1.76
High vs passive	1.19 ^e	0.89–1.61	1.30	0.87–1.96	1.08	0.71–1.64
High vs active	1.24 ^e	0.88–1.75	1.37	0.86–2.17	1.03	0.62–1.71

^a Model I: adjusted for age, tenure period.

^b Bonferroni adjusted pairwise comparisons.

^c Model II: adjusted for age, tenure period and occupational group.

^d Model III: adjusted for age, tenure period, occupational group, and other covariates (chronic diseases, physical activity, alcohol intake, body mass index and smoking status).

^e Adjusted for items in model III including gender.

Discussion

The findings of this longitudinal study of a cohort of Finnish public sector employees provides a strong evidence that high job strain and biomechanical exposure in working life separately and jointly predict the severity of disability in later life. The higher level of both exposures was decidedly deleterious in both genders. The interactions of the high/high level of exposures compared to low-level in midlife was highly associated with a unit increase in severity of disability in later life among both women and men.

Although occupational activity and exposures is a crucial part of the daily lives of most adults, studies on occupational exposures as a potential source of ADL and IADL disability later in life are scarce. Only a limited number of earlier studies have acknowledged work-related exposures as the long-term risk factors of old age disability and impaired functioning (6, 14, 20, 25, 26). Heavy physical load and high job demands




Figure 1. Prevalence proportion ratios (PR) and 95% confidence intervals (95% CI) for the association of later life disability and different combinations of joint work-related exposures (job strain and biomechanical exposure) of midlife among all subjects; model I: adjusted for age and tenure period; model II: further adjusted for gender, occupational group, chronic diseases, body mass index, alcohol intake, smoking status and physical activity; low/low category of exposures serve as referent (ref); Finnish Longitudinal study on Aging Municipal Employees (FLAME), 1981–2009.

were reported to be a risk factor for a decline in physical functioning (27) and functional limitations (28) among the Dutch cohorts. Biomechanical exposure and job strain in working life significantly predicted post-retirement disability among the GAZEL cohort (14) and work disability among Norwegian employees (6, 29). Likewise, higher level of work-related exposures was found to be long-term risk factors for musculoskeletal disability (30), disability pensioning (12, 31–33), and other chronic conditions (34). The findings of the present study corroborate these earlier findings, however, the respondents of our study were older and the outcome of interest was disability related to ADL.

Even though studies on the association between work-related exposures in working life and disability are limited, there is a lack of consistency in the findings. The results of our study contradict the findings of a Swedish study, which reported that occupational physical activity had no significant association with ADL disability (35). In our study, biomechanical exposure, which includes frequent occupational physical activities, was associated with high likelihood of higher disability. The risk of disability and disorders depends on the extent of exposures and varies by occupational class (6, 16, 36). Physically demanding work in midlife was found to be associated with ADL disability among women in a study in Ireland (26). Similarly, longest held manual occupations carried high risk of ADL disability among a Taiwanese cohort (6) and the risk of disability pensioning in Swedish construction workers (37). Consequently, our findings




Figure 2. Prevalence proportion ratios (PR) and 95% confidence intervals (95% CI) for the association of later life disability and different combinations of joint work-related exposures (job strain and biomechanical exposure) of midlife among women and men; model I: adjusted for age and tenure period; model II: further adjusted for occupational group, chronic diseases, body mass index, alcohol intake, smoking status and physical activity; low/low category of exposures serve as referent (ref); Finnish Longitudinal study on Aging Municipal Employees (FLAME), 1981–2009.

are parallel with these earlier studies because the high risk of disability was attenuated but persisted after we adjusted our results for occupational class (white- versus blue-collar) in model II. Work-related exposures as risk factors for future health outcomes may vary by gender (14, 16, 25, 38). The gender specific variation was clearly seen in our findings. The independent likelihood of higher disability due to high biomechanical exposure was higher among women and due to high strain was higher among men, which are parallel to the findings of (38).

An interaction effect is equally important along with the independent effects of work-related exposure on health functioning outcomes (33). Biomechanical exposure and job strain in working life jointly predicted post-retirement disability in the GAZEL cohort (14). Likewise, elevated risk of musculoskeletal disability was associated with combined high physical and psychosocial work exposures in a working population in Sweden (38) and in the UK (39). Our findings are in line with these earlier studies, although there were some differences in the follow-up period, methodological

considerations, the age of the respondents, and type of disability. The joint effect was biological with a significant RERI of 0.32 in the GAZEL cohort (14), and we found a RERI of 0.27 among women for the high/high combination of exposures in the age-adjusted model, but it was not significant in the fully adjusted model. Similarly, the statistical estimate of joint exposure (risk ratio) was 3.6 for musculoskeletal disability among women in a Swedish cohort (38) and 1.91 in a mixed population in the GAZEL cohort (14) and the same estimate was 1.28 (PR) in our study. These variations could be explained by the differences in methods and outcome of interest. The exposure of interest used in (14, 38) was cumulative, but we used the exposures from baseline only. Furthermore, the outcome in those studies were more related to musculoskeletal-pain-related disabilities compared to ADL in our study. Our study has added some epidemiological evidence to the current body of literature regarding the risk of disability in old age associated with the joint work-related exposures in midlife. Prevention of high work-related exposures in midlife could be significant not only for better physical

Table 4. Estimates of excessive prevalence of higher disability due to interaction between job strain and biomechanical exposure expressed as relative excessive risk due to interaction (RERI) and 95% confidence interval (95% CI) for the association between joint work-related exposures (biomechanical and job strain) and later life disability, Finnish Longitudinal study on Aging Municipal Employees (FLAME), 1981–2009.

Interactions	All		Women		Men	
	RERI	95 % CI	RERI	95 % CI	RERI	95 % CI
Job strain / biomechanical exposure						
Passive / High						
Model I ^a	0.05	-0.09–0.19	0.07	-0.12–0.26	0.07	-0.19–0.33
Model II ^b	0.05 ^c	-0.19–0.09	-0.21	-0.41– -0.01	0.16	-0.07–0.39
Active / High						
Model I ^a	0.22	0.06–0.38	0.57	0.38–0.77	-0.10	-0.42–0.21
Model II ^b	0.11 ^c	-0.05–0.28	0.38	0.18–0.58	-0.14	-0.44–0.16
High / High						
Model I ^a	0.09	-0.09–0.25	0.27	0.04–0.49	0.06	-0.22–0.34
Model II ^b	-0.06 ^c	-0.22–0.10	-0.04	-0.26–0.19	0.05	-0.18–0.29

^a Adjusted for age and tenure period.

^b Adjusted for age, tenure period, occupational group, smoking status, alcohol intake, body mass index, physical activity and chronic diseases.

^c Adjusted for items in model II including gender.

functioning and good health in midlife, but also and equally for better health outcomes in later life as well (25). Thus midlife is the best phase of life to start prevention of health problems of old age (5). Most of our findings highlight the significance of working conditions and low work-related exposures in midlife not only for proper physical functioning and good health in midlife but also for better physical functioning and reduction of the risk of disability in later life as well. Future studies should shed more light on the interactive predictability of these work-related exposures using different exposure models.

Strengths and limitations

One of the strengths of this study is a long prospective follow-up of almost three decades. Another strength is that our study involved a large representative sample of diverse occupations. Exploring the predictors of old age disability, a public health indicator among the rapidly growing elderly population is another potential strength. The information on retirement and mortality were taken from the national register rather than other sources, which increases the reliability of the study. This study contributes to the existing literature demanding interaction studies of longitudinal design investigating the interactions of work-related exposures in midlife to predict functional limitations and disability in old age. Among those who were alive, the response rate was significantly higher in all waves and almost half of the respondents from 1981 also responded in 2009 and the non-respondents were older than the respondents.

A potential limitation of the design was the availability of disability data from the last survey (2009) only and its self-reported nature, which could have possibly been a subject of information and reporting bias. Nonetheless, the assessment of ADL task and using the severity of difficulty in performing the task are taken as a valid and widely used measurement of old age health and functional status (18). Although both exposure variables were self-reported and were subject to reporting bias, we used the widely accepted Karasek model to create job strain. On the other hand, previous findings indicate at correlation between subjective and expert ratings on work-related exposures (40). The measurement of the exposure variable at a single time point at baseline could be the subject of an underestimation of the association, but we believe that the exposure level did not change much during the follow-up because almost 3/4 of the respondents did not change their jobs for >15 years (41). Furthermore, no major changes in their work were seen from the baseline survey until their retirement (41). The other drawback could be the lack of data on items like social support and supportive leadership at work, which has been shown to be associated with disability (8). The respondents in our study worked in the public sector, but the findings are relevant to private sector employees as well because in Finland the labor legislation applies equally to all sectors and occupations. The results could be generalized to countries with labor policies similar to those in Finland.

Concluding remarks

Job strain and biomechanical exposure in working life separately and jointly predicted disability in later life. The higher the level of work-related biomechanical exposures and job strain in midlife the higher was the chance of increase in the severity of disability in later life. These findings imply the need for mitigating both of these exposures at work through proper workplace interventions that could help in beginning to prevent old age disability already in working age.

Acknowledgements

Monika E von Bonsdorff was funded by the Academy of Finland (grant number 294530; 307114; 303920); Mikaela B von Bonsdorff was funded by EU H2020-PHC-2014-DynaHealth (grant number 633595). Taina Rantanen was funded by Academy of Finland (grant numbers 132597; 130285; 255403). Päivi Leino-Arjas and Jorma Seitsamo were funded by a grant from the Social Insurance Institution, Finland (grant number 53/26/2013).

The contents of this article do not necessarily exemplify the views of the funding institutions and are exclusively the responsibility of the authors. The authors declare no conflicts of interest.

References

- Tas U, Verhagen AP, Bierma-Zeinstra S MA, Oding E, Koes BW. Prognostic factors of disability in older people: a systematic review. *Br J Gen Pract.* 2007;57(537):319–23.
- Chan KS, Kasper JD, Brandt J, Pezzin LE. Measurement Equivalence in ADL and IADL difficulty across international surveys of aging: Findings from the HRS, SHARE, and ELSA. *J Gerontol B Psychol Sci Soc Sci.* 2012;67B(1):121–32. <https://doi.org/10.1093/geronb/gbr133>.
- Prakash KC, Neupane S, Leion-Arjas P, von Bonsdorff MB, Rantanen T, von Bonsdorff ME, et al. Midlife job profiles and disabilities in later life: a 28-year follow-up of municipal employees in Finland. *Int Arch Occup Environ Health.* 2016;89(6):997–1007. <https://doi.org/10.1007/s00420-016-1133-7>.
- den Ouden ME, Schuurmans MJ, Arts IE, van der Schouw YT. Physical performance characteristics related to disability in older persons: A systematic review. *Maturitas.* 2011;69(3):208–19. <https://doi.org/10.1016/j.maturitas.2011.04.008>.
- Vermeulen J, Neyens J CL, van Rossum E, Spreeuwenberg MD, de Witte LP. Predicting ADL disability in community-dwelling elderly people using physical frailty indicators: a systematic review. *BMC Geriatrics.* 2011;11:33. <https://doi.org/10.1186/1471-2318-11-33>.
- Li C, Wu SC, Wen SW. Longest held occupation in a lifetime and risk of disability in activities of daily living. *Occup Environ Med.* 2000;57(8):550–4. <https://doi.org/10.1136/oem.57.8.550>.
- Sterud T. Work-related psychosocial and mechanical risk factors for work disability: a 3-year follow-up study of the general working population in Norway. *Scand J Work Environ Health.* 2013;39(5):468–76. <https://doi.org/10.5271/sjweh.3359>.
- Guilley E and Lalive d'Epinau CJ. Social status and mortality with activity of daily living disability in later life. *J Gerontol B Psychol Sci Soc Sci.* 2008; 63(3):S192–6. <https://doi.org/10.1093/geronb/63.3.S192>.
- Groffen DAI, Koster A, Bosma H, van den Akker M, Aspelund T, Siggeirsdóttir K, et al. Socioeconomic factors from midlife predict mobility limitation and depressed mood three decades later; Findings from the AGES-Reykjavik Study. *BMC Public Health.* 2013;13:101–6. <https://doi.org/10.1186/1471-2458-13-101>.
- Idland G, Pettersen R, Avlund K, Bergland A. Physical performance as long-term predictor of onset of activities of daily living (ADL) disability: A 9 year longitudinal study among community-dwelling older women. *Archives of Gerontology and Geriatrics.* 2013;56(3):501–6. <https://doi.org/10.1016/j.archger.2012.12.005>.
- Hinrichs T, von Bonsdorff MB, Törmäkangas T, von Bonsdorff ME, Kulmala J, Seitsamo J, et al. Inverse effects of Midlife Occupational and leisure time physical activity on mobility limitation in old age- A 28-Year prospective follow up study. *J Am Geriatr Soc.* 2014;62(5):812–20. <https://doi.org/10.1111/jgs.12793>.
- Ropponen A, Samuelsson Å, Alexanderson K, Svedberg P. Register- based data of psychosocial working conditions and occupational groups as predictors of disability pension due to musculoskeletal diagnoses: a prospective cohort study of 24543 Swedish twins. *BMC Musculoskelet Disord.* 2013;14:268. <https://doi.org/10.1186/1471-2474-14-268>.
- Messing K, Stock SR, Tissot F. Should studies of risk factors for musculoskeletal disorders be stratified by gender? Lessons from the 1998 Québec health and social survey. *Scand J Work Environ Health.* 2009; 35(2):96–112. <https://doi.org/10.5271/sjweh.1310>.
- Sabbath EL, Glymour MM, Descatha A, Leclerc A, Zins M, Goldberg M, et al. Biomechanical and psychosocial occupational exposures: Joint predictors of post-retirement functional health in the French GAZEL cohort. *Advances in Life Course Research.* 2013;18:235–43. <https://doi.org/10.1016/j.alcr.2013.07.002>.
- Ilmarinen J, Suurnakki T, Nygård C-H, Landau K. Classification of municipal occupations. *Scand J Work Environ Health.* 1991;17(suppl 1):12–29.
- von Bonsdorff MB, Seitsamo J, Ilmarinen J, Nygård CH, von Bonsdorff ME, Rantanen T. Work ability in midlife as a predictor of mortality and disability in later life: a 28-year prospective follow-up study. *CMAJ.* 2011;183(4):E235–42. <https://doi.org/10.1503/cmaj.100713>.
- von Bonsdorff MB, Seitsamo J, von Bonsdorff ME, Ilmarinen J, Nygård CH, Rantanen T. Job strain among blue-collar and white-collar employees as a determinant of total mortality: a 28-year population based follow-up. *BMJ Open.* 2012;2(2):e000860. <https://doi.org/10.1136/bmjopen-2012-000860>.
- Spector WD, Fleishman JA. Combining activities of daily living with instrumental activities of daily living to measure functional disability. *J Gerontol B Psychol Sci Soc Sci.* 1998;53B(1):S46–57. <https://doi.org/10.1093/geronb/53B.1.S46>.
- Jagger C, Matthews R, Matthews F, Robinson T, Robine J-M, Brayne C. The burden of disease on disability-free life expectancy in later life. *J Gerontol A Biol Sci Med Sci.* 2007;62(4):408–14. <https://doi.org/10.1093/gerona/62.4.408>.
- Kulmala J, von Bonsdorff MB, Stenholm S, Törmäkangas T, von Bonsdorff ME, Nygård C-H, et al. Perceived stress symptoms in midlife predict disability in old age: a 28-year prospective cohort study. *J Gerontol A Biol Sci Med Sci.* 2013;68(8):984–91. <https://doi.org/10.1093/gerona/gls339>.
- Katz S. Assessing self-maintenance: activities of daily living, mobility, and instrumental activities of daily

- living. *J Am Geriatr Soc.* 1983;31:721–7. <https://doi.org/10.1111/j.1532-5415.1983.tb03391.x>.
22. KCP, Neupane S, Leion-Arjas P, von Bonsdorff MB, Rantanen T, von Bonsdorff ME, et al. Work-related biomechanical exposure and job strain as separate and joint predictors of musculoskeletal diseases: A 28-year prospective follow-up study. *Am J Epidemiol* kwx189. <https://doi.org/10.1093/aje/kwx189>.
 23. Karasek RA. Job demands, job decision latitude, and mental strain: implications for job redesign. *Admin Sci Q.* 1979;24(2):285–308. <https://doi.org/10.2307/2392498>.
 24. van der Weele TJ, Knol MJ. A tutorial on Interaction. *Epidemiol. Methods.* 2014;3(1):33–72.
 25. Warhrendorf M, Sembajwe G, Zins M, Berkman L, Goldberg M, Siegrist J. Long-term effects of psychosocial work stress in midlife on health functioning after labor market exit – results from the GAZEL study. *J Gerontol B Psychol Sci Soc Sci.* 2012;67(4): 471–80. <https://doi.org/10.1093/geronb/gbs045>.
 26. McCarthy VJC, Perry IJ, Greiner BA. Has your work worked you too hard? Physically demanding work and disability in a sample of the older Irish population. *Ir J Med Sci.* 2013;182:47–55. <https://doi.org/10.1007/s11845-012-0824-7>.
 27. Geuskens GA, Burdorf A, Barendregt PJ, Hazes JMW. A high physical workload and high job demands hamper the good prognosis in physical functioning in persons with early inflammatory joint conditions. *Rheumatology.* 2011;50:789–98. <https://doi.org/10.1093/rheumatology/keq359>.
 28. Rijs KJ, van der pas S, Geuskens GA, Cozijnsen R, Koppes LLJ, van der Beek AJ, et al. Development and validation of physical and psychosocial job-exposure matrix in older and retired workers. *Ann Occup Hyg.* 2013;58(2):152–70.
 29. Støver M, Pape K, Johnsen R, Fleten N, Sund ER, Ose So, et al. Work environment and disability pension- an 18 year follow-up study in a Norwegian working population. *Scand J Public Health.* 2013;41(6):587–96. <https://doi.org/10.1177/1403494813486965>.
 30. Grooten WJA, Mulder M, Josephson M, Alfredsson L, Wiktorin C. The influence of work-related exposures on the prognosis of neck/shoulder pain. *Eur Spine J.* 2007;16(2):2083–91. <https://doi.org/10.1007/s00586-007-0481-0>.
 31. Canivet C, Choi B, Karasek R, Moghaddassi M, Staland-Nyman C, Östergren P. Can high psychological job demands, low decision latitude, and high job strain predict disability pensions? A 12-year follow-up of middle-aged Swedish workers. *Int Arch Occup Environ Health.* 2013;86(3):307–19. <https://doi.org/10.1007/s00420-012-0766-4>.
 32. Benavides FG, Duran X, Gimeno D, Vanroelen, Martinez JM. Labour market trajectories and early retirement due to permanent disability: a study based on 14972 new cases in Spain. *European Journal of Public Health.* 2015;25(4):673–7. <https://doi.org/10.1093/eurpub/cku204>.
 33. Clausen T, Burr H, Borg V. Does affective organizational commitment and experience of meaning at work predict risk of disability pensioning? An analysis of register based outcomes using pooled data on 40,554 observations in four occupational group. *Am J Ind Med.* 2014;57(6):709–17. <https://doi.org/10.1002/ajim.22313>.
 34. Dembe AE, Yao X, Wickizer TM, Shoben AB, Dong X. Using O*NET to estimate the association between work exposures and chronic diseases. *Am J Ind Med.* 2014;57(9):1022–31. <https://doi.org/10.1002/ajim.22342>.
 35. Rydwik E, Welmer A-K, Angleman S, Fratiglioni L, Wang H-X. Is Midlife Occupational Physical Activity Related to Disability in Old Age? The SNAC-Kungsholmen Study. *PLoS ONE.* 2013;8(7):e70471. <https://doi.org/10.1371/journal.pone.0070471>.
 36. Arvidsson I, Gremark SJ, Dahlqvist, Axmon A, Karlson Björn, Björk J, et al. Cross-sectional associations between occupational factors and musculoskeletal pain in women teachers, nurses and sonographers. *BMC Musculoskelet Disord.* 2016;17:35. <https://doi.org/10.1186/s12891-016-0883-4>.
 37. Järholm B, Stattin M, Robroek SJ, Janlert U, Karlsson B, Burdorf A. Heavy work and disability pension – a long term follow-up of Swedish construction workers. *Scand J Work Environ Health.* 2014;40(4):335–42. <https://doi.org/10.5271/sjweh.3413>.
 38. Vingård E, Alfredsson L, Hagberg M, Kilbom A, Theorell T, Waldenström M, et al. To what extent do current and past physical and psychosocial occupational factors explain care-seeking for low back pain in a working population? Results from the musculoskeletal intervention Norrälje study. *Spine (Phila Pa 1976).* 2000;25(4):493–500. <https://doi.org/10.1097/00007632-200002150-00017>.
 39. Devereux JJ, Vlachonikolis IG, Buckle PW. Epidemiological study to investigate potential interaction between physical and psychosocial factors at work that may increase the risk of symptoms of musculoskeletal disorder of the neck and upper limb. *Occup Environ Med.* 2002;59(4):269–77. <https://doi.org/10.1136/oem.59.4.269>.
 40. Huuhtanen P, Nygård C-H, Tuomi K, Martikainen R. Changes in stress symptoms and their relationship to changes at work in 1981–1992 among elderly workers in municipal occupations. *Scand J Work Environ Health.* 1997;23(Suppl 1):36–48.
 41. Nygård C-H, Huuhtanen P, Tuomi K, Martikainen R. Perceived work changes between 1981 and 1992 among aging workers in Finland. *Scand J Work Environ Health.* 1997;23(suppl 1):12–19.

Received for publication: 13 February 2017