Scand J Work Environ Health 2017;43(6):560-568

<https://doi.org/10.5271/sjweh.3666>

Published online: 07 Sep 2017, Issue date: 01 Nov 2017

Shift work and the incidence of prostate cancer: a 10-year follow-up of a German population-based cohort study

by Behrens T, Rabstein S, Wichert K, Erbel R, Eisele L, Arendt M, Dragano N, Brüning T, Jöckel K-H

Affiliation: Institute for Prevention and Occupational Medicine of the German Social Accident Insurance (IPA), Institute of the Ruhr-Universität Bochum (IPA), Bürkle-de-la-Camp-Platz 1, 44789 Bochum, Germany. behrens@ipa-dguv.de

Refers to the following texts of the Journal: 2007;33(5):336-343
2008;34(1):5-22 2010;36(2):163-179 2014;40(5):502-510

Key terms: cancer; chronotype; circadian disruption; cohort study; Germany; Heinz Nixdorf Recall Study; night work; population-based cohort study; prostate cancer; shift work; sleep preference; vitamin D

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/28879368

Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](#).

This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

Shift work and the incidence of prostate cancer: a 10-year follow-up of a German population-based cohort study

by Thomas Behrens, MD, MPH,¹ Sylvia Rabstein, PhD,¹ Katharina Wichert, MSc,¹ Raimund Erbel, MD,² Lewin Eisele, MD,² Marina Arendt, MSc,² Nico Dragano, PhD,³ Thomas Brüning, MD,¹ Karl-Heinz Jöckel, PhD²

Behrens T, Rabstein S, Wichert K, Erbel R, Eisele L, Arendt M, Dragano N, Brüning T, Jöckel K-H. Shift work and the incidence of prostate cancer: a 10-year follow-up of a German population-based cohort study. *Scand J Work Environ Health*. 2017;43(6):560–568. doi:10.5271/sjweh.3666

Objectives We investigated the association of shift and night work with the incidence of prostate cancer using data of the population-based prospective Heinz Nixdorf Recall Study from the highly industrialized Ruhr area in Germany.

Methods Participants of the baseline survey were recruited between 2000–2003. A follow-up survey including, a detailed interview on shift and night work, was conducted from 2011–2014. We included 1757 men who did not report a history of prostate cancer at baseline. We assessed shift- and night-work exposure up to time of the baseline interview. Incident prostate cancers were recorded from baseline through September 2014. We calculated hazard ratios (HR) of shift- and night-work exposure using Cox proportional hazards regression with age at event as timescale, adjusting for smoking status, family history of prostate cancer, education (≤ 13 , 14–17, ≥ 18 years), and equivalent income (low, medium, high).

Results We observed a twofold increased HR for prostate cancer among shift and night workers. Ever employment in shift work was associated with HR 2.29, 95% confidence interval (CI) 1.43–3.67 and night work with HR 2.27, 95% CI 1.42–3.64. HR increased steadily with duration of employment in shift or night work. Stratifying analyses by preferred midpoint of sleep, yielded strongly elevated HR among subjects with early sleep preference, although these analyses were limited by small number of cases.

Conclusions We identified increased risks for prostate cancer among men with employment in shift or night work. HR were strongly elevated among long-term employed shift workers and men with early preferred midpoint of sleep.

Key terms chronotype; circadian disruption; Germany; Heinz Nixdorf Recall Study; night work; sleep preference; vitamin D.

In 2007, the International Agency for Research on Cancer (IARC) concluded that shift work involving circadian disruption was probably carcinogenic in humans based upon evidence of an increased breast cancer risk among women. The IARC expert panel judged that there was "sufficient evidence" in experimental animals but "limited evidence" in humans (Group 2A) (1).

Possible mechanisms linking shift work and cancer center around the "light-at-night" (LAN) hypothesis, which postulates that exposure to artificial LAN sup-

presses pineal melatonin secretion and subsequently leads to an increase of sex hormones, which in turn could increase the susceptibility to hormone-dependent cancers (2). Also, anti-proliferative effects of melatonin that could influence tumor origin and progression have been demonstrated *in vivo* and *in vitro* for prostate cancer cells (3).

Although much of the research regarding the carcinogenicity of circadian disruption focused on breast cancer (4, 5), an increasing body of evidence has emerged associating prostate cancer with shift work. A meta-

¹ Institute for Prevention and Occupational Medicine of the German Social Accident Insurance (IPA), Institute of the Ruhr-Universität Bochum, Bochum, Germany.

² Institute for Medical Informatics, Biometry and Epidemiology, University Hospital of Essen, University Duisburg-Essen, Essen, Germany.

³ Institute for Medical Sociology, Centre for Health and Society, Faculty of Medicine, Heinrich Heine University, Düsseldorf, Germany.

Correspondence to: Thomas Behrens, Institute for Prevention and Occupational Medicine of the German Social Accident Insurance (IPA), Institute of the Ruhr-Universität Bochum (IPA), Bürkle-de-la-Camp-Platz 1, 44789 Bochum, Germany. [E-mail: behrens@ipa-dguv.de]

analysis of earlier research judged that the overall evidence was suggestive for a positive association between proxies of circadian disruption and prostate cancer risk (6). However, results from subsequent reports remained inconsistent (7–14).

Chronotype or diurnal preference may influence an individual's sensitivity to circadian disruption (with "evening types" showing better adaptation to shift work) which in turn may affect cancer risks (15). However, previous study results on prostate cancer were not consistent with this hypothesis showing higher risks rather in evening types (13) or similarly increased risks for men with morning and evening, but not intermediate type (11). In addition, methods to assess diurnal preference differed between studies, which limits the direct comparability of results.

Therefore, we investigated the association of shift and night work with the incidence of prostate cancer using data of the population-based Heinz Nixdorf Recall Cohort Study (HNR) from the Ruhr area in Germany. In contrast to industry-based cohorts, we studied a variety of jobs associated with shift and night work, following recommendations by an IARC Working Group how to define shift work in epidemiological studies (16). We considered various non-occupational confounders that may affect the incidence of prostate cancer. In addition, we studied the differential effect of diurnal preference, assessed by preferred midpoint of sleep, on prostate cancer risk.

Methods

Study population

The detailed methods of the HNR study have been described elsewhere (17). Briefly, the HNR is a population-based cohort study involving a random sample of 45–74 year-old inhabitants of the highly industrialized Ruhr area in Germany. Participants for the baseline survey were recruited from 2000–2003 and a follow-up survey was conducted from 2011–2014.

Of the 4100 men eligible to participate in the study, 2395 participated in the baseline survey. We excluded 39 participants who were diagnosed with prostate cancer before baseline. Of 2356 men without a history of prostate cancer at baseline, 1481 participated in the follow-up investigation. For 1438 men shift-work information was available from the follow-up interview. For an additional 319 men with insufficient shift-work information from the follow-up interview, shift-work exposure was complemented from the baseline interview. Therefore, 1757 men with information on shift and night work could be analyzed (figure 1).

All incident prostate tumors were monitored from

Figure 1. Formation of the analysis cohort

baseline through September 2014. An independent endpoint committee of the HNR study evaluated patient records and, in the case of deceased participants, death certificates to assess the validity of the incident endpoints. Agreement of questionnaire-based incident cancer cases with medical records was evaluated for an 8-year follow-up and was considered to be satisfactory (18). The Ethics Committee of the University of Duisburg-Essen approved the study, and all subjects provided written informed consent.

Prostate cancer risk was higher among men who were not included in the analysis cohort (analysis cohort: $414.8 / 10^5$ person-years versus $855.1 / 10^5$ person-years). In addition, non-participants tended to be older (mean age 65.5 versus 57.5 years), had lower educational levels (66.1% versus 48% with education ≤ 13 years), and lower income (52.4 versus 33% in the lowest category).

Exposure assessment

For the follow-up interview, we employed a supplemental shift-work module, which included a detailed phase-by-phase assessment of the shift-work history

and specific exposure characteristics, following recommendations by an IARC working group (16). Shift work was defined as any regular employment in shift systems including work hours outside 07:00–18:00 hours, whereas night work was defined as a shift that included work between 24:00–05:00 hours.

For subjects not participating in the follow-up, we used two questions from the baseline survey soliciting employment in shift work from all subjects who had been gainfully employed or unemployed for less than two years: (1) "Have you ever been employed in shift work?" (never; yes, not involving night work; yes, involving night work), and (2) "How long have you been involved in shift work?"

We included the following shift work variables in the analysis: 1) ever shift work (≥ 1 year) up to the baseline interview, 2) ever night work (≥ 1 year) up to baseline, 3) duration of shift work, and 4) duration of night work up to baseline, respectively. We quantified total duration of shift/night as follows: 0–<1, 1–<10, 10–<20 years, ≥ 20 years. To calculate duration of shift-work exposure, shift-work information from the follow-up interview was censored at baseline, ie, job periods involving shift work after baseline were not considered in the analysis. Information from the baseline interview up to the time of baseline was used for 319 men (figure 1).

Subjects were analyzed as having ever been employed in shift or night work during their work life. We also analyzed the association of employment in shift work without night work ever, although we observed this pattern in only 22% of shift workers,

To assess risk estimates for particular jobs associated with shift or night work, we studied subjects' full job history. We analyzed the subject's longest-held job (coded according to the "International Standard Classification of Occupations, revision 1968 (ISCO-68) that involved shift or night work according to the shift-work module. Due to the great variety of jobs in this population-based cohort (and the small number of single job titles), we present hazard ratios (HR) only for the three most frequent longest-held jobs in this cohort, as well as an analysis pooling jobs according to industrial sector: industries (eg, jobs in the metal, chemical, construction, electrical, and mining industry etc.) and public services (such as police officers, administrative and public transport employees).

To estimate a subject's diurnal preference, we employed two questions from the follow-up interview soliciting the preferred bed- and wake-up time during days off work. From these two time specifications, we calculated the preferred midpoint of sleep. To define early, intermediate, and late midpoint of sleep, we determined age-specific tertiles separately in age groups 55–<60, 60–<65, 65–<70, 70–<75, 75–<80, and ≥ 80

years. Cut-offs at the 25% and 75% percentile were used to define early, intermediate, and late preferred sleep midpoint, resulting in age-specific sleep midpoints <02:30–03:00 hours for the early, 02:30–04:00 hours for the intermediate, and >03:30–04:00 hours for the late preference, respectively.

Total serum 25(OH)D at baseline was measured in 2007 on thawed samples using Liaison assay (DiaSorin). Vitamin D status was categorized as "low" or "high" based on a cut-off at the median concentration (15.3 ng/ml).

Statistical analysis

We performed Cox proportional hazards regression analysis with age at event as timescale to determine the association between shift work before baseline and prostate cancer incidence. We considered the age-at-event model as more suitable than the time-to-event model with linear age adjustment, because prostate cancer is a tumor of the elderly, and one would rather expect a difference between older and younger men than two subjects of the same age with different follow-up times. For the purpose of sensitivity analysis, we also calculated the Cox model with time to event as timescale and adjusting for age at baseline.

As potential confounders, we considered smoking status (never, former, current smoker), body mass index (<20, 20–<25, 25–<30, ≥ 30 kg/m²), physical activity [metabolic equivalent task (MET)-hours, assessed by an interview question regarding sports/physical activities during the last four weeks, combined with additional questions directed at duration and frequency of these activities, linear], alcohol consumption (g alcohol/week, linear), family history of prostate cancer, school education (≤ 13 , 14–17 (reference), ≥ 18 years) adapted to the International Standard Classification of Education (ISCED), version 1997 (19), and equivalent income according to tertiles of income among men without prostate cancer [in German marks (DM)] [<2567.60; 2567.60–<3666.70 (reference); 3 666.70+ DM]. Next to family history of prostate cancer, potential confounders were included if risk estimates changed by >10% or if the P-value of the factor was <0.15.

We present three models: model 1, with age at event as timescale, an adjusted model including smoking status and family history of prostate cancer, and the fully adjusted model adding indicators of socioeconomic status (SES). Analyses were stratified by preferred midpoint of sleep and vitamin D-status. Tests for trend were calculated continuously.

Furthermore, we ran a sensitivity analysis restricted to men who had answered the supplemental shift-work module in the follow-up survey.

Results

We observed 76 incident primary prostate cancer cases during follow-up. The mean age at prostate cancer diagnosis was 66.8 years [standard deviation (SD) 6.83]. Among men without prostate cancer mean age was 77.3 years (SD 7.18) at the time of interview. At follow-up, 106 men without prostate cancer were ≥ 80 years (age at interview), whereas only one subject was ≥ 80 years at age of prostate cancer diagnosis. 34% of men reported a history of shift work, whereas a history of night work was indicated by 27%. The preferred midpoint of sleep was late for 14.1% and early for 13% of subjects. For further descriptive results see table 1.

Associations between shift work and prostate cancer incidence are reported in table 2. In the fully adjusted model including SES indicators, we observed increased HR among shift workers (HR 2.29, 95% CI 1.43–3.67) and night workers (HR 2.27, 95% CI 1.42–3.64). Being employed for ≥ 20 years in shift or night work was associated with three- to fourfold increased risks (HR 3.08 95% CI 1.67–5.69 and HR 3.76, 95% CI 2.04–6.93, respectively). The analysis of men employed in shift work never entailing night work showed slightly smaller HR. Because most exposed men (78%) had exposure to both shift and night work, the potential to assess their independent effects was limited and based on only six men with prostate cancer (table 2). Including body mass index, physical activity, and alcohol consumption as confounders changed the overall results only slightly (not shown).

The most frequent jobs associated with shift-work history were in mining [N=45 (7.5%), prostate cancer cases N=5], engine building/production engineering [N=41 (6.8%), prostate cancer cases N=3], and metal working (eg, metal-cutting, metal-forming, foundries) [N=38 (6.3%), prostate cancer cases N=5], which were associated with HR of 2.34 (95% CI 0.92–5.96), HR 1.47 (95% CI 0.46–4.71), and HR 3.68 (95% CI 1.46–9.27), respectively (fully adjusted model). Shift work in industries was associated with an HR 2.45 (95% CI 1.46–4.09) based on 22 prostate cancer cases. Public service jobs involving shift work showed HR 2.26 (95% CI 0.90–5.63) based on five men with prostate cancer who were employed in these jobs (supplemental table S4, www.sjweh.fi/index.php?page=data-repository).

Stratification by preferred midpoint of sleep revealed a trend with shift- and night-work duration particularly for the early sleep preference. We observed smaller, but still increased HR for intermediate and late sleep preferences (table 3).

No differences in prostate cancer risk due to shift or night work were seen in the analysis stratified by vitamin D level (table 4).

Analyzing the associations with time to event as timescale did not change the results substantially (sup-

Table 1. Descriptive data of the analysis cohort, stratified by shift work status at baseline, 1757 men. [DM=Deutchmarks]

	Total population		Prostate cancer		Never shift work		Ever shift work	
	N	%	N	%	N	%	N	%
Total	1757	100	76	100	1156	100	601	100
Shift work ever								
Never/less <1 year	1156	65.8	38	50.0	1156	100		
Yes	601	34.2	38	50.0			601	100
Duration of shift work (years) ^a								
0–<1	1156	65.8	38	50.0	1156	100		
1–<10	246	14.0	13	17.1			246	40.9
10–<20	127	7.2	8	10.5			127	21.1
≥ 20	219	12.5	17	22.4			219	36.4
Missing	9	0.5					9	1.5
Night work ever								
0–<1 year	1285	73.1	44	57.9	1156	100	129	21.5
Yes	472	26.9	32	42.1			472	78.5
Duration of night work (years) ^a								
0–<1	1285	73.5	44	57.9	1156	100	129	21.5
1–<10	204	11.7	11	14.5			204	33.9
10–<20	99	5.7	5	6.6			99	16.5
≥ 20	161	9.2	16	21.1			161	26.8
Missing	8	0.5					8	1.3
Age (years)								
45–<55	700	39.8	16	21.1	453	39.2	247	41.1
55–<65	734	41.8	31	40.8	490	42.4	244	40.6
65–75	323	18.4	29	38.2	213	18.4	110	18.3
Preferred midpoint of sleep ^a								
Early	228	13.0	13	17.1	147	12.7	81	13.5
Intermediate	909	51.7	42	55.3	619	53.6	290	48.3
Late	248	14.1	8	10.5	166	14.4	82	13.6
Missing	372	21.2	13	17.1	224	19.4	148	24.6
Family history of prostate cancer								
Yes	23	1.3	2	2.6	13	1.1	10	1.7
No	1734	98.7	74	97.4	1143	98.9	591	98.3
Smoking status								
Never	507	28.9	33	43.4	374	32.4	133	22.1
Former	809	46.0	34	44.7	506	43.8	303	50.4
Current	441	25.1	9	11.8	276	23.9	165	27.5
Vitamin D status								
Low	731	41.6	26	34.2	491	42.5	240	39.9
High	747	42.5	33	43.4	480	41.5	267	44.4
Missing	279	15.9	17	22.4	185	16.0	94	15.6
Education (years)								
≤ 13	843	48.0	32	42.1	470	40.7	373	62.1
14–17	622	35.4	34	44.7	430	37.2	192	32.0
≥ 18	290	16.5	10	13.2	255	22.1	35	5.8
Missing	2	0.1			1	0.1	1	0.2
Income (DM)								
<2567.60	579	33.0	19	25.0	313	27.1	266	44.3
2567.60–<3666.70	568	32.3	32	42.1	366	31.7	202	33.6
≥ 3666.70	553	31.5	23	30.3	440	38.1	113	18.8
Missing	57	3.2	2	2.6	37	3.2	20	3.3

^a Assessed at 2011–2014 follow-up.

plemental tables S1 and S2). When we restricted the analysis to men participating in the follow-up survey, we observed HR that were similar as compared to the full analysis cohort (results not shown).

Table 2. Associations between shift and night work in the Heinz Nixdorf Recall Cohort [HR=hazard ratio; 95% CI=95% confidence interval]

Factor	Person-years	Prostate cancer (N)	HR ^a	95% CI	P for trend	HR ^b	95% CI	P for trend	HR ^c	95% CI	P for trend
0<1 year of shift work	12 168	38	1.0	Reference		1.0	Reference		1.0	Reference	
Ever ≥1 years of shift work	6154	38	2.0	1.28–3.14		2.15	1.36–3.38		2.29	1.43–3.67	
1<10 years of shift work	2500	13	1.72	0.92–3.23	0.0012	1.82	0.97–3.42	0.0004	1.87	0.99–3.55	0.0001
10<20 years of shift work	1312	8	2.03	0.95–4.35		2.05	0.96–4.41		2.18	1.01–4.72	
≥20 years of shift work	2245	17	2.38	1.34–4.21		2.66	1.49–4.77		3.08	1.67–5.69	
0<1 year of shift work	12 168	38	1.0	Reference		1.0	Reference		1.0	Reference	
Ever ≥1 year of shift work without night work during work life	1343	6	1.54	0.65–3.64		1.57	0.66–3.72		1.91	0.79–4.64	
0<1 year of night work	13 511	44	1.0	Reference		1.0	Reference		1.0	Reference	
Ever ≥1 year of night work	4811	32	2.01	1.28–3.18		2.18	1.37–3.45		2.27	1.42–3.64	
1<10 years of night work	2089	11	1.60	0.83–3.11	0.0003	1.70	0.88–3.30	<0.0001	1.72	0.88–3.35	<0.0001
10<20 years of night work	1031	5	1.48	0.59–3.74		1.59	0.63–4.01		1.68	0.66–4.26	
≥20 years of night work	1606	16	2.99	1.68–5.29		3.32	1.86–5.95		3.76	2.04–6.93	

^a Modelled with age at event as timescale only.^b Modelled with age at event and adjusted for smoking (never, former smoker, current smoker) and family history of prostate cancer (yes, no).^c As in b, additionally adjusted for level of school education (≤13, 14–17, ≥18 years) and equivalent income (low, medium, high).**Table 3.** Associations between shift work and prostate cancer according to preferred midpoint of sleep. [HR=hazard ratio; 95% CI=95% confidence interval]

Factor	Person-years	Prostate cancer (N)	HR ^a	95% CI	P for trend	HR ^b	95% CI	P for trend	HR ^c	95% CI	P for trend
Preferred midpoint of sleep ^d											
Early	2475	13	1.18			1.20	0.64–2.23		1.17	0.63–2.18	
Intermediate	9829	42	1.0	Reference		1.0	Reference		1.0	Reference	
Late	2674	8	0.65	0.31–1.39		0.70	0.34–1.50		0.69	0.32–1.48	
Early											
0<1 year of shift work	1616	4	1.0	Reference		1.0	Reference		1.0	Reference	
Ever, ≥1 year of shift work	859	9	4.56	1.40–14.89		4.14	1.21–14.21		5.47	1.45–20.71	
1<10 years of shift work	383	4	4.52	1.12–18.34	0.0155	3.14	0.68–14.60	0.0150	4.09	0.80–20.97	
≥10 years of shift work	476	5	4.59	1.23–17.14		5.05	1.33–19.23		6.94	1.55–31.13	
0<1 year of night work	1865	6	1.0	Reference		1.0	Reference		1.0	Reference	
Ever, ≥1 year of night work	610	7	3.72	1.25–11.10		4.68	1.44–15.21		6.43	1.81–22.80	
1<10 years of night work	239	2	2.60	0.52–13.0	0.0117	2.93	0.54–15.77	0.0047	4.48	0.74–27.11	0.0031
≥10 years of night work	372	5	4.48	1.37–14.73		5.98	1.71–21.0		7.72	1.92–30.97	
Intermediate											
0<1 year of shift work	6752	20	1.0	Reference		1.0	Reference		1.0	Reference	
Ever, ≥1 year of shift work	3078	22	2.23	1.22–4.09		2.36	1.28–4.35		2.37	1.26–4.45	
1<10 years of shift work	108	8	2.25	0.99–5.13	0.0112	2.31	1.01–5.26	0.0071	2.26	0.98–5.19	0.0079
≥10 years of shift work	1914	14	2.29	1.16–4.55		2.45	1.22–4.09		2.52	1.23–5.19	
0<1 year of night work	7467	24	1.0	Reference		1.0	Reference		1.0	Reference	
Ever, ≥1 year of night work	2362	18	2.12	1.15–3.91		2.26	1.21–4.20		2.30	1.22–4.35	
1<10 years of night work	974	8	2.26	1.01–5.05	0.0238	2.33	1.04–5.22	0.0150	2.38	1.06–5.39	0.0151
≥10 years of night work	1344	10	2.09	1.00–4.38		2.26	1.07–4.79		2.31	1.07–4.99	
Late											
0<1 year of shift work	1794	5	1.0	Reference		1.0	Reference		1.0	Reference	
Ever, ≥1 year of shift work	880	3	1.29	0.30–5.44		1.23	0.29–5.20		1.20	0.27–5.29	
0<1 year of night work	1938	5	1.0	Reference		1.0	Reference		1.0	Reference	
Ever, ≥1 year of night work	736	3	1.63	0.39–6.87		1.60	0.38–6.71		1.42	0.33–6.20	

^a Modelled with age at event as timescale only.^b Modelled with age at event and adjusted for smoking (never, former smoker, current smoker) and family history of prostate cancer (yes, no).^c As in b, additionally adjusted for level of school education (≤13, 14–17, ≥18 years) and equivalent income (low, medium, high).^d Preferred midpoint of sleep according to 5-year age groups.

Table 4. Associations between shift work and prostate cancer according to vitamin D status.

Factor	Person-years	Prostate cancer (N)	HR ^a	95% CI	HR ^b	95% CI	HR ^c	95% CI
"Low" vitamin D								
0–<1 year of shift work	5161	13	1.0	Reference	1.0	Reference	1.0	Reference
Ever, ≥1 year of shift work	2453	13	2.12	0.98–4.58	2.21	1.01–4.87	2.42	1.06–5.51
0–<1 year of night work	5663	15	1.0	Reference	1.0	Reference	1.0	Reference
Ever ≥1 year of night work	1951	11	2.03	0.93–4.43	2.08	0.94–4.63	2.22	0.98–5.04
"High" vitamin D								
0–<1 year of shift work	5072	16	1.0	Reference	1.0	Reference	1.0	Reference
Ever, ≥1 year of shift work	2768	17	2.04	1.03–4.05	2.19	1.10–4.37	2.41	1.18–4.92
0–<1 year of night work	5726	18	1.0	Reference	1.0	Reference	1.0	Reference
Ever ≥1 year of night work	2114	15	2.31	1.16–4.59	2.45	1.23–4.90	2.63	1.29–5.35

^a Modelled with age at event as timescale only.

^b Modelled with age at event and adjusted for smoking (never, former smoker, current smoker) and family history of prostate cancer (yes, no).

^c as in b, additionally adjusted for level of school education (≤13, 14–17, ≥18 years) and equivalent income (low, medium, high).

^d Vitamin D status stratified at median at baseline.

Discussion

Using data from the prospective HNR, we identified increased risks for prostate cancer among men who were employed in shift or night work. Prostate cancer risks were particularly pronounced for early sleep preference.

Strengths of this analysis include that we adapted questions from a shift-work module, which was created according to recommendations by an IARC working group and included detailed information to differentiate between shift and night work (16). In contrast to many earlier publications, we were able to consider a large number of potential confounders in our analyses (such as physical activity, family history of prostate cancer, alcohol consumption, and smoking behavior). Furthermore, we were able to test the influence of preferred midpoint of sleep and low vitamin D levels on prostate cancer risk, which are discussed as possible effect modifying factors (16, 20).

Limitations include that the number of incident cases was still small, stratified analyses by duration of shift or night work could not be performed in all subgroups, and the numbers of subjects in subgroups were small and results therefore tended to show wide confidence intervals.

Further, shift work was assessed retrospectively, which may introduce the possibility of reverse causation bias. Reverse causation may occur if prodromal symptoms of prostate cancer affect sleep patterns and subsequently influence the choice of shift worked. We consider the choice of shift work type due to sleep disturbances rather unlikely because the lifelong shift-work history was assessed before baseline, whereas incident cancer was monitored prospectively from the baseline interview. However, we cannot rule out completely that urinary tract symptoms such as nocturia may have modified shift-work exposure as the induction time of prostate cancer is rather lengthy.

It is noteworthy that we observed a much higher

prostate cancer risk among men who were not included in the analysis cohort, mainly due to loss to follow-up. Non-participants tended to be older, had lower educational levels, and lower income. Although age may explain part of these differences, a spurious protective association with low SES indicators would rather overestimate the association between shift work and prostate cancer risk. We therefore analyzed a separate model to assess the influence of age, income, education and smoking on prostate cancer risk in the entire HNR cohort (N=2356 men), which showed a decreased HR for low income (HR 0.76, 95% CI 0.48–1.18) and smoking (former smokers: HR 0.60, 95% CI 0.40–0.90; current smokers: HR 0.60, 95% CI 0.35–1.02), but a small increase for low educational level (HR 1.15, 95% CI 0.77–1.74, results not shown). Excluding income and education from the model (equivalent to the absence of confounding by SES indicators) still yielded estimates which were elevated two- to threefold (see table 2). Although we cannot predict a further attenuating effect on risk estimates if low SES indicators were true risk factors for prostate cancer, it would require very strong confounding to reduce these associations to a null effect.

Furthermore, we were not able to include more detailed information on shift work as recommended by IARC, such as time schedules of different shift systems, number of days off after shifts, or direction of shift rotation. These questions were either insufficiently answered in our questionnaire or hard to classify retrospectively due to the complexity of reported shift systems.

It has been pointed out that even early day shifts may be associated with significant circadian disruption in some workers, posing them at increased cancer risk (21). This possibility was implied by our finding that ever shift work, defined as working outside regular hours between 07:00–18:00 hours, showed similar risk estimates to night work taking place at work hours involving substantial circadian disruption.

To assess circadian disruption among shift workers by current diurnal preference is difficult because preferences may change with age. To estimate individual diurnal preference, we solicited men's preferred midpoint of sleep during leisure days at the 2011 follow-up, which is not an ideal approach (22). Because the subjects' age in the HNR was >55 years in the follow-up survey, we only determined the current age-specific sleep preference and did not assess the subjects' past preferred midpoint of sleep. However, the majority of men in the HNR were already retired, limiting entrainment of the circadian clock by occupational and environmental influences (22).

Early reports on the association between shift work and the risk of prostate cancer found heterogeneous results (23–27), but many of these early studies were hampered by methodological concerns, such as misclassification of shift-work exposure and insufficient confounder control (27, 28). However, more recently several studies have been published which support our findings. A Canadian case-control study reported increased odds ratios (OR) for prostate cancer among men performing night work (OR 2.77, 95% CI 1.96–3.92). Estimates were increased across all categories of cumulative duration of night work, although a clear time-dependent trend was not seen (12). In a Spanish case-control study increased prostate cancer risk estimates after 28 years of night work (OR 1.37, 95% CI 1.05–1.81) were reported, but the authors did not observe clear risk differences between workers performing permanent or rotating night shifts (11). Further, the Age, Gene, Environment Susceptibility (AGES) cohort from Iceland showed a two-fold increased prostate cancer risk for men who were suffering from sleep disruption (10). The recent Older Finnish Twin cohort observed an increased risk of prostate cancer among rotating shift workers who were classified as evening chronotypes. In contrast to our results, morning types rather showed a decrease in prostate cancer risk (13). However, the Finnish cohort used self-assessed chronotype, which is not equivalent to the assessment of preferred sleep midpoint and limits direct comparison of these results.

The US Cancer Prevention cohort II observed moderately increased relative risks (RR) for fatal prostate cancer with fixed afternoon/evening shift systems (RR 1.27, 95% CI 0.97–1.65) in the most recent follow-up (1991–2010), whereas rotating shift systems showed RR 1.11, 95% CI 0.97–1.26. The study identified increased risks with insomnia or short sleep duration only in the first, but not the second follow-up, though (7). However, this study concentrated on fatal prostate cancer and therefore does not allow the consideration of less aggressive cancer forms and their association with proxies of circadian disruption.

A large industrial cohort from the BASF company

in Germany that did not identify a difference in prostate cancer risk between rotating shift and day time workers is in contrast to our findings. Notably, standardized incidence ratios (SIR) in this cohort were increased in both, rotating shift and day workers (SIR 1.44 and 1.51, respectively) (8). An advantage of this study certainly is that shift schedules at BASF are uniformly fast forward-rotating, which is not the general rule in all industries. In contrast to the BASF cohort, men in the HNR were employed in many different jobs. The most frequent jobs were mining, metal work, and production engineering, which are associated with a variety of shift systems.

Several mechanisms have been discussed regarding the association between shift work and prostate cancer. The "LAN" hypothesis postulates an effect on hormone levels due to circadian disruption and subsequently an influence on sex-hormone levels and hormone-dependent cancers. Much of the research with respect to circadian disruption so far focused on female breast cancer, whereas relatively little research investigated possible mechanisms in male cancers. In support of the LAN-hypothesis, a case-control study nested in the Icelandic AGES cohort found that low first morning void urinary 6-sulfatoxymelatonin levels were associated with sleep problems as well as the risk for prostate cancer (9). Similarly, a cross-sectional survey from the US National Health and Nutrition Survey (NHANES) observed an association between increased prostate-specific antigen (PSA) levels and night or rotating shift schedule, which may indicate an increased risk for future and aggressive prostate cancer (30).

Low vitamin D levels are discussed as possible effect modifying or mediating factor on cancer risk (16, 20). A recent survey from Italy identified lower vitamin D levels among 96 shift workers as compared to 100 workers engaging in regular daily shifts (31). We studied the influence of vitamin D-deficiency on the association between shift/night work and prostate cancer but found that low vitamin D-levels at baseline had no effect modifying effect.

In our data, we observed confounding particularly for low educational levels and low income which both exerted protective effects, thereby increasing the association between shift work and prostate cancer in the fully adjusted model. This rather surprising finding may be due to the fact that subjects of low SES were less likely to participate in the HNR follow-up (see above). Alternatively, we also may have over-ascertained incident prostate cancer among men with high SES (32). In line with this hypothesis, testing for PSA has been demonstrated to be more frequently performed among more affluent men (33, 34), which is important in countries like Germany where PSA testing is generally not covered by the statutory health insurance. Unfortunately, we were not able to investigate further whether lower

grade, clinically insignificant cancer was more prevalent among the group of men of higher SES because prostate tumor stage was not solicited in the HNR.

In summary, we identified increased risks for prostate cancer among men engaged in shift or night work. We observed the strongest risks for men with early preferred midpoint of sleep. Smoking behavior and proxies of SES strongly influenced the association between shift/night work and prostate cancer, whereas other lifestyle factors were no important confounders.

Acknowledgements

We would like to thank the investigative group and the entire study staff of the Heinz Nixdorf Recall Study. We also thank Evelyn Heinze at IPA for data management and performance of descriptive analyses.

The study's follow-up was funded by the German Social Accident Insurance (DGUV) (grant number FP 295). We thank the Heinz Nixdorf Foundation (Germany) for the generous support of the Heinz Nixdorf Study. This study was also supported by the German Ministry of Education and Science (BMBF) and the Kulturstiftung Essen. The German Research Foundation supported the study (DFG project: ER 155/6-2) and funded the study of psychosocial factors (DFG project SI 236/8-1 and SI 236/9-1).

The authors do not declare any conflict of interest.

TB, SR, KW, and TBr, as staff of the Institute for Prevention and Occupational Medicine (IPA), are employed at the "Berufsgenossenschaft Rohstoffe und chemische Industrie" (BG RCI), a public body, which is a member of the study's main sponsor, the German Social Accident Insurance. IPA is an independent research institute of the Ruhr-Universität Bochum. The authors are independent from the German Social Accident Insurance in study design, access to the collected data, responsibility for data analysis and interpretation, and the right to publish. The views expressed in this paper are those of the authors and not necessarily those of the sponsor.

References

1. Straif K, Baan R, Grosse Y, Secretan B, El Ghissassi F, Bouvard V, et al. Carcinogenicity of shift-work, painting, and fire-fighting. *Lancet Oncology*. 2007;8(12):1065–6. [https://doi.org/10.1016/S1470-2045\(07\)70373-X](https://doi.org/10.1016/S1470-2045(07)70373-X).
2. Stevens RG. Testing the light-at-night (LAN) theory for breast cancer causation. *Chronobiol Int*. 2011;28(8):653–6. <https://doi.org/10.3109/07420528.2011.606945>.
3. Jung-Hynes B, Schmit TL, Reagan-Shaw SR, Siddiqui IA, Mukhtar H, Ahmad N. Melatonin, a novel Sirt1 inhibitor, imparts antiproliferative effects against prostate cancer in vitro in culture and in vivo in TRAMP model. *J Pineal Res*. 2011;50(2):140–9.
4. Wang F, Yeung KL, Chan WC, Kwok CC, Leung SL, Wu C, et al. A meta-analysis on dose-response relationship between night shift work and the risk of breast cancer. *Ann Oncol*. 2013;24(11):2724–32. <https://doi.org/10.1093/annonc/mdt283>.
5. He C, Anand ST, Ebell MH, Vena JE, Robb SW. Circadian disrupting exposures and breast cancer risk: a meta-analysis. *Int Arch Occup Environ Health*. 2015;88(5):533–47. <https://doi.org/10.1007/s00420-014-0986-x>.
6. Sigurdardottir LG, Valdimarsdottir UA, Fall K, Rider JR, Lockley SW, Schernhammer E, et al. Circadian disruption, sleep loss, and prostate cancer risk: a systematic review of epidemiologic studies. *Cancer Epidemiol. Biomarkers Prev*. 2012;21(7):1002–11. <https://doi.org/10.1158/1055-9965.EPI-12-0116>.
7. Gapstur SM, Diver WR, Stevens VL, Carter BD, Teras LR, Jacobs EJ. Work schedule, sleep duration, insomnia, and risk of fatal prostate cancer. *Am J Prev Med*. 2014;46(3 Suppl 1):33. <https://doi.org/10.1016/j.amepre.2013.10.033>.
8. Yong M, Blettner M, Emrich K, Nasterlack M, Oberlinner C, Hammer GP. A retrospective cohort study of shift work and risk of incident cancer among German male chemical workers. *Scand J Work Environ Health*. 2014;40(5):502–10. <https://doi.org/10.5271/sjweh.3438>.
9. Sigurdardottir LG, Markt SC, Rider JR, Haneuse S, Fall K, Schernhammer ES, et al. Urinary melatonin levels, sleep disruption, and risk of prostate cancer in elderly men. *Eur Urol*. 2015;67(2):191–4. <https://doi.org/10.1016/j.eururo.2014.07.008>.
10. Sigurdardottir LG, Valdimarsdottir UA, Mucci LA, Fall K, Rider JR, Schernhammer E, et al. Sleep disruption among older men and risk of prostate cancer. *Cancer Epidemiol Biomarkers Prev*. 2013;22(5):872–9. <https://doi.org/10.1158/1055-9965.EPI-12-1227-T>.
11. Papantoniou K, Casta-o-Vinyals G, Espinosa A, Aragonés N, Pérez-Gómez B, Burgos J, et al. Night shift work, chronotype and prostate cancer risk in the MCC-Spain case-control study. *Int J Cancer*. 2015;137(5):1147–57. <https://doi.org/10.1002/ijc.29400>.
12. Parent M-É, El-Zein M, Rousseau M-C, Pintos J, Siemiatycki J. Night work and the risk of cancer among men. *Am J Epidemiol*. 2012;176(9):751–9. <https://doi.org/10.1093/aje/kws318>.
13. Dickerman BA, Markt SC, Koskenvuo M, Hublin C, Pukkala E, Mucci LA, et al. Sleep disruption, chronotype, shift work, and prostate cancer risk and mortality: A 30-year prospective cohort study of Finnish twins. *Cancer Causes Control*. 2016;27(11):1361–70. <https://doi.org/10.1007/s10552-016-0815-5>.
14. Markt SC, Flynn-Evans EE, Valdimarsdottir UA, Sigurdardottir LG, Tamimi RM, Batista JL, et al. Sleep Duration and

- Disruption and Prostate Cancer Risk: A 23-Year Prospective Study. *Cancer Epidemiol Biomarkers Prev.* 2016;25(2):302–8. <https://doi.org/10.1158/1055-9965.EPI-14-1274>.
15. Saksvik IB, Bjorvatn B, Hetland H, Sandal GM, Pallesen S. Individual differences in tolerance to shift work—a systematic review. *Sleep Med Rev.* 2011;15(4):221–35. <https://doi.org/10.1016/j.smrv.2010.07.002>.
 16. Stevens RG, Hansen J, Costa G, Haus E, Kauppinen T, Aronson KJ, et al. Considerations of circadian impact for defining ‘shift work’ in cancer studies: IARC Working Group Report. *Occup Environ Med.* 2011;68(2):154–62. <https://doi.org/10.1136/oem.2009.053512>.
 17. Schermund A, Möhlenkamp S, Stang A, Grönemeyer D, Seibel R, Hirche H, et al. Assessment of clinically silent atherosclerotic disease and established and novel risk factors for predicting myocardial infarction and cardiac death in healthy middle-aged subjects: rationale and design of the Heinz Nixdorf RECALL Study. Risk Factors, Evaluation of Coronary Calcium and Lifestyle. *Am Heart J.* 2002;144(2):212–8. <https://doi.org/10.1067/mhj.2002.123579>.
 18. Bokhof B, Eisele L, Erbel R, Moebus S. Agreement between different survey instruments to assess incident and prevalent tumors and medical records - results of the Heinz Nixdorf Recall Study. *Cancer Epidemiol.* 2014;38(2):181–92. <https://doi.org/10.1016/j.canep.2014.01.002>.
 19. UNESCO Institute for Statistics. International Standard Classification of Education - ISCED 1997; 2006 [Accessed 2015 June 30]. Available from: URL:<http://www.uis.unesco.org/Library/Documents/iscsed97-en.pdf>.
 20. Fritschi L, Glass DC, Heyworth JS, Aronson K, Girschik J, Boyle T, et al. Hypotheses for mechanisms linking shiftwork and cancer. *Med Hypotheses.* 2011;77(3):430–6. <https://doi.org/10.1016/j.mehy.2011.06.002>.
 21. Erren TC, Morfeld P. Computing chronodisruption: how to avoid potential chronobiological errors in epidemiological studies of shift work and cancer. *Chronobiol Int.* 2014;31(4):589–99. <https://doi.org/10.3109/07420528.2013.877921>.
 22. Roenneberg T, Mrosovsky M. Entrainment of the human circadian clock. *Cold Spring Harb Symp Quant Biol.* 2007;72:293–9. <https://doi.org/10.1101/sqb.2007.72.043>.
 23. Kubo T, Oyama I, Nakamura T, Kunimoto M, Kadowaki K, Otomo H, et al. Industry-based retrospective cohort study of the risk of prostate cancer among rotating-shift workers. *Int J Urol.* 2011;18(3):206–11. <https://doi.org/10.1111/j.1442-2042.2010.02714.x>.
 24. Kubo T, Ozasa K, Mikami K, Wakai K, Fujino Y, Watanabe Y, et al. Prospective cohort study of the risk of prostate cancer among rotating-shift workers: findings from the Japan collaborative cohort study. *Am J Epidemiol.* 2006;164(6):549–55. <https://doi.org/10.1093/aje/kwj232>.
 25. Conlon M, Lightfoot N, Kreiger N. Rotating shift work and risk of prostate cancer. *Epidemiology.* 2007;18(1):182–3. <https://doi.org/10.1097/01.ede.0000249519.33978.31>.
 26. Kakizaki M, Inoue K, Kuriyama S, Sone T, Matsuda-Ohmori K, Nakaya N, et al. Sleep duration and the risk of prostate cancer: the Ohsaki Cohort Study. *Br J Cancer.* 2008;99(1):176–8. <https://doi.org/10.1038/sj.bjc.6604425>.
 27. Schwartzbaum J, Ahlbom A, Feychting M. Cohort study of cancer risk among male and female shift workers. *Scand J Work Environ Health.* 2007;33(5):336–43. <https://doi.org/10.5271/sjweh.1150>.
 28. Kolstad HA. Nightshift work and risk of breast cancer and other cancers—a critical review of the epidemiologic evidence. *Scand J Work Environ Health.* 2008;34(1):5–22. <https://doi.org/10.5271/sjweh.1194>.
 29. Costa G, Haus E, Stevens R. Shift work and cancer - considerations on rationale, mechanisms, and epidemiology. *Scand J Work Environ Health.* 2010;36(2):163–79. <https://doi.org/10.5271/sjweh.2899>.
 30. Flynn-Evans EE, Mucci L, Stevens RG, Lockley SW. Shiftwork and prostate-specific antigen in the National Health and Nutrition Examination Survey. *J Natl Cancer Inst.* 2013;105(17):1292–7. <https://doi.org/10.1093/jnci/djt169>.
 31. Romano A, Vigna L, Belluigi V, Conti DM, Barberi CE, Tomaino L, et al. Shift work and serum 25-OH vitamin D status among factory workers in Northern Italy: Cross-sectional study. *Chronobiol Int.* 2015;32(6):842–7. <https://doi.org/10.3109/07420528.2015.1048867>.
 32. Islami F, Moreira DM, Boffetta P, Freedland SJ. A systematic review and meta-analysis of tobacco use and prostate cancer mortality and incidence in prospective cohort studies. *Eur Urol.* 2014;66(6):1054–64. <https://doi.org/10.1016/j.eururo.2014.08.059>.
 33. Morgan RM, Steele RJC, Nabi G, McCowan C. Socioeconomic variation and prostate specific antigen testing in the community: a United Kingdom based population study. *J Urol.* 2013;190(4):1207–12. <https://doi.org/10.1016/j.juro.2013.04.044>.
 34. Guessous I, Cullati S, Fedewa SA, Burton-Jeangros C, Courvoisier DS, Manor O, et al. Prostate cancer screening in Switzerland: 20-year trends and socioeconomic disparities. *Prev Med.* 2016;82:83–91. <https://doi.org/10.1016/j.ypmed.2015.11.009>.

Received for publication: 20 January 2017