Scand J Work Environ Health 2017;43(6):550-559

<https://doi.org/10.5271/sjweh.3676>

Published online: 06 Oct 2017, Issue date: 01 Nov 2017

Mesothelioma incidence and asbestos exposure in Italian national priority contaminated sites

by [Binazzi A](#), [Marinaccio A](#), [Corfiati M](#), [Bruno C](#), [Fazzo L](#), [Pasetto R](#), [Pirastu R](#), [Biggeri A](#), [Catelan D](#), [Comba P](#), [Zona A](#)

The originality of this paper concerns the detection of mesothelioma excesses not only in sites where asbestos is reported as a source of contamination but also in areas labeled for environmental remediation due to other sources of pollution. This confirms the wide range of working and living environments affected by asbestos exposure and the need for communication plans with the affected communities.

Affiliation: Department of Occupational and Environmental Medicine, Epidemiology, Hygiene. National Institute for Insurance against Accidents at Work (INAIL), Via Stefano Gradi 55, 00143 Rome, Italy. a.binazzi@inail.it

The following article refers to this text: [2019;45\(5\):444-449](#)

Key terms: [asbestos](#); [asbestos exposure](#); [contaminated site](#); [environmental risk](#); [epidemiological surveillance](#); [Italy](#); [mesothelioma](#); [occupational disease](#); [polluted area](#)

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/28985440

This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Mesothelioma incidence and asbestos exposure in Italian national priority contaminated sites

by Alessandra Binazzi, PhD,¹ Alessandro Marinaccio, MSc,¹ Marisa Corfiati, PhD,¹ Caterina Bruno, MD,² Lucia Fazzo, MSc,² Roberto Pasetto, DSC,² Roberta Pirastu, MSc,³ Annibale Biggeri, MD,⁴ Dolores Catelan, PhD,⁴ Pietro Comba, PhD,² Amerigo Zona, MD²

Binazzi A, Marinaccio A, Corfiati M, Bruno C, Fazzo L, Pasetto R, Pirastu R, Biggeri A, Catelan D, Comba P, Zona A. Mesothelioma incidence and asbestos exposure in Italian national priority contaminated sites. *Scand J Work Environ Health*. 2017;43(6):550–559. doi:10.5271/sjweh.3676

Objectives This study aimed to (i) describe mesothelioma incidence in the Italian national priority contaminated sites (NPCS) on the basis of data available from the Italian National Mesothelioma Registry (ReNaM) and (ii) profile NPCS using Bayesian rank analysis.

Methods Incident cases of mesothelioma and standardized incidence ratios (SIR) were estimated for both genders in each of the 39 selected NPCS in the period 2000–2011. Age-standardized rates of Italian geographical macro areas were used to estimate expected cases. Rankings of areas were produced by a hierarchical Bayesian model. Asbestos exposure modalities were discussed for each site.

Results In the study period, 2683 incident cases of mesothelioma (1998 men, 685 women) were recorded. An excess of mesothelioma incidence was confirmed in sites with a known past history of direct use of asbestos (among men) such as Balangero (SIR 197.1, 95% CI 82.0–473.6), Casale Monferrato (SIR 910.7, 95% CI 816.5–1012.8), and Broni (SIR 1288.5, 95% CI 981.9–1691.0), in sites with shipyards and harbors (eg, Trieste, La Spezia, Venice, and Leghorn), and in settings without documented direct use of asbestos. The analysis ranked the sites of Broni and Casale Monferrato (both genders) and Biancavilla (only for women) the highest.

Conclusions The present study confirms that asbestos pollution is a risk for people living in polluted areas, due to not only occupational exposure in industrial settings with direct use of asbestos but also the presence of asbestos in the environment. Epidemiological surveillance of asbestos-related diseases is a fundamental tool for monitoring the health profile in NPCS.

Key terms environmental risk; epidemiological surveillance; Italy; occupational disease; polluted area.

The World Health Organization (WHO) has defined contaminated sites as areas that have produced or might produce in the future an environmental contamination, which results or could result in human health impacts (1). The European Environmental Agency (EEA) estimates there are 342 000 contaminated sites (CS) in the 39 EEA countries on the basis of the soil contamination regulation; one third has already been identified and about 15% have been remediated (2). In Italy, a total of 57 sites have been defined as national priority contaminated sites

(NPCS) in 2009, due to the relevance of soil and water contamination and the resulting potential health impact on the basis of European and national regulation. Epidemiological surveillance systems are essential in analyzing the health profile of populations residing in contaminated sites and for testing the associations between environmental exposure and health outcomes (3).

In the framework of the national strategic program “Environment and Health”, established by the Italian Ministry of Health and coordinated by the National

¹ Department of Occupational and Environmental Medicine, Epidemiology and Hygiene, National Institute for Insurance against Accidents at Work (INAIL), Rome, Italy.

² Department of Environment and Health, Italian National Institute of Health, Rome, Italy.

³ Department of Biology and Biotechnologies Charles Darwin, “Sapienza” University of Rome, Italy.

⁴ Department of Statistics, Computer Science, Applications “G. Parenti”, University of Florence, Italy.

Correspondence to: Alessandra Binazzi, PhD, Department of Occupational and Environmental Medicine, Epidemiology, Hygiene, National Institute for Insurance against Accidents at Work (INAIL), Via Stefano Gradi 55, 00143 Rome, Italy. [E-mail: a.binazzi@inail.it]

Health Institute, the “SENTIERI project” (an epidemiological study of residents health in NPCS) was conducted with the aim of monitoring the health effects of environmental contaminations. The SENTIERI project measured an excess of deaths (9969 for all causes and 4309 for cancer) in the period 1995–2002 among residents in NPCS (4). In 18 NPCS served by a cancer registry, an excess of cancer incidence (about 9% among men and 7% among women) was observed in the framework of the same project (5). Among the contaminants listed by specific decrees that define the NPCS boundaries, asbestos (or asbestos-like fibres) was reported together with additional pollutant sources in 12 of the 44 sites investigated by the SENTIERI project, and as the only pollutant source in 4 (namely, the sites of Broni in Lombardy, Casale Monferrato in Piedmont, Bari in Apulia and Biancavilla in Sicily). In Italy, 3 748 550 tons of raw asbestos have been produced up to 1992 (year of definite asbestos ban), with a peak between 1976 and 1980 of >160 000 tons/year (6). For this reason, Italy is one of the countries more sensitive to the issues of monitoring and preventing asbestos-related health effects. Malignant mesothelioma (MM) is an uncommon neoplasm with high mortality that typically originates in mesothelial cells coating the serous cavities, mainly the pleura and the peritoneum and, to a lesser extent, also the pericardium and testicular tunica vaginalis. The risk of MM attributable to occupational asbestos exposure has been reported to range between 86–95%, according to epidemiological studies (7–9), but such percentages must be considered with caution as they are actually place- and time-specific.

The relevance of environmental exposure to asbestos for the occurrence of mesothelioma in Italy was recently estimated (10), and residences near asbestos-cement plants were frequently reported as a documented source of risk for mesothelioma in analytical epidemiological studies carried out in Casale Monferrato (11), Bari (12), Broni (13), La Spezia (included in the NPCS of Pitelli) (14). While the extent of occupational exposure to asbestos is expected to decrease in the next years, the contribution of different patterns of non-occupational exposures is likely underestimated due to exposure levels that are usually lower than in the workplace, although not negligible and reliably associated with the risk of mesothelioma (15).

As a consequence of the relevant and widespread use of asbestos in Italy, a permanent MM surveillance system has been active since 2002 through the National Register of Malignant Mesotheliomas (“Registro Nazionale dei Mesoteliomi – ReNaM” in Italian) at the National Institute for Insurance against Accidents at Work (INAIL). ReNaM performs active research of MM cases and asbestos-exposure assessment under standardized procedures and methods (16). Information

about demographic, clinical, occupational, residential, and familial histories of MM cases are collected, and analytical studies about the incidence, epidemiological parameters of the diseases, territorial distribution of cases, and the modalities of asbestos exposure have been published (6, 10, 17). The aim of this study was to estimate the relative risk of MM incidence, provided by the ReNaM, and rank the risk of diseases for the population residing in NPCS.

Methods

The observed MM cases were extracted from the ReNaM archives in the period of diagnosis 2000–2011. In accordance with the law, the national register obtains data of incident MM cases from the Regional Operating Centres [Centri Operativi Regionali (COR) in Italian], currently established in all Italian regions. Diagnostic coding criteria are established by means of a 3-class scale of decreasing level of certainty: certain, probable, and possible MM (18). Occupational history, lifestyle habits, and residential histories are obtained using a standardized questionnaire completed by the subject or the next of kin. The classification of occupational exposure is qualitative, identified as: definite, probable, or possible. In the present analysis, the three levels of occupational exposure (definite, probable, possible) were considered together. Specific codes were

Figure 1. Geographical distribution of the 39 national priority contaminated sites and number of malignant mesothelioma cases in the period 2000–2011.

assigned to environmental exposure (for a MM subject residing near a source of asbestos pollution, without any work-related exposure), familial exposure (when patients have lived with a cohabitant who has been occupationally exposed), and leisure activities exposure (other non-occupational exposures, likely due to leisure-time activities). Details are extensively described in the ReNaM national guidelines (19).

ReNaM is operational across all of Italy, but MM case lists from the regions of Sardinia and Calabria cannot be considered complete in the study period. For this reason, of the 44 Italian NPCS previously evaluated as eligible for the epidemiological assessment in the context of the SENTIERI projects, we excluded 3 sites (Sulcis-Iglesiente-Guspinese, Aree industriali Porto Torres in Sardinia, and Crotone in Calabria). Moreover, the corresponding COR did not detect any MM cases in 2 sites (Emaiese in Valle d'Aosta and Tito in Basilicata), in the period 2000–2011, and they too have been excluded (figure 1). The time period of study has been restricted to 2001–2011 for the sites of Bacino idrografico fiume Sacco, Litorale Domizio Flegreo and Area Litorale Vesuviano, to 2005–2011 for the site of Bolzano, and to 2006–2011 for the site of Terni, due to unavailability of incidence data in the previous years.

Age-standardized incidence rates of mesothelioma (with direct method using the Italian population in 2010 as reference; all anatomical sites included) and trends over time for the overall NPCS were calculated in the period 2000–2011 by gender.

In each NPCS included in the analyses, the SIR for MM in any anatomical site (pleura, peritoneum, pericardium, and tunica vaginalis testis) were estimated for both genders separately. Gender- and age-specific rates of MM in the four Italian geographical macro areas (North-East, North-West, Centre, South-Islands) were calculated to derive expected numbers. The population of the macro areas was used as a reference in order to highlight the excesses of mesothelioma risk with a non-homogenous territorial distribution, taking into account the different industrial development across Italy. Therefore, each macro area must be considered individually, and the SIR emphasize the impact of asbestos exposure in every analyzed NPCS.

Confidence interval (CI) at 95% were estimated for each SIR according to Poisson distribution (20) and applying Byar's approximation when >100 cases were observed (21).

Hierarchical Bayesian models, described elsewhere in detail (22), were specified on SIR. In brief, let Y_i , the observed number of cases in the i^{th} NPCS, follow a Poisson distribution with mean $E_i\theta_i$, the expected number of cases times the parameter θ_i , the unknown relative risk (RR) for the i^{th} NPCS. We assumed that for each area, $\log(\theta_i)$ was independently drawn from a normal distribu-

tion with mean parameter following a weakly informative normal hyperprior distribution with zero mean and precision 0.0001 and precision parameter as inverse Gamma (23). Rank posterior distributions of RR were obtained from MCMC runs. The rank of the parameter of interest (RR_i) was computed at each iteration. The MCMC ran approximate the joint cumulative posterior distribution of $RR_i F(\theta_i|Y)$ and hence it was possible to approximate the posterior distribution $[R_i|Y] \forall i$ and its summaries (for example the posterior mean as point estimate of the rank). Notice the posterior mean of the rank is usually not integer and is shrunken toward the mid-rank. From the posterior rank distribution, 80% CI can directly be obtained (24).

All statistical analyses were carried out by IBM SPSS statistics software (IBM Statistical Package for Social Sciences for Windows, Version 22.0. Armonk, New York, USA) and WinBugs14 (25).

Results

In the analysis of the 39 NPCS, 2683 MM cases (1998 men and 685 women) were detected in the period 2000–2011. The corresponding figure in Italy in the same period is 16 837 MM incident cases (4872 women and 11 965 men).

Pleural site is largely predominant (2535 cases, 94.5%), but peritoneal cases (142), pericardial (3), and tunica vaginalis testis (3) cases were also registered. The overall study findings are reported in table 1. A statistically significant ($P<0.05$) excess of mesothelioma incidence was estimated in 20 NPCS for men (of which 9 also for women), namely: Casale Monferrato (men and women), Cengio e Saliceto, Broni (men and women) and Pitelli in the North West of Italy; Venezia (men and women), Laguna di Grado e Marano and Trieste (men and women) in the North East; Massa Carrara, Livorno (men and women), Piombino, Falconara Marittima and Basso bacino Fiume Chienti in Central Italy; Litorale Domizio Flegreo, Area Litorale Vesuviano, Bari (men and women), Taranto (men and women), Milazzo, Gela, Biancavilla (men and women), and Priolo (men and women) in Southern Italy. A defect in male MM incidence cases was observed in the sites of Bolzano and Trento Nord (North East). Further excesses, but not statistically significant, were found in 8 NPCS for men, and 11 for women.

Trends in incidence for the period 2000–2011 reveal a peak in 2007 (age-standardized rate: 8.67 and 2.89 per 100 000 inhabitants among men and women, respectively), with a minimum in 2000 for men (5.79), whereas the trend is constant for women. In 2011, age-standardized rates are equal to 7.26 among men and 2.68 among women.

Table 1. Incident mesothelioma cases (N), standardized incidence ratio (SIR) and 95% confidence interval (95% CI) in Italian national priority contaminated sites (NPCSs) by gender.

Region	NPCS	Acronym	Men			Women		
			N	SIR	95% CI	N	SIR	95% CI
Piedmont	North-Western Italy							
	Balangero	BAL	5	197.1	82.0–473.6	.	.	..
	Casale Monferrato	CAS	340	910.7	816.5–1012.8	248	1338.1	1176.7–1515.4
	Serravalle Scrivia	SER	1	28.2	4.0–200.0	1	64.4	9.1–456.9
Piedmont, Liguria	Pieve Vergonte	PIV	1	43.6	6.1–309.2	.	.	..
	Cengio e Saliceto	CES	31	165.8	116.6–235.7	8	98.0	49.0–195.9
	Cerro al Lambro	CER	2	83.2	20.8–332.5	1	97.9	13.8–695.1
	Pioltello Rodano	PIR	11	101.4	56.2–183.1	6	123.9	55.6–275.7
Lombardy	Sesto San Giovanni	SES	37	80.6	58.4–111.2	18	81.5	51.3–129.3
	Brescia Caffaro	BRE	50	68.8	52.1–90.8	26	65.7	44.7–96.5
	Broni	BRO	52	1288.5	981.9–1691.0	43	2006.7	1488.3–2705.8
	Laghi Di Mantova E Polo Chimico	LMN	14	61.4	36.4–103.7	7	55.2	26.3–115.9
Liguria	Cogoleto Stoppani	COS	14	154.6	91.6–261.0	3	69.9	22.5–216.7
	Pitelli	PIT	201	445.4	385.9–511.4	30	124.5	87.1–178.1
	North-Eastern Italy							
A.P. Bolzano	Bolzano	BOL	4	17.5	6.6–46.7	.	.	..
A.P. Trento	Trento Nord	TRE	7	30.5	14.5–64.0	3	37.1	12.0–115.1
Veneto	Venezia	VEN	127	181.0	150.9–215.4	36	144.5	104.2–200.3
Friuli Venezia Giulia	Laguna di Grado e Marano	LGM	15	196.7	118.6–326.2	2	82.3	20.6–329.0
Emilia Romagna	Trieste	TRI	209	374.5	325.4–428.9	42	204.4	151.0–276.5
	Fidenza	FID	11	98.2	54.4–177.4	7	185.5	88.4–389.2
	Sassuolo - Scandiano	SAS	18	80.6	50.8–127.9	2	29.0	7.3–116.1
Tuscany	Central Italy							
	Massa Carrara	MSC	46	243.4	182.3–324.9	10	149.6	80.5–278.1
	Livorno	LIV	113	429.6	354.0–516.5	25	277.1	187.3–410.1
	Piombino	PIO	17	287.6	178.8–462.7	2	100.4	25.1–401.6
Umbria	Orbetello	ORB	2	88.0	22.0–351.8	1	128.7	18.1–913.7
	Terni – Papigno ^a	TER	20	118.7	76.6–184.0	3	51.9	16.7–160.9
Marche	Falconara Marittima	FAL	17	401.0	249.3–645.1	1	72.2	10.2–512.6
	Basso Bacino Fiume Chienti	BBC	17	128.2	138.1–357.4	3	70.9	38.7–372.5
Lazio	Bacino Idrografico Fiume Sacco ^b	BFS	18	149.9	94.4–237.9	1	26.3	3.7–186.6
Campania	Southern Italy and Isles							
	Litorale Domizio Flegreo e Agro Aversano ^b	LDF	154	150.0	130.4–180.0	36	105.5	79.5–152.7
Apulia	Area Litorale Vesuviano ^b	ALV	133	336.3	281.5–398.5	20	152.7	98.5–236.6
	Manfredonia	MAN	6	90.2	40.5–200.8	1	48.3	6.8–343.2
	Bari - Fibronit	BAR	88	271.3	220.2–334.3	35	322.2	231.4–448.8
	Taranto	TAR	85	417.0	337.1–515.8	25	355.2	136.0–297.8
Basilicata	Brindisi	BRI	13	154.9	89.9–266.7	3	107.0	34.5–331.7
	Aree Industriali Val Basento	AVB	7	173.4	82.7–363.8	.	.	..
	Milazzo	MIL	11	238.8	132.2–431.2	2	138.1	34.5–552.3
	Gela	GEL	14	215.3	127.5–363.5	3	157.7	50.9–489.1
Sicily	Biancavilla	BIA	9	436.6	227.2–839.1	13	1941.4	1127.3–3343.5
	Priolo	PRI	78	447.7	358.6–559.0	18	330.2	208.1–524.1

^a Incidence period limited to 2006–2011.^b Incidence period limited to 2001–2011.

Most of the cases in the NPCS were diagnosed as certain MM (84% and 79% among men and women, respectively, figure 2).

An individual questionnaire was used to investigate the modalities of asbestos exposure in 2150 MM cases (80.1% of the whole case list: 81.7% among men and decreasing to 75.6% among women). The amount of non-interviewed subjects vary according to the NPCS, with a decreasing trend from the North to the South of Italy, and non-negligible percentages (from 20% to 50%) in critical NPCS, such as Casale Monferrato, Litorale Domizio Flegreo or Area Litorale Vesuviano (table 2a).

Among cases evaluated for exposure, an occupational exposure to asbestos has been ascertained for

70.6% of cases from the overall case list, with a significant difference by gender (87% among men and 20% among women). The COR assigned non-occupational exposure (environmental, family and leisure-time related) to 409 cases (19%), with the most being represented by women (55%) (figure 3). In a percentage of MM, asbestos exposure could not be identified on the basis of available information, with different patterns between men (6%) and women (25%). The modalities of asbestos exposure by NPCS for all MM cases included in the analyses are reported in table 2b.

Profiling NPCS by Bayesian rank analysis highlights the firsts positions, consistently on both sexes (figures 4a, 4b). The site of Broni has the worse position for both

Figure 2. Level of diagnostic certainty for malignant mesothelioma cases in Italian NPCCs in the period 2000–2011, by gender.

Figure 3. Modalities of asbestos exposure for malignant mesothelioma cases in Italian NPCCs in the period 2000–2011, by gender.

Table 2a. Mesothelioma cases with exposure defined by interview in Italian national priority contaminated sites (NPCS), by gender. See table 1 for complete names.

NPCS	Men			Women		
	All	Interviewed		All	Interviewed	
	N	N	%	N	N	%
BAL	5	5	100.0	.	.	.
CAS	340	272	80.0	248	176	71.0
SER	1	0	0.0	1	.	.
PIV	1	0	0.0	.	.	.
CES	31	25	80.6	8	5	62.5
CER	2	2	100.0	1	1	100.0
PIR	11	10	90.9	6	5	83.3
SES	37	37	100.0	18	16	88.9
BRE	50	49	98.0	26	22	84.6
BRO	52	52	100.0	43	43	100.0
LMN	14	14	100.0	7	6	85.7
COS	14	11	78.6	3	2	66.7
PIT	201	199	99.0	30	29	96.7
BOL	4	4	100.0	.	.	.
TRE	7	6	85.7	3	3	100.0
VEN	127	120	94.5	36	32	88.9
LGM	15	15	100.0	2	2	100.0
TRI	209	194	92.8	42	32	76.2
FID	11	10	90.9	7	6	85.7
SAS	18	18	100.0	2	1	50.0
MSC	46	45	97.8	10	9	90.0
LIV	113	109	96.5	25	25	100.0
PIO	17	17	100.0	2	2	100.0
ORB	2	2	100.0	1	1	100.0
TER	20	20	100.0	3	3	100.0
FAL	17	16	94.1	1	1	100.0
BBC	17	16	94.1	3	2	66.7
BFS	18	13	72.2	1	.	.
LDF	154	62	40.3	36	12	33.3
ALV	133	71	53.4	20	10	50.0
MAN	6	5	83.3	1	1	100.0
BAR	88	79	89.8	35	31	88.6
TAR	85	77	90.6	25	20	80.0
BRI	13	13	100.0	3	3	100.0
AVB	7	6	85.7	.	.	.
MIL	11	10	90.9	2	1	50.0
GEL	14	6	42.9	3	1	33.3
BIA	9	4	44.4	13	10	76.9
PRI	78	18	23.1	18	5	27.8
Total	1998	1632	81.7	685	518	75.6

genders. The sites of Casale Monferrato (both genders), and Biancavilla (only for women), resulted also among the higher positions, followed by the sites of Livorno (5th in men and 7th in women), Taranto (6th in men and 4th in women), Trieste (9th in men and 8th in women). Among the other sites, the degree of overlap of credible intervals is so large that it is impossible to clearly state an ordering by SIR.

Discussion

Through the SENTIERI approach, this study is the first attempt to combine the analyses of health profiles in the NPCS population with individual data from a cancer register (ReNaM), which includes the exposure and anamnestic history of the affected people.

The epidemiological surveillance of the health impact of contamination in NPCS contributes to the definition of priorities for remedial programs and public health interventions, and the SENTIERI project has been developed for this purpose. SENTIERI studied mortality, cancer incidence, and hospital discharge records of residents in NPCS specifically focusing on causes of death for which environmental exposure is suspected or ascertained to play an etiologic role (4). The epidemiological evidence of the causal association between diseases death and environmental exposure was classified a priori into three categories: sufficient (S), limited (L), and inadequate (I). The WHO defined this approach as a first-level stage in the description of the health status of residents in contaminated sites (1).

The systematic surveillance of mesothelioma incidence in Italy by means of a national register, which allows a qualitative assessment of asbestos exposure through the analysis of the entire occupational, residential and family history of each MM case, is an original

Table 2b. Modalities of asbestos exposure in mesothelioma cases with exposure defined by interview in Italian national priority contaminated sites (NPCS), by gender. See table 1 for complete names.

NPCS	Men						Women					
	Work-related		Non work-related		Unlikely or unknown		Work-related		Non work-related		Unlikely or unknown	
	N	%	N	%	N	%	N	%	N	%	N	%
BAL	3	60.0	2	40.0	.	.	.	.	.	.	.	.
CAS	198	72.8	71	26.1	3	1.1	37	21.0	135	76.7	4	2.3
SER	.	.	.	.	.	.	.	.	.	.	.	.
PIV	.	.	.	.	.	.	.	.	.	.	.	.
CES	24	96.0	.	.	1	4.0	2	40.0	.	.	3	60.0
CER	1	50.0	.	.	1	50.0	.	.	.	.	1	100.0
PIR	10	100.0	.	.	.	.	2	40.0	1	20.0	2	40.0
SES	34	91.9	1	2.7	2	5.4	4	25.0	1	6.3	11	68.8
BRE	37	75.5	2	4.1	10	20.4	4	18.2	1	4.5	17	77.3
BRO	40	76.9	12	23.1	.	.	8	18.6	35	81.4	0	0
LMN	10	71.4	.	.	4	28.6	.	.	.	.	6	100.0
COS	9	81.8	.	.	2	18.2	1	50.0	.	.	1	50.0
PIT	189	95.0	2	1.0	8	4.0	3	10.3	17	58.6	9	31.0
BOL	3	75.0	1	25.0	.	.	.	.	.	.	.	.
TRE	4	66.7	.	.	2	33.3	.	.	2	66.7	1	33.3
VEN	109	90.8	5	4.2	6	5.0	14	43.8	11	34.4	7	21.9
LGM	14	93.3	.	.	1	6.7	.	.	2	100.0	.	.
TRI	175	90.2	.	.	19	9.8	4	12.5	12	37.5	16	50.0
FID	9	90.0	.	.	1	10.0	2	33.3	3	50.0	1	16.7
SAS	16	88.9	.	.	2	11.1	1	100.0	.	.	.	.
MSC	41	91.1	.	.	4	8.9	2	22.2	3	33.3	4	44.4
LIV	105	96.3	1	0.9	3	2.8	6	24.0	7	28.0	12	48.0
PIO	16	94.1	1	5.9	.	.	.	.	1	50.0	1	50.0
ORB	2	100.0	.	.	.	.	.	.	.	.	1	100.0
TER	20	100.0	.	.	.	.	2	66.7	.	.	1	33.3
FAL	14	87.5	1	6.3	1	6.3	.	.	.	.	1	100.0
BBC	14	87.5	.	.	2	12.5	.	.	1	50.0	1	50.0
BFS	13	100.0	.	.	.	.	.	.	.	.	.	.
LDF	54	87.1	4	6.5	4	6.5	3	25.0	4	33.3	5	41.7
ALV	63	88.7	5	7.0	3	4.2	2	20.0	4	40.0	4	40.0
MAN	5	100.0	.	.	.	.	1	100.0	.	.	.	.
BAR	63	79.7	9	11.4	7	8.9	4	12.9	19	61.3	8	25.8
TAR	71	92.2	2	2.6	4	5.2	.	.	13	65.0	7	35.0
BRI	13	100.0	.	.	.	.	.	.	2	66.7	1	33.3
AVB	6	100.0	.	.	.	.	.	.	.	.	.	.
MIL	7	70.0	2	20.0	1	10.0	.	.	1	100.0	.	.
GEL	4	66.7	.	.	2	33.3	.	.	.	.	1	100.0
BIA	2	50.0	1	25.0	1	25.0	.	.	9	90.0	1	10.0
PRI	16	88.9	1	5.6	1	5.6	1	20.0	2	40.0	2	40.0
Total	1414	86.6	123	7.5	95	5.8	103	19.9	286	55.2	129	24.9

and substantial experience in the international framework for managing and monitoring asbestos-related diseases (26). The occurrence of asbestos-related diseases in NPCS is relevant considering that in 10 out of 39 NPCS included in this analysis asbestos is one of the contaminants cited in the decrees defining the sites' boundaries and the sources of pollution. In four sites (Casale Monferrato, Broni, Bari-Fibronit and Biancavilla), asbestos pollution is the only source of health risk identified in the decrees. In the Biancavilla site, the amphibole fluoro-edenite, mined from the local quarries and found in the soil and building materials, has been identified as responsible for the outbreak of mesotheli-

oma observed in the town, as previously discussed (27).

Trends of incidence of mesothelioma in the overall NPCS show high figures among both genders that in 2011 are almost twice that of the national values (16). Such evidence is validated by the inclusion among the NPCS of sites such as Broni, Casale Monferrato, Biancavilla, whose incidence rates are very high due to the presence of important asbestos-cement plants in the first two sites and to the environmental exposure from the presence of a quarry with fluoro-edenite contamination in Biancavilla.

The reliability of diagnosis is similar in NPCS and in the rest of the ReNaM database, where the figures of the three levels of diagnostic certainty ("certain", "probable", "possible") are: 82%, 9%, 9% for men, and 76%, 12%, 12% for women, respectively.

However, some critical limitations in the exposure assessment need to be discussed. The modalities of identifying asbestos exposure are not always fully uniform in the ReNaM network, and the percentage of interviewed subjects varies between 45% and 95% according to the COR, depending on their available resources and knowledge. As a consequence, the level of interviewed subjects in the NPCS is heterogeneous among different areas. It is one of the main limitation of this paper, and regards mainly NPCS such as Casale Monferrato, where the expositive history is certainly significant.

Also the procedures for coding and classifying the diagnosis of MM, established at national level, still lack of homogeneity among the CORs. Another limitation of this study is the use of the municipality of residence (at the time of diagnosis) to allocate the distribution of MM cases, and this is problematic when the subject resides outside the contaminated site where the past working activity was carried out.

Moreover, another limit is represented by the features of the ReNaM, which is designated by order of Italian law to record any mesothelioma case exclusively arising from the pleura, peritoneum, pericardium, and tunica vaginalis testis. No other anatomical sites are included. In perspective the ReNaM should extend the network of recording mesothelioma also to other asbestos-related neoplasms (eg, lung, larynx, ovary) (28).

A significant excess of MM cases has been found in 20 of the 39 NPCS selected for this study (in 9 sites for both men and women, in 11 sites only for men). Remarkably, in 12 of these (60%), asbestos was not included among the sources of pollution by the official decree. These sites were: Cengio e Saliceto, Venezia, Laguna Grado e Marano, Trieste, Falconara Marittima, Livorno, Piombino, Bacino Idrografico Fiume Sacco, Litorale Domizio Flegreo ed Agro Aversano, Taranto, Milazzo, and Gela.

The percentages of "definite", "probable", "possible" occupational exposure are 42.5%, 5.2%, and

Figure 4. Posterior rank estimates and 80% credibility intervals for malignant mesothelioma in Italian NPCSs in the period 2000-2011; (A) men, (B) women.

8.9% respectively in the overall NPCS, corresponding to 38.4%, 5.7%, and 11% in the ReNaM database. The presence of occupational exposure to asbestos in a wide spectrum of economic activities, and not only in industrial settings where asbestos was directly used (asbestos cement plants, asbestos textile production, shipbuilding and repair, rail stock and rail carriage insulation, mining and work with friction products), has been repeatedly demonstrated in Italy (29) and elsewhere (30, 31). Our results support and confirm these findings, demonstrating that asbestos exposure is a relevant public health issue for a relevant number of contaminated sites, also without any mention of asbestos as direct cause of contamination.

In this study, we used 95% CI as opposed to the 90% CI generally applied in the SENTIERI project. That choice was made to show the range of uncertainty for risk estimators, reducing the use of CI as surrogate of hypothesis testing.

Reporting the NPCS ranking allowed a simple and immediate way to summarize the information about the profile of mesothelioma RR among the resident populations. Methodologically, it is important to underline that we provide an estimate of both the rank and its uncertainty. This is not trivial, and the degree of overlap in the CI is able to communicate to the reader the reliability of a given classification.

With regard to the methodology applied, the standardization procedure is motivated by the need to take into account not only the different age structure of the studied populations but also the presence of large-scale geographical variations. The ranking is therefore “relative” to the presence of large-scale geographical patterns. In Italy, the industrialization process included important difference between northern, central and

southern regions, and asbestos exposures historically varied accordingly.

We therefore specified age and macro area-specific set of reference rates in order to gain sensitivity in the less industrialized macro areas. By observing the rank, the health impact of asbestos exposure in populations residing in the NPCS with a past history of direct asbestos use [asbestos cement plants (Broni, Casale Monferrato, Bari) or shipyards and harbours areas (Livorno, Taranto, Trieste, Piombino)] and the populations residing in territory with an environmentally diffuse contamination of fluoro-edenite (Biancavilla) emerged clearly (32).

In the present study about 20% of cases had a non-occupational exposure, predominantly among women (apart from Biancavilla, mainly in the sites of Casale Monferrato and Broni). The percentage of MM with a non-identified asbestos exposure is more relevant among women (25%) than men (6%). Such a difference underlines the need for tools investigating the modalities of exposure (ie, questionnaires) for the female population, which is more complex than the male one.

The ReNaM archive recently provided a reliable estimation of 10% of MM due to a non-occupational exposure to asbestos, based on more than 15 845 detected cases, of which 12 065 were individually interviewed (10). With respect to specific non-occupational exposures, the percentages of familial and environmental exposures are slightly higher than the corresponding figures in the ReNaM database for the same period (familial: 8% versus 5%; environmental: 10% versus 4% respectively). Non-occupational exposure to asbestos could be due to (i) naturally occurring contamination, (ii) the presence in residential areas of industrial sites previously involving asbestos use, (iii) the diffuse pres-

ence of asbestos industry by-products for insulation and for road and courtyard paving, (iv) cohabitation with exposed people, or (v) the accidental use of asbestos-containing materials. Such non occupational exposures have been analyzed extensively elsewhere (10). These various patterns pose different concerns with respect to the welfare protection framework and deserve special attention. Individuals were especially likely to be unaware of their exposure or of the associated hazard, as in the case of people living around industrial sources of asbestos pollution or with asbestos workers. The lower level of control over certain non-occupational circumstances, associated with the presence of asbestos *in-situ*, in buildings, make it possible for such exposures to persist. There is a need to discuss how to deal with compensation rights (currently reserved in many countries only for occupationally exposed people) for malignant mesothelioma cases induced by a non-occupational exposure to asbestos.

There is an open issue in Italy regarding MM cases of occupational origin that do not seek compensation (33). Furthermore, around 29% of MM cases due to non-occupational exposure in Italy has been found in NPCS and the percentage of non-occupational cases with respect to all cases investigated for exposure is higher in NPCS than in the ReNaM archive for the same period (19% versus 11% respectively). It is remarkable that recently the compensation of all MM cases, regardless of the modality or different patterns of exposures (naturally occurring contamination, industrial activities involving asbestos use existing in residential areas, cohabitation with exposed people, accidental use of asbestos-containing materials), has been implemented in France (34).

Our analyses confirm that asbestos pollution is a real concern in the NPCS, due not only to the occupational exposure of people working inside industrial plants where asbestos was directly used as a raw material or in workplaces where asbestos was present but also to the environmental exposure. Furthermore, non-occupational exposure to asbestos is largely the predominant cause of female mesothelioma cases in NPCS. A non-negligible proportion of mesothelioma cases identified at national level is concentrated in the NPCS: 16% of the whole ReNaM database (in the period 2000–2011), against an overall NPCS population representing the 9% of the Italian one. Moreover, the detection of significant mesothelioma excesses not only in NPCS, where asbestos is explicitly reported as a source of contamination, but also in a number of areas with other sources of pollution, confirms the wide range of activities and working and living environments affected by asbestos exposure, which are not restricted to the industrial sectors characterized by the direct use of this material.

The evaluation of occupational and environmental

risk for people living in polluted areas, and the assessment of the risk produced by industrial settlements, could also be achievable by means of data available from surveillance systems of occupational tumors. The ReNaM-SENTIERI approach underlines the relevant role of such systems in the investigation of the associations between environmental exposures and health effects in polluted sites and reveals how this synergy can be successful for the permanent epidemiologic surveillance of occupational risks in contaminated sites.

In future, the aim is to extend the surveillance beyond mesothelioma to all occupational diseases in the NPCS by using the archives of compensated cases of occupational diseases available at INAIL. As a consequence, a permanent program for the epidemiological surveillance of occupational safety and health could be implemented in these areas where, when integrated with the analysis of environmental risks, results can improve and refine the interpretation of the health profile of populations affected by NPCS.

Acknowledgements

The authors would like to thank COR staff for data collection and transmission to the ReNaM and Dr. Alberto Scarselli of the Department of Occupational and Environmental Medicine, Epidemiology, Hygiene of INAIL for editing figure 1.

Conflict of interest

The authors declare no conflict of interest.

References

1. Pasetto R, Martin Olmedo P, Martuzzi M, editors. Contaminated sites and health. In: Report of two WHO workshops (Syracuse, Italy, 18 November 2011; Catania, Italy, 21–22 June 2012). Copenhagen (Denmark): WHO Regional Office for Europe; 2013. Available from: www.euro.who.int/__data/assets/pdf_file/0003/186240/e96843e.pdf.
2. van Liedekerke M, Prokop G, Rabl-Berger S, Kibblewhite M, Louwagie G, editors. Progress in the Management of Contaminated Sites in Europe. Reference Report by the Joint Research Centre of the European Commission. Luxembourg: Publications Office of the European Union; 2014. Available from: http://eusoiis.jrc.ec.europa.eu/ESDB_Archive/eusoiis_docs/other/EUR26376EN.pdf.
3. Pirastu R, Ancona C, Iavarone I, Mitis F, Zona A, Comba P; SENTIERI Working Group SENTIERI Project. Mortality study of residents in Italian polluted sites: evaluation of the epidemiological evidence. *Epidemiol Prev*. 2010;34(5–6

- Suppl 3):1-2. English, Italian. Available from: www.epiprev.it/publicazione/epidemiol-prev-2010-34-5-6-suppl-3.
4. Pirastu R, Pasetto R, Zona A, Ancona C, Iavarone I, Martuzzi M et al. The health profile of populations living in contaminated sites: SENTIERI approach. *J Environ Public Health*. 2013;2013:939267. <http://dx.doi.org/10.1155/2013/939267> PubMed.
5. Comba P, Ricci P, Iavarone I, Pirastu R, Buzzoni C, Fusco M et al.; ISS-AIRTUM Working Group for the study of cancer incidence in contaminated sites. Cancer incidence in Italian contaminated sites. *Ann Ist Super Sanita*. 2014;50(2):186–91.
6. Marinaccio A, Binazzi A, Marzio DD, Scarselli A, Verardo M, Mirabelli D et al.; ReNaM Working Group. Pleural malignant mesothelioma epidemic: incidence, modalities of asbestos exposure and occupations involved from the Italian National Register. *Int J Cancer*. 2012 May;130(9):2146–54. PubMed <http://dx.doi.org/10.1002/ijc.26229>.
7. Rushton L, Hutchings SJ, Fortunato L, Young C, Evans GS, Brown T et al. Occupational cancer burden in Great Britain. *Br J Cancer*. 2012 Jun;107 Suppl 1:S3–7. PubMed <http://dx.doi.org/10.1038/bjc.2012.112>.
8. Rake C, Gilham C, Hatch J, Darnton A, Hodgson J, Peto J. Occupational, domestic and environmental mesothelioma risks in the British population: a case-control study. *Br J Cancer*. 2009 Apr;100(7):1175–83. <http://dx.doi.org/10.1038/sj.bjc.6604879>.
9. Lacourt A, Gramond C, Rolland P, Ducamp S, Audignon S, Astoul P et al. Occupational and non-occupational attributable risk of asbestos exposure for malignant pleural mesothelioma. *Thorax*. 2014 Jun;69(6):532–9. <http://dx.doi.org/10.1136/thoraxjnl-2013-203744>.
10. Marinaccio A, Binazzi A, Bonafede M, Corfiati M, Di Marzio D, Scarselli A et al.; ReNaM Working Group. Malignant mesothelioma due to non-occupational asbestos exposure from the Italian national surveillance system (ReNaM): epidemiology and public health issues. *Occup Environ Med*. 2015 Sep;72(9):648–55. PubMed <http://dx.doi.org/10.1136/oemed-2014-102297> </jrn>
11. Magnani C, Dalmasso P, Biggeri A, Ivaldi C, Mirabelli D, Terracini B. Increased risk of malignant mesothelioma of the pleura after residential or domestic exposure to asbestos: a case-control study in Casale Monferrato, Italy. *Environ Health Perspect*. 2001 Sep;109(9):915–9. <http://dx.doi.org/10.1289/ehp.01109915>.
12. Musti M, Pollice A, Cavone D, Dragonieri S, Bilancia M. The relationship between malignant mesothelioma and an asbestos cement plant environmental risk: a spatial case-control study in the city of Bari (Italy). *Int Arch Occup Environ Health*. 2009 Mar;82(4):489–97. <http://dx.doi.org/10.1007/s00420-008-0358-5>.
13. Mensi C, Riboldi L, De Matteis S, Bertazzi PA, Consonni D. Impact of an asbestos cement factory on mesothelioma incidence: global assessment of effects of occupational, familial, and environmental exposure. *Environ Int*. 2015 Jan;74:191–9. <http://dx.doi.org/10.1016/j.envint.2014.10.016>.
14. Dodoli D, Del Nevo M, Fiumalbi C, Iaia TE, Cristaudo A, Comba P et al. Environmental household exposures to asbestos and occurrence of pleural mesothelioma. *Am J Ind Med*. 1992;21(5):681–7. <http://dx.doi.org/10.1002/ajim.4700210508>.
15. Goldberg M, Luce D. The health impact of nonoccupational exposure to asbestos: what do we know? *Eur J Cancer Prev*. 2009 Nov;18(6):489–503. <http://dx.doi.org/10.1097/CEJ.0b013e32832f9bee>.
16. Marinaccio A, Binazzi A, Bonafede M, Branchi C, Corfiati M, Di Marzio D et al., editors. [Il Registro Nazionale dei Mesoteliomi (ReNaM). Quinto Rapporto]. Rome: INAIL Monograph; 2015. (Italian).
17. Corfiati M, Scarselli A, Binazzi A, Di Marzio D, Verardo M, Mirabelli D et al.; ReNaM Working Group. Epidemiological patterns of asbestos exposure and spatial clusters of incident cases of malignant mesothelioma from the Italian national registry. *BMC Cancer*. 2015 Apr;15:286. <http://dx.doi.org/10.1186/s12885-015-1301-2>.
18. Conti S, Minelli G, Ascoli V, Marinaccio A, Bonafede M, Manno V et al. Peritoneal mesothelioma in Italy: trends and geography of mortality and incidence. *Am J Ind Med*. 2015 Oct;58(10):1050–8. <http://dx.doi.org/10.1002/ajim.22491>.
19. Nesti M, Adamoli S, Ammirabile F, Ascoli V, Barbieri PG, Cacciarini V et al. Guidelines for the identification and definition of malignant mesothelioma cases and the transmission to Ispesl by Regional Operating centres. Rome: ISPEL Monograph; 2003. Available from: <https://www.inail.it/cs/internet/docs/renamlineeguida2005-pdf.pdf>.
20. Sahai H, Khurshid A. Confidence Intervals for the Mean of a Poisson Distribution: A Review. *Biometrical J*. 1993;35:857–67. <http://dx.doi.org/10.1002/bimj.4710350716>.
21. Rothman JJ, Boice JD. *Epidemiologic Analysis with a Programmable Calculator*. Washington DC: US Government Printing Office; 1979. NIH Publication, 76-1649.
22. Catelan D, Buzzoni C, Coviello E, Crocetti E, Pasetto R, Pirastu R et al. [Risk profiling in cancer surveillance in contaminated sites: an example from the ISS-AIRTUM collaborative study] [Italian.]. *Epidemiol Prev*. 2014 Mar-Apr;38(2 Suppl 1):162–70.
23. Wakefield J. Disease mapping and spatial regression with count data. *Biostatistics*. 2007 Apr;8(2):158–83. <http://dx.doi.org/10.1093/biostatistics/kxl008>.
24. Shen W, Louis TA. Triple-goal estimates in two-stage hierarchical models. *J R Stat Soc B*. 1998;60(2):455–71. <http://dx.doi.org/10.1111/1467-9868.00135>.
25. Lunn DJ, Thomas A, Best N, Spiegelhalter D. WinBUGS – a Bayesian modelling framework: concepts, structure, and extensibility. *Stat Comput*. 2000;10:325–37. <http://dx.doi.org/10.1023/A:1008929526011>.
26. Ferrante P, Binazzi A, Branchi C, Marinaccio A. [National epidemiological surveillance systems of mesothelioma cases]. *Epidemiol Prev*. 2016 Sep-Oct;40(5):336–43.
27. Comba P, Gianfagna A, Paoletti L. Pleural mesothelioma

- cases in Biancavilla are related to a new fluoro-edenite fibrous amphibole. *Arch Environ Health*. 2003 Apr;58(4):229–32. <http://dx.doi.org/10.3200/AEOH.58.4.229-232>.
28. Arsenic, metals, fibres, and dusts. volume 100 C. IARC Working Group on the evaluation of Carcinogenic Risks to Humans. Lyon, France; 2009. Available from: <http://monographs.iarc.fr/ENG/Monographs/vol100C/mono100C-11.pdf>. [Accessed 13 September 2017].
 29. Binazzi A, Scarselli A, Corfiati M, Di Marzio D, Branchi C, Verardo M et al.; Gruppo di lavoro ReNaM. Sorveglianza epidemiologica dei mesoteliomi per la prevenzione dell'esposizione ad amianto anche in attività non tradizionalmente coinvolte. *Epidemiol Prev*. 2013 Jan-Feb;37(1):35–42.
 30. Whitehouse AC, Black CB, Heppe MS, Ruckdeschel J, Levin SM. Environmental exposure to Libby Asbestos and mesotheliomas. *Am J Ind Med*. 2008 Nov;51(11):877–80. PubMed <http://dx.doi.org/10.1002/ajim.20620>.
 31. de Klerk N, Alfonso H, Olsen N, Reid A, Sleith J, Palmer L et al. Familial aggregation of malignant mesothelioma in former workers and residents of Wittenoom, Western Australia. *Int J Cancer*. 2013 Mar;132(6):1423–8. <http://dx.doi.org/10.1002/ijc.27758>.
 32. Bruno C, Comba P, Zona A. Adverse health effects of fluoro-edenitic fibers: epidemiological evidence and public health priorities. *Ann N Y Acad Sci*. 2006 Sep;1076:778–83. PubMed <http://dx.doi.org/10.1196/annals.1371.020>.
 33. Marinaccio A, Scarselli A, Merler E, Iavicoli S. Mesothelioma incidence surveillance systems and claims for workers' compensation. Epidemiological evidence and prospects for an integrated framework. *BMC Public Health*. 2012 Jul;12:314. <http://dx.doi.org/10.1186/1471-2458-12-314>.
 34. Goldberg M. Indennizzi per gli esposti all'amianto: il modello francese. *Epidemiol Prev*. 2008 May-Jun;32(3):129–31.

Received for publication: 22 May 2017