



## Original article

Scand J Work Environ Health 2018;44(4):385-393

doi:10.5271/sjweh.3737

### Risk of injury after evening and night work - findings from the Danish Working Hour Database

by [Nielsen HB](#), [Larsen AD](#), [Dyreborg J](#), [Hansen ÅM](#), [Pompeii LA](#), [Conway SH](#), [Hansen J](#), [Kolstad HA](#), [Nabe-Nielsen K](#), [Garde AH](#)

This study sheds new light on the association between injury and evening or night work by studying the past week and reducing unmeasured confounding. Results show a higher risk of injury after evening or night work in the past week compared to day work during the past week. Thus evening and night work should be kept to a minimum.

**Affiliation:** National Research Centre for the Working Environment, Lersø Parkallé 105, 2100 Copenhagen Ø, Denmark. [hbn@nrcwe.dk](mailto:hbn@nrcwe.dk)

Refers to the following texts of the Journal: [2015;41\(3\):268-279](#)  
[2011;37\(3\):173-185](#) [2008;34\(3\):206-212](#) [2006;32\(3\):232-240](#)  
[2001;27\(2\):97-105](#)

The following article refers to this text: [2019;45\(2\):166-173](#)

**Key terms:** [accident](#); [Danish Working Hour Database](#); [evening shift](#); [evening work](#); [injury](#); [night shift](#); [night work](#); [risk of injury](#); [shift work](#); [work pattern](#); [work schedule tolerance](#)

This article in PubMed: [www.ncbi.nlm.nih.gov/pubmed/29777613](http://www.ncbi.nlm.nih.gov/pubmed/29777613)

### Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](#).



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

## Risk of injury after evening and night work – findings from the Danish Working Hour Database

by Helena B Nielsen, MSc,<sup>1</sup> Ann D Larsen, PhD,<sup>1</sup> Johnny Dyreborg, PhD,<sup>1</sup> Åse M Hansen, PhD,<sup>1,2</sup> Lisa A Pompeii, PhD,<sup>3</sup> Sadie H Conway, PhD,<sup>3</sup> Johnni Hansen, PhD,<sup>4</sup> Henrik A Kolstad, PhD,<sup>5</sup> Kirsten Nabe-Nielsen, PhD,<sup>2</sup> Anne H Garde, PhD<sup>1,2</sup>

Nielsen HB, Larsen AD, Dyreborg J, Hansen ÅM, Pompeii LA, Conway SH, Hansen J, Kolstad HA, Nabe-Nielsen K, Garde AH. Risk of injury after evening and night work – findings from the Danish Working Hour Database. *Scand J Work Environ Health* 2018;44(4):385–393. doi:10.5271/sjweh.3737

**Objectives** Evening and night work have been associated with higher risk of injury than day work. However, previous findings may be affected by recall bias and unmeasured confounding from differences between day, evening and night workers. This study investigates whether evening and night work during the past week increases risk of injury when reducing recall bias and unmeasured confounding.

**Methods** We linked daily working hours at the individual level of 69 200 employees (167 726 person years from 2008–2015), primarily working at hospitals to registry information on 11 834 injuries leading to emergency room visits or death. Analyses were conducted with Poisson regression models in the full population including permanent day, evening and night workers, and in two sub-populations of evening and night workers, with both day and evening or night work, respectively. Thus, the exchangeability between exposure and reference group was improved in the two sub-populations.

**Results** Risk of injury was higher after a week with evening work [incidence rate ratio (IRR) 1.32, 95% confidence interval (CI) 1.26–1.37] and night work (IRR 1.33, 95% CI 1.25–1.41) compared with only day work. Similar, although attenuated, estimates were found for evening work among evening workers (IRR 1.18, 95% CI 1.12–1.25), and for night work among night workers (IRR 1.10, 95% CI 1.01–1.20).

**Conclusion** There is an overall increased risk of injury after a week that has included evening or night work compared with only day work. Though attenuated, the higher risk remains after reducing unmeasured confounding.

**Key terms** accident; evening shift; evening work; night shift; shift work; work pattern; work schedule tolerance.

Work patterns that differ from standard daytime hours, such as work in the evening or night as part of both rotating or permanent work patterns, have been associated with increased risk of injury (1–4). These work patterns are common, with 20% of employees in Denmark working schedules other than regular daytime shifts, of whom 8% work permanent night shifts (5). Among healthcare professionals in Denmark, 45% work patterns other than permanent dayshift, of whom 30% work night shift (5). The healthcare and social work sector has the high-

est number of occupational injury claims in Denmark (6). Injury, the physical harm as a consequence of an accident, both at work and outside of work is a leading cause of death among young adults in Denmark (7) and may result in disabilities (8, 9), lost work days (10) and poorer general health (11).

Sleepiness has been linked to reduced performance (12) and a higher risk of injury (12–14) and is suggested as a mechanism linking work patterns and risk of injury (15, 16). Employees working night shifts experience

<sup>1</sup> The National Research Centre for the Working Environment, Copenhagen, Denmark.

<sup>2</sup> Department of Public Health, University of Copenhagen, Copenhagen, Denmark.

<sup>3</sup> Division of Epidemiology, Human Genetics and Environmental Sciences, The University of Texas Health Science Center at Houston, School of Public Health, Houston, TX, USA.

<sup>4</sup> Danish Cancer Society Research Centre, Copenhagen, Denmark.

<sup>5</sup> Department of Occupational Medicine, Danish Ramazzini Centre, University of Aarhus, Aarhus, Denmark.

Correspondence to: Helena Breth Nielsen, National Research Centre for the Working Environment, Lersø Parkallé 105, 2100 Copenhagen Ø, Denmark. [E-mail: hbn@nrcwe.dk]

shorter and impaired sleep and poorer performance (17), which in turn may affect their alertness and decision-making abilities during work activities, as well as outside work (eg, impaired driving performance). This implies that night work and rotating work patterns may also affect the risk of injury outside work as sleepiness and poor performance carry over to commuting and leisure time.

Previous studies have found an increased risk of injury among individuals who work night compared to day shifts (18–20) or shifts other than the night shift (21, 22); similar results have been observed among workers with non-regular day work compared with those with regular day work (23, 24). Furthermore, the number of consecutive night shifts (4) and night shifts in the past week have been associated with a higher risk of injury (25, 26). However, few studies have been longitudinal (18, 19, 27) and most are based on self-reported, retrospectively collected information on work patterns (18, 22–24, 26, 27) and/or injury (18, 22–24, 26, 27), introducing possible bias due to non-participation, recall and reverse causality in cross-sectional designs. Furthermore, most previous studies have assessed the effects of evening or night work by comparing risk of injury among evening or night workers with day

workers (18, 21–24, 26, 27). However, studies suggest that day, evening and night workers differ in terms of personality, lifestyle (28–30), as well as work tasks and work environment (29, 31, 32). Also, shift workers may be a highly selected group of people due to the "healthy worker effect" (29). These differences, which are unknown or rarely measured, may confound the association between work patterns and injury, as the group of exposed and the reference may not be exchangeable. Daily collection of working hours (eg, payroll data) provide a unique opportunity for longitudinal cohort studies with objective measures of the start and end times of work each day for each employee (33, 34). This daily information allows the same individual to contribute with risk time to both exposure and reference.

The purpose of this study was to (i) assess the association between working evening or night during the past seven days and risk of injury (occupational, transport and leisure time), compared to only day work during the past week; (ii) determine the association between the number of evening or night shifts worked during the past seven days and the risk of injury; and (iii) examine these associations when reducing unmeasured confounding in two restricted sub-populations of evening and night workers, respectively.


## Methods

In this longitudinal study, we used a dynamic cohort of Danish hospital and administrative workers from 2007 to 2015. By use of the Danish personal identification number (35), work patterns were linked at the individual level to information on all injuries (occupational, commuting and leisure time injury) from national health and death registers. Daily information on exposure and injury allowed us to classify the number of evening or night shifts during the past week for each day at risk for each individual.

### Total population

We included all employees in two regions of Denmark, covering urban and rural areas, registered with  $\geq 1$  shift during 2007–2015 ( $N=152\,995$ ). We excluded employees who were  $<18$  or  $>65$  years of age ( $N=1530$ ), worked part-time (annual average  $<30$  work hours/work week) ( $N=44\,607$ ), or worked  $<1$  year according to the Danish Working Hour Database (DWHD) ( $N=19\,710$ ). The total population included 69 200 participants (see figure 1) with rotating or permanent day, evening and night workers.

Employees entered the study one year after their first registration in DWHD if they were primarily employed (eg, not self-employed or student) based on source for



**Figure 1.** Flow chart of included employees (N) and numbers of person years (PY).

main income and 18 years old. They were followed until first injury, death, emigration from Denmark, 66 years of age, termination of employment or the end of follow-up, whichever came first. Participants were allowed to leave and re-enter the study population throughout follow-up. They were excluded temporarily (ie, did not contribute to risk time) when they were not primarily employed or worked <20 hours the previous week (eg, due to holiday, maternity leave and long-term sickness absence).

### Sub-populations

To improve exchangeability between exposed and unexposed, two sub-populations of evening and night workers (with no permanent day, evening or night workers) were constructed. Based upon definitions by Härmä et al (36), the evening-worker population consisted of 40 646 employees with  $\geq 12$  days with evening shifts and  $\geq 12$  days with day shifts annually, evaluated monthly (figure 1). The night-worker population included 24 515 employees with  $\geq 12$  days with night shift and  $\geq 12$  days with day shifts annually (figure 1). Employees could contribute to both the evening- and night-worker population if they met both inclusion criteria.

### Exposure assessment

Information on working hours was obtained from the DWHD, which includes payroll data, with exact time of start and end of each shift, primarily from all public hospital employees in Denmark from 2007–2015.

We defined three shifts based on the highest correlations in the study population between all actual clock hours at work: Day shift:  $\geq 3$  hours of work between 08:00–13:59 hours ( $\rho$ : 0.83–0.99); evening shift:  $\geq 3$  hours of work between 16:00–21:59 hours ( $\rho$ : 0.81–0.98); night shift:  $\geq 3$  hours of work between 00:00–05:59 hours ( $\rho$ : 0.98–1.00).

For each calendar day, we evaluated if the employee had worked day, evening or night shifts during the past seven days. We then constructed four exposure variables related to the past seven days: evening work (yes/no), night work (yes/no), number of evening shifts ( $0\text{--}\geq 5$  days with evening work), and number of night shifts ( $0\text{--}\geq 5$  days with night work). Thus, employees could have both evening and night work in the same past week. On average the evening-worker population worked 38 evening shifts and 15 night shifts per person year while the night-worker population worked 41 evening shifts and 31 night shifts per person year.

### Injury

Information on injuries was obtained from the National Patient Register (37) and the Danish Register of Causes

of Death (38) and included both injuries at work and outside of work combined (ie, occupational, commuting and leisure-time injuries). The National Patient Register contains information on all patient contacts in both public and private Danish hospitals (37). We included all emergency room contacts with "accident" as the reason for the visit (39). All deaths of residents in Denmark are registered in the Danish Register of Causes of Death (38). We included all deaths with "accident" as cause of death or where the primary or supplementary cause of death was classified as ICD10 codes: S00-T79 and T90-T99 (injury, poisoning and certain other consequences of external causes), X00-X59, V01-W99 and Y10-Y34 (external causes of morbidity and mortality). For exact codes, see the study protocol (40).

### Covariates

Information on covariates was derived from DWHD (age, sex, season, calendar year, previous sick leave, region and time since last shift) and Statistic Denmark's Employment Classification Module (occupation and socioeconomic status, SES) (41). All included covariates were time-dependent by calendar day. Age was categorized as 18–24, 25–34, 35–44, 45–54 and 55–65 years. We classified SES as high (managers and high skilled jobs), intermediate (intermediate skilled jobs), and low (low or unskilled jobs) (for exact codes see supplementary material, [www.sjweh.fi/show\\_abstract.php?abstract\\_id=3737](http://www.sjweh.fi/show_abstract.php?abstract_id=3737)). Occupation was classified as administrative work (eg, secretaries, managers and scientific workers), jobs with patient contact (eg, doctors, nurses and porters) and technical staff (eg, technicians, chefs and cleaners), based on the Danish extended version of the classification of ISCO codes (41) with information on annual main source of income (see supplementary material). Previous sick leave ( $>10$  days of sick leave within the previous year), weekly working hours (continuous), time since last shift [ $\leq 24$ , 25–72 and  $>72$  hours from end of last shift till start of calendar day (00:00)], evening work in the past week (yes/no) and night work in the past week (yes/no) were based on DWHD. We used the first year of data for all employees to calculate baseline values for previous sick leave, evening workers, night workers as well as evening and night shifts.

### Statistical method

We used Poisson regression to model incidence rate ratios (IRR) of injury as a function of evening or night work, with the log (person days) as offset. This reflects the time during which the response could occur (1), since injuries could happen over the entire day (both at and outside of work). Generalized estimating equations (GEE) with an independent correlation structure were used to account for repeated measures within employees.

IRR with 95% confidence intervals (CI) were presented in three models. In model 1, estimates were adjusted for season and calendar year. Model 2 included model 1 and age, sex and occupation. Model 3 included model 2 and previous sick leave, weekly working hours, time since last shift, SES, region and – depending on the independent variable – night work in the past week in analyses of evening work or evening work in the past week in analyses of night work. Furthermore, we performed two sensitivity analyses based on model 2. In the first, weeks with night work were excluded from the analyses of evening work and, in the second, weeks with evening work were excluded from the analyses of night work. All analyses were conducted in SAS 9.4 (SAS Institute, Cary, NC, USA) and performed on the total population as well as the two sub-populations of evening and night workers.

## Results

Characteristics of the three populations are shown in table 1. In the total population, 167 726 person years (61 219 861 person days) and 11 834 first injuries were registered. The time-in-study ranged from 1–2830 days per employee, with a mean of 885 [standard deviation (SD) 757] days, and mean age at baseline was 40.5 (SD 11.2) years. Women, employees with patient contact, and employees with an intermediate SES contributed

the majority of person years.

The restricted population of evening workers included 40 646 employees, 5834 injuries and 70 843 person years. Compared with the total population, employees with patient contact or intermediate SES contributed a larger proportion of person years. The restricted population of night workers consisted of 24 515 employees resulting in 3380 cases and 36 873 person years. Compared with the total population at baseline, the night-worker population was younger [mean age 37.1 (SD 10.2) years] and a higher proportion had occupations with patient contact.

Table 2 shows the associations between injury and evening or night work in the total population, within evening as well as night workers. In the total population, employees had higher risk of injury after a week including evening work (IRR 1.32, 95% CI 1.26–1.37), and after a week including night work (IRR 1.33, 95% CI 1.25–1.41) compared with after a week with only day work, when adjusted for confounding (model 2).

Similar, although attenuated estimates were observed for evening work in the past week among evening workers (IRR 1.18, 95% CI 1.12–1.25) and for night work among night workers (IRR 1.10, 95% CI 1.01–1.20).

In terms of number of evening and night shifts in the total population, a linear trend were found in the risk of injury across 1–≥5 evening shifts in the past week ( $P<0.001$ ). The risk of injury was highest after one night shift during the past week (IRR 1.38, 95% CI 1.28–1.50)

**Table 1.** Characteristics of the study population at baseline, cases and time in the study [PY=person years; SES=socioeconomic status].

	Total population <sup>a,b</sup>					Evening-workers population <sup>b</sup>					Night-workers population <sup>b</sup>				
	Baseline <sup>c</sup> N=69 200		Cases N=11 834	Person years <sup>d</sup> PY=167 726		Baseline <sup>c</sup> N=40 646		Cases N=5834	Person years <sup>d</sup> PY= 70 843		Baseline <sup>c</sup> N=24 515		Cases N=3380	Person years <sup>d</sup> PY=36 873	
	%	N	N	%	PY	%	N	N	%	PY	%	N	N	%	PY
Age (years)															
18–24	5.4	3699	369	1.4	2392	5.1	2083	200	1.6	1162	5.6	1383	126	1.9	700
25–34	30.5	21 084	2863	18.0	30 187	34.1	13 840	1889	24.4	17 288	43.7	10 708	1458	34.7	12 804
35–44	26.0	17 989	3054	26.2	43 960	26.0	10 573	1535	27.8	19 659	24.9	6114	901	28.5	10 524
45–54	24.9	17 230	3223	31.9	53 453	23.9	9714	1411	29.7	21 027	18.8	4597	596	23.8	8761
55–65	13.3	9198	2325	22.5	37 734	10.9	4436	799	16.5	11 708	7.0	1713	299	11.1	4084
Female gender	75.6	52 292	8655	72.7	121 918	76.5	31 101	4349	73.7	52 228	76.4	18 723	2557	74.0	27 272
Previous sick leave (>10 days)	23.0	15 947	3144	21.3	35 793	23.9	9705	1608	23.0	16 278	22.8	5600	911	23.4	8624
Occupation															
Administration	23.0	15 945	2244	25.1	42 075	12.7	5157	477	11.3	7997	3.7	897	74	2.8	1024
Patient contact	63.0	43 566	7603	61.1	102 483	76.5	31 105	4710	78.8	55 837	90.9	22 279	3128	91.6	33 784
Technical staff	14.0	9689	1987	13.8	23 168	10.8	4384	647	9.9	7009	5.5	1339	178	5.6	2065
SES															
High	28.3	19 586	3042	32.6	54 703	27.2	11 069	1412	29.2	20 710	31.1	7625	913	31.0	11 437
Intermediate	41.8	28 893	5362	44.7	75 036	49.4	20 061	3026	50.5	35 758	52.3	12 809	1925	53.5	19 732
Low	29.9	20 721	3430	22.7	37 987	23.4	9516	1396	20.3	14 375	16.7	4081	542	15.5	5704
Danish region															
Urban	62.6	43 333	7475	63.9	107 086	62.5	25 412	3668	63.0	44 609	64.4	15 780	2261	64.7	23 870
Rural	37.4	25 867	4359	36.2	60 639	37.5	15 234	2166	37.0	26 235	35.6	8735	1119	35.3	13 003

<sup>a</sup> The total population consists of both permanent and rotating day, evening and night workers.

<sup>b</sup> Median weekly working hours (p25–p75) for total population: 37.0 (30.5–37.5); evening workers 36.5 (30.0–40.0) and night workers 35.7 (30.0–40.50).

<sup>c</sup> The first registration employees had when they entered the study.

<sup>d</sup> Each employee may contribute person time in several categories throughout the study period.

and lowest after three night shifts in the past week (IRR 1.20, 95% CI 1.04–1.38). Similar patterns were found for both 1–≥5 evening shifts in evening workers (linear trend:  $P=0.005$ ) and 1–≥5 night shifts in the night workers (linear trend:  $P=0.522$ ), though estimates were lower than in the total population.

The sensitivity analyses showed no major changes in estimates (evening work during the past week within evening workers: IRR 1.19, 95% CI 1.12–1.26,  $P<0.001$ ; night work during the past week within night workers: IRR 1.10, 95% CI 0.99–1.22,  $P=0.084$ ).

## Discussion

In this large cohort study of hospital and administrative workers, a higher risk of injury was observed after evening and night work in the past week compared with only day work. Lower, but still elevated risks of injury after evening and night work persisted when we reduced group differences by restricting the population to evening and night workers, respectively. Risk of injury increased with more evening shifts in the past week.

**Table 2.** Evening and night work during the last week and IRR of injury in three models in the total population and two subpopulations [PY=person years; IRR=incidence rate ratio; CI=confidence interval].

	Cases		PY			Model 1 <sup>a</sup>			Model 2 <sup>b</sup>			Model 3 <sup>c</sup>		
	N	% (N)	IRR	95% CI	P-value	IRR	95% CI	P-value	IRR	95% CI	P-value			
<b>Total population</b>														
Evening work <sup>d1</sup>					<0.001			<0.001			<0.001			
Only day	7965	77.9 (125 020)	1			1			1					
Evening work	3195	22.1 (35 514)	1.41	1.35–1.47		1.32	1.26–1.37		1.23	1.17–1.29				
Number of evening shifts					<0.001			<0.001			<0.001			
0	8639	78.8 (132 212)	1			1			1					
1	1405	9.9 (16 540)	1.33	1.26–1.41		1.25	1.18–1.32		1.22	1.15–1.30				
2	880	5.7 (9624)	1.43	1.33–1.53		1.32	1.23–1.42		1.23	1.14–1.32				
3	504	3.1 (5249)	1.50	1.37–1.64		1.40	1.28–1.53		1.25	1.14–1.37				
4	235	1.5 (2447)	1.50	1.32–1.71		1.41	1.24–1.60		1.23	1.07–1.40				
≥5	171	0.9 (1654)	1.61	1.38–1.88		1.54	1.32–1.80		1.28	1.09–1.49				
Test for trend: 1–≥5 shifts					0.002			<0.001			0.225			
<b>Night work<sup>d2</sup></b>														
Only day	7965	89.1 (125 020)	1		<0.001	1		<0.001	1		<0.001			
Night work	1445	10.9 (15 362)	1.47	1.39–1.56		1.33	1.25–1.41		1.21	1.12–1.32				
Number of night shifts					<0.001			<0.001			<0.001			
0	10 389	90.8 (152 363)	1			1			1					
1	691	4.2 (7099)	1.53	1.41–1.65		1.38	1.28–1.50		1.37	1.22–1.53				
2	389	2.5 (4214)	1.44	1.30–1.60		1.27	1.15–1.41		1.20	1.06–1.35				
3	201	1.4 (2328)	1.35	1.17–1.55		1.20	1.04–1.38		1.08	0.93–1.25				
4	97	0.6 (1016)	1.49	1.22–1.81		1.37	1.12–1.67		1.19	0.97–1.46				
≥5	67	0.4 (706)	1.47	1.16–1.88		1.45	1.13–1.85		1.15	0.89–1.48				
Test for trend: 1–≥5 shifts					0.339			0.876			0.294			
<b>Evening worker-population</b>														
Evening work <sup>d3</sup>					<0.001			<0.001			<0.001			
Only day	2479	52.1 (34 045)	1			1			1					
Evening work	2802	47.9 (31 252)	1.23	1.16–1.30		1.18	1.12–1.25		1.13	1.07–1.20				
Number of evening shifts					<0.001			<0.001			<0.001			
0	3032	55.9 (39 592)	1			1			1					
1	1303	21.8 (15 405)	1.16	1.09–1.24		1.13	1.06–1.21		1.12	1.04–1.21				
2	818	12.6 (8914)	1.26	1.16–1.36		1.20	1.11–1.30		1.14	1.04–1.24				
3	426	6.2 (4392)	1.33	1.20–1.47		1.26	1.14–1.40		1.16	1.04–1.29				
4	174	2.5 (1761)	1.35	1.16–1.58		1.28	1.10–1.50		1.15	0.98–1.35				
≥5	81	1.2 (779)	1.42	1.14–1.77		1.37	1.10–1.71		1.17	0.94–1.47				
Test for trend: 1–≥5 shifts					0.002			0.005			0.424			
<b>Night-worker population</b>														
Night work <sup>d4</sup>					0.003			0.021			0.498			
Only day	1053	49.5 (12 814)	1											
Night work	1219	50.5 (13 075)	1.13	1.04–1.23		1.10	1.01–1.20		1.04	0.93–1.15				
Number of night shifts					0.089			0.214			0.521			
0	2161	64.5 (23 799)	1			1			1					
1	603	17.3 (6359)	1.16	1.05–1.28		1.14	1.03–1.26		1.12	0.98–1.28				
2	352	10.3 (3784)	1.13	1.00–1.27		1.08	0.95–1.21		1.02	0.88–1.17				
3	171	5.2 (1930)	1.07	0.91–1.26		1.02	0.87–1.21		0.95	0.80–1.12				
4	64	1.9 (698)	1.11	0.86–1.43		1.09	0.84–1.40		1.00	0.78–1.30				
≥5	29	0.8 (303)	1.15	0.79–1.67		1.21	0.83–1.76		1.08	0.73–1.59				
Test for trend: 1–≥5 shifts					0.508			0.522			0.256			

<sup>a</sup> Model 1: adjusted for calendar year and season.

<sup>b</sup> Model 2: adjusted for model 1 + age, sex and occupation.

<sup>c</sup> Model 3: adjusted for model 2 + previous sick leave, weekly working hours, time since last shift, SES, region and evening or night work past week.

<sup>d</sup> Missing PY (not in evening/night work or only day work). D1: 7192, D2: 27 344; D3 5547; D4 10 985.

For night shifts, the risk of injury was lowest after three night shifts in the past week. Excluding days with night and evening work during the past week within analyses of evening and night work, respectively, did not change the conclusions.

With all workers included, our results of a higher risk of injury after evening and night work during the past week compared with day work, are generally in line with previous studies (19–22). Another study using payroll information found a higher risk of occupational injuries in afternoon shifts (IRR 1.66, 95% CI 1.23–2.25) and midnight shifts (IRR 1.72, 95% CI 1.26–2.36) compared with day shifts among US police officers (19). Where previous studies have mainly focused on occupational injuries (18, 19, 22–24), we analyzed all injuries. Our results on night work are consistent with a recent study among randomly selected Danish employees that used all injuries as the outcome [relative risk (RR) 1.11, 95% CI 1.06–1.17] (21).

A newly published systematic review found no linear risk of occupational injuries across number of consecutive afternoon/evening shifts based on three studies with only 2062 cases (4). Based on 5834 cases, we found an increasing risk of injury across the number of evening shifts in the past week. In model 3, no linear trend was observed. However, we suspect the model includes possible mediators and may be over adjusted.

We observed the lowest risk of injury after three night shifts in the past week, whereas previous cross-sectional studies have found a higher risk after three or more night shifts compared with none (25, 26). This deviation may be explained by a cruder categorization of night shifts and different confounding adjustment in previous studies. The present indication of a U-shaped risk could be a result of a combination of two different mechanisms. Sleepiness is most pronounced after the first and last (consecutive) night shifts (42), and thus the relative contribution of sleepiness is larger with few night shifts. On the other hand, sleep debt accumulates with increasing number of night shifts. Accordingly, sleepiness and therefore risk of injury is expected to be higher with more night shifts. When night work is unavoidable, our results indicate that three night shifts during the past week are preferable. However, more studies are needed to verify our results.

Most previous studies compare evening or night workers with day workers (18, 21–24, 26, 27). Even when results are adjusted for covariates, such as age, SES and weekly working hours, group differences between day, evening and night workers may still persist and lead to biased results. In our analysis of evening and night workers, employees could contribute with both exposed and unexposed risk time. This improved the exchangeability and reduced potential unmeasured confounding. The lower relative risk of injury follow-

ing evening and night work within evening and night workers compared with the total population, suggests additional unmeasured confounding not accounted for by the confounding adjustment.

A previous, well-designed study, which also took group differences into account, found a higher risk of occupational injuries during afternoon shifts (RR 1.15, 95% CI 1.07–1.23), and during night shifts (RR 1.23, 95% CI 1.14–1.31) compared with morning shifts, when studying the shifts during which the injuries occurred (43). These findings are consistent with our results on evening work within evening workers and night work within night workers.

### Strengths and limitations

The strengths of this study include the large study population, longitudinal design and the objective measures of both exposure and outcomes from reliable nationwide registers. This eliminates all recall bias. Detailed information from payroll data allowed us to calculate daily time-varying exposures thereby reducing misclassification. Moreover, by using a one year wash-out period prior to entry and only the first injury, we reduced the risk of reverse causation due to previous injuries, which may alter work patterns afterwards (44). All injuries (occupational, commuting and leisure time injuries) were included from national emergency room and death registers reducing the risk of bias from underreporting (45, 46). Lastly, little misclassification of timing of work in DWHD and shift categorizing is expected as these were based on actual shift schedules, largely covering definition from previous studies (19, 25, 34).

Unmeasured confounding was reduced by improving the exchangeability between exposed and unexposed. This may also reduce bias from self-selection into shift work. In addition, we adjusted for many potential confounders including season, sex and occupation. We also controlled for short job tenure and part-time work by excluding these from the study population. Working hours from multiple jobs within the regions were combined, but we had no information on jobs outside of the regions. However, by only including fulltime workers, we expect to have adjusted for most of the confounding from multiple jobs. Yet, task, workload, work environment, leisure-time activities, and traffic may be additional potential confounders if associated with shift patterns. Most of these differences are expected to decrease in the restricted populations, due to exclusion of, for example, daytime working administrative employees. Nonetheless, confounding from unmeasured individual differences may persist, since employees did not contribute with an equal amount of time to the exposure and reference groups and our results need to be confirmed in future studies. Furthermore, different characteristics

of work patterns are highly correlated and the effects are therefore difficult to disentangle. Especially long shifts may have contributed to the higher risk of injury after evening work during the past week. Moreover, rest breaks, on-call shifts, quick-returns and rotating work may be of interest to consider in future studies.

No information on exact time of the injury was available, thus there is a risk of assessing work patterns after the injury. However, injuries are only admitted to emergency rooms in Denmark within 24 hours after occurrence. By assessing the exposures up until the day before the injury the risk of reverse causation was reduced.

The populations of evening and night workers did not include permanent evening or night workers. Whether risks are similar in permanent evening and night workers remains to be settled. Thus, the results are generalizable to workers with some day work in Denmark and should be interpreted with caution in other countries and with respect to job groups with similar work patterns. Furthermore, we expect the mechanisms to be similar for minor injuries.

In conclusion, we find an increased risk of injury after evening and night work among Danish hospital employees compared with day work during the past week. The higher risks persist, though attenuated, when unmeasured confounding from differences between day, evening and night workers was reduced. While a dose-response for number of evening shifts was found, in our study, three night shifts during the past week seem to be preferable in relation to risk of injury.

### Acknowledgements

The authors have no conflicts of interest to declare.

The Danish Working Environment Research Fund supported this study [grant number 31-2014-09/20140072847] but was not involved in the design, execution, writing or decision to publish this study. The Danish National Data Protection Agency approved the study (J.nr. 2015-57-0074).

The DWHD was established in collaboration between National Research Centre for the Working Environment, the Danish Cancer Society, Danish Ramazzini Centre, Department of Occupational Medicine, Aarhus University Hospital, and Department of Public Health, University of Copenhagen. The establishment of the DWHD was financed by research grants from the Danish Working Environment Research Fund (23-2012-09), NordForsk, Nordic Program on Health and Welfare (74809), and the National Research Centre for the Working Environment. The regions have partially financed the transfer of data to the cohort.

We would like to acknowledge the Danish regions for their participation and willingness to provide data to the DWHD. Moreover, we thank Jens Worm Begtrup, Anders Ørberg and Lisbeth Nielsen from The National Research Centre for the Working Environment for valuable work with data management in establishing the DWHD.

### References

1. Zhao I, Bogossian F, Turner C. Shift work and work related injuries among health care workers: A systematic review. *Aust J Adv Nurs* 2010;27(3):62–74.
2. Folkard S, Lombardi DA. Modeling the impact of the components of long work hours on injuries and “accidents”. *Am J Ind Med* 2006 Nov;49(11):953–63. <http://dx.doi.org/10.1002/ajim.20307>.
3. Wagstaff AS, Sigstad Lie JA. Shift and night work and long working hours—a systematic review of safety implications. *Scand J Work Environ Health* 2011 May;37(3):173–85. <http://dx.doi.org/10.5271/sjweh.3146>.
4. Fischer D, Lombardi DA, Folkard S, Willetts J, Christiani DC. Updating the “Risk Index”: A systematic review and meta-analysis of occupational injuries and work schedule characteristics. *Chronobiol Int* 2017;34(10):1423–38. <http://dx.doi.org/10.1080/07420528.2017.1367305>.
5. The national research centre for The working environment. [Working Environment in Denmark] Arbejdsmiljø i Danmark [Internet]. 2017 [cited 2017 Jun 1]. Available from: <https://arbejdsmiljoindanmark.nfa.dk/>.
6. Labour Market Insurance. Arbejdsskade-statistik 2016 [Occupational injury statistics 2016] [Internet]. Copenhagen; 2017 [cited 2017 Oct 30]. Available from: [https://aes.dk/da/Presse-og-nyheder/Nyhedsarkiv/2017/7\\_april\\_Arbejdsskade-statistik\\_2016.aspx](https://aes.dk/da/Presse-og-nyheder/Nyhedsarkiv/2017/7_april_Arbejdsskade-statistik_2016.aspx).
7. Møller H, Damm M, Laursen B. Ulykker i Danmark 1990-2009 [Accidents in Denmark 1990-2009]. Kbh.: Statens Institut for Folkesundhed; 2012.
8. O’Donnell ML, Varker T, Holmes AC, Ellen S, Wade D, Creamer M et al. Disability after injury: the cumulative burden of physical and mental health. *J Clin Psychiatry* 2013 Feb;74(2):e137–43. <http://dx.doi.org/10.4088/JCP.12m08011>.
9. Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012 Dec;380(9859):2197–223. [http://dx.doi.org/10.1016/S0140-6736\(12\)61689-4](http://dx.doi.org/10.1016/S0140-6736(12)61689-4).
10. Meerding WJ, Looman CW, Essink-Bot ML, Toet H, Mulder S, van Beeck EF. Distribution and determinants of health and work status in a comprehensive population of health and work status in a comprehensive population of injury patients. *J Trauma* 2004 Jan;56(1):150–61. <http://dx.doi.org/10.1097/01.TA.0000062969.65847.8B>.

11. Toft AM, Møller H, Laursen B. The years after an injury: long-term consequences of injury on self-rated health. *J Trauma* 2010 Jul;69(1):26–30. <http://dx.doi.org/10.1097/TA.0b013e3181d3cbf2>.
12. Williamson A, Lombardi DA, Folkard S, Stutts J, Courtney TK, Connor JL. The link between fatigue and safety. *Accid Anal Prev* 2011 Mar;43(2):498–515. <http://dx.doi.org/10.1016/j.aap.2009.11.011>.
13. Robb G, Sultana S, Ameratunga S, Jackson R. A systematic review of epidemiological studies investigating risk factors for work-related road traffic crashes and injuries. *Inj Prev* 2008 Feb;14(1):51–8. <http://dx.doi.org/10.1136/ip.2007.016766>.
14. Zhang T, Chan AH. Sleepiness and the risk of road accidents for professional drivers: A systematic review and meta-analysis of retrospective studies. *Saf Sci* 2014;70:180–8. <http://dx.doi.org/10.1016/j.ssci.2014.05.022>.
15. Åkerstedt T. Work hours, sleepiness and accidents Introduction and summary. *J Sleep Res* 1995 Dec;4(2):1–3. <http://dx.doi.org/10.1111/j.1365-2869.1995.tb00219.x>.
16. Uehli K, Mehta AJ, Miedinger D, Hug K, Schindler C, Holsboer-Trachsler E et al. Sleep problems and work injuries: a systematic review and meta-analysis. *Sleep Med Rev* 2014 Feb;18(1):61–73. <http://dx.doi.org/10.1016/j.smrv.2013.01.004>.
17. Rajaratnam SM, Arendt J. Health in a 24-h society. *Lancet* 2001 Sep;358(9286):999–1005. [http://dx.doi.org/10.1016/S0140-6736\(01\)06108-6](http://dx.doi.org/10.1016/S0140-6736(01)06108-6).
18. Dembe AE, Erickson JB, Delbos RG, Banks SM. Nonstandard shift schedules and the risk of job-related injuries. *Scand J Work Environ Health* 2006 Jun;32(3):232–40. <http://dx.doi.org/10.5271/sjweh.1004>.
19. Violanti JM, Fekedulegn D, Andrew ME, Charles LE, Hartley TA, Vila B et al. Shift work and the incidence of injury among police officers. *Am J Ind Med* 2012 Mar;55(3):217–27. <http://dx.doi.org/10.1002/ajim.22007>.
20. Folkard S, Akerstedt T. Trends in the risk of accidents and injuries and their implications for models of fatigue and performance. *Aviat Space Environ Med* 2004 Mar;75(3 Suppl):A161–7.
21. Larsen AD, Hannerz H, Møller SV, Dyreborg J, Bonde JP, Hansen J et al. Night work, long work weeks, and risk of accidental injuries. A register-based study. *Scand J Work Environ Health* 2017 Nov;43(6):578–86.
22. Stimpfel AW, Brewer CS, Kovner CT. Scheduling and shift work characteristics associated with risk for occupational injury in newly licensed registered nurses: an observational study. *Int J Nurs Stud* 2015 Nov;52(11):1686–93. <http://dx.doi.org/10.1016/j.ijnurstu.2015.06.011>.
23. de Castro AB, Fujishiro K, Rue T, Tagalog EA, Samaco-Paquiz LP, Gee GC. Associations between work schedule characteristics and occupational injury and illness. *Int Nurs Rev* 2010 Jun;57(2):188–94. <http://dx.doi.org/10.1111/j.1466-7657.2009.00793.x>.
24. Trinkoff AM, Le R, Geiger-Brown J, Lipscomb J. Work schedule, needle use, and needlestick injuries among registered nurses. *Infect Control Hosp Epidemiol* 2007 Feb;28(2):156–64. <http://dx.doi.org/10.1086/510785>.
25. Hopcia K, Dennerlein JT, Hashimoto D, Orechia T, Sorensen G. Occupational injuries for consecutive and cumulative shifts among hospital registered nurses and patient care associates: a case-control study. *Workplace Health Saf* 2012 Oct;60(10):437–44.
26. Fransén M, Wilsmore B, Winstanley J, Woodward M, Grunstein R, Ameratunga S et al. Shift work and work injury in the New Zealand Blood Donors' Health Study. *Occup Environ Med* 2006 May;63(5):352–8. <http://dx.doi.org/10.1136/oem.2005.024398>.
27. Dembe AE, Delbos R, Erickson JB. Estimates of injury risks for healthcare personnel working night shifts and long hours. *Qual Saf Health Care* 2009 Oct;18(5):336–40. <http://dx.doi.org/10.1136/qshc.2008.029512>.
28. Knutsson A, Nilsson T. Tobacco use and exposure to environmental tobacco smoke in relation to certain work characteristics. *Scand J Soc Med* 1998 Sep;26(3):183–9. <http://dx.doi.org/10.1177/14034948980260030801>.
29. Knutsson A. Methodological aspects of shift-work research. *Chronobiol Int* 2004;21(6):1037–47. <http://dx.doi.org/10.1081/CBI-200038525>.
30. Nabe-Nielsen K, Garde AH, Tüchsen F, Hogh A, Diderichsen F. Cardiovascular risk factors and primary selection into shift work. *Scand J Work Environ Health* 2008 Jun;34(3):206–12. <http://dx.doi.org/10.5271/sjweh.1230>.
31. Nabe-Nielsen K, Tüchsen F, KB C, AH G, Diderichsen F. Differences between day and non-day workers in exposure to physical and psychosocial work factors in the Danish eldercare sector. *Scand J Work Environ Health* 2009;35(1):48–55.
32. Bøggild H, Burr H, Tüchsen F, Jeppesen HJ. Work environment of Danish shift and day workers. *Scand J Work Environ Health* 2001 Apr;27(2):97–105. <http://dx.doi.org/10.5271/sjweh.595>.
33. Härmä M, Ropponen A, Hakola T, Koskinen A, Vanttola P, Puttonen S et al. Developing register-based measures for assessment of working time patterns for epidemiologic studies. *Scand J Work Environ Health* 2015 May;41(3):268–79. <http://dx.doi.org/10.5271/sjweh.3492>.
34. Garde AH, Hansen J, Kolstad HA, Larsen AD, Hansen ÅM. How do different definitions of night shift affect the exposure assessment of night work? *Chronobiol Int* 2016;33(6):595–8. <http://dx.doi.org/10.3109/07420528.2016.1167729>.
35. Pedersen CB. The Danish Civil Registration System. *Scand J Public Health* 2011 Jul;39(7 Suppl):22–5. <http://dx.doi.org/10.1177/1403494810387965>.
36. Härmä M, Koskinen A, Ropponen A, Puttonen S, Karhula K, Vahtera J et al. Validity of self-reported exposure to shift work. *Occup Environ Med* 2017 Mar;74(3):228–30. <http://dx.doi.org/10.1136/oemed-2016-103902>.
37. Lynge E, Sandegaard JL, Rebolj M. The Danish National

- Patient Register. *Scand J Public Health* 2011 Jul;39(7 Suppl):30–3. <http://dx.doi.org/10.1177/1403494811401482>.
38. Helweg-Larsen K. The Danish Register of Causes of Death. *Scand J Public Health* 2011 Jul;39(7 Suppl):26–9. <http://dx.doi.org/10.1177/1403494811399958>.
  39. Sundhedsdatastyrelsen. Fællesindhold for basisregistrering af sygehuspatienter [Join content for basic registration of hospital patients] [Internet]. Copenhagen: Sundhedsdatastyrelsen; 2017 [cited 2017 Oct 26]. Report No.: 6. Available from: <https://sundhedsdatastyrelsen.dk/da/rammer-og-retningslinjer/om-patientregistrering/patientregistrering-feallesindhold>.
  40. Nielsen HB, Larsen AD, Dyreborg J, Hansen ÅM, Hansen J, HenrikKolstad, et al. Working hours and the risk of injuries – study protocol [Internet]. 2017 [cited 2017 Oct 26]. Available from: [https://figshare.com/articles/Working\\_hours\\_and\\_the\\_risk\\_of\\_injuries\\_study\\_protocol/4801891](https://figshare.com/articles/Working_hours_and_the_risk_of_injuries_study_protocol/4801891).
  41. Petersson F, Baadsgaard M, Thygesen LC. Danish registers on personal labour market affiliation. *Scand J Public Health* 2011 Jul;39(7 Suppl):95–8. <http://dx.doi.org/10.1177/1403494811408483>.
  42. Waage S, Harris A, Pallesen S, Saksvik IB, Moen BE, Bjorvatn B. Subjective and objective sleepiness among oil rig workers during three different shift schedules. *Sleep Med* 2012 Jan;13(1):64–72. <http://dx.doi.org/10.1016/j.sleep.2011.04.009>.
  43. Smith L, Folkard S, Poole CJ. Increased injuries on night shift. *Lancet* 1994 Oct;344(8930):1137–9. [http://dx.doi.org/10.1016/S0140-6736\(94\)90636-X](http://dx.doi.org/10.1016/S0140-6736(94)90636-X)
  44. Matérne M, Lundqvist L-O, Strandberg T. Opportunities and barriers for successful return to work after acquired brain injury: A patient perspective. *Work Read Mass*. 56(1):125–34.
  45. Brown JG, Trinkoff A, Rempher K, McPhaul K, Brady B, Lipscomb J et al. Nurses' inclination to report work-related injuries: organizational, work-group, and individual factors associated with reporting. *AAOHN J* 2005 May;53(5):213–7.
  46. Wuellner SE, Bonauto DK. Exploring the relationship between employer recordkeeping and underreporting in the BLS Survey of Occupational Injuries and Illnesses. *Am J Ind Med* 2014 Oct;57(10):1133–43. <http://dx.doi.org/10.1002/ajim.22350>.

Received for publication: 7 March 2018