



## Short communication

Scand J Work Environ Health 2019;45(2):203-208

doi:10.5271/sjweh.3780

### Early part-time sick leave results in considerable savings in social security costs at national level: an analysis based on a quasi-experiment in Finland

by Viikari-Juntura E, Leinonen T, Virta LJ, Hiljanen I, Husgafvel-Pursiainen K, Autti-Rämö I, Rissanen P, Burdorf A, Solovieva S

Use of part-time sick leave instead of full-time sick leave at the early stage of work disability due to a musculoskeletal disease or mental disorder leads to both gain in work participation, extension of working life by avoiding full retirement, and social security cost savings during a two year period. Part-time sick leave can be recommended from an economic perspective.

**Affiliation:** Finnish Institute of Occupational Health. PO Box 40, FI-00032 Työterveyslaitos, Finland. eira.viikari-juntura@ttl.fi

Refers to the following texts of the Journal: 2008;34(4):239-249  
2012;38(2):134-143 2017;43(5):447-456

The following article refers to this text: [online first; 12 April 2019]

**Key terms:** cost saving; disability; disability cost; Finland; mental disorder; musculoskeletal disease; musculoskeletal disorder; part-time sick leave; quasi-experiment; sick leave; sick leave cost; sickness absence; sickness absence cost; sickness benefit; social security; social security cost

This article in PubMed: [www.ncbi.nlm.nih.gov/pubmed/30338336](http://www.ncbi.nlm.nih.gov/pubmed/30338336)

### Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](http://www.sjweh.fi).



This work is licensed under a [Creative Commons Attribution 4.0 International License](http://creativecommons.org/licenses/by/4.0/).

## Early part-time sick leave results in considerable savings in social security costs at national level: an analysis based on a quasi-experiment in Finland

by Eira Viikari-Juntura, MD, PhD,<sup>1</sup> Taina Leinonen, PhD,<sup>1</sup> Lauri J Virta, MD, PhD,<sup>2</sup> Ismo Hiljanen, MSc,<sup>2</sup> Kirsti Husgafvel-Pursiainen, PhD,<sup>1</sup> Ilona Autti-Rämö, MD, PhD,<sup>2</sup> Pekka Rissanen, PhD,<sup>3</sup> Alex Burdorf, PhD,<sup>4</sup> Svetlana Solovieva, PhD<sup>1</sup>

Viikari-Juntura E, Leinonen T, Virta LJ, Hiljanen I, Husgafvel-Pursiainen K, Rissanen P, Burdorf A, Solovieva S. Early part-time sick leave results in considerable savings in social security costs at national level: an analysis based on a quasi-experiment in Finland. *Scand J Work Environ Health*. 2019;45(2):203–208. doi:10.5271/sjweh.3780

**Objectives** We analyzed social security costs based on an earlier quasi-experiment that compared work participation between partial sickness beneficiaries and a matched group of full sickness beneficiaries.

**Methods** Utilizing a population-based 70% representative sample, 1878 persons with part-time sick leave (intervention group) due to musculoskeletal diseases or mental disorders at an early stage of work disability and their propensity-score-matched controls with full-time sick leave were followed for two years. The outcome was the difference (absolute and relative) in social security costs between the intervention and control groups during follow-up. Costs of sickness absence, vocational rehabilitation, unemployment, and retirement days were calculated from national administrative registers.

**Results** A cost reduction of €2395 per person per year [95% confidence interval (CI) -2890–-1899] was observed in the intervention compared with the control group. The cost ratio was 0.512 (95% CI 0.511–0.513). The largest savings were attributable to differences in the costs of retirement and vocational rehabilitation. The savings were higher for the second compared with the first follow-up year. Costs were saved across both genders and diagnostic groups, however, savings for women with musculoskeletal diseases were lowest.

**Conclusions** Part-time instead of full-time sick leave, at the early stage of work disability due to musculoskeletal diseases or mental disorders, leads to considerable social security cost savings during two years, in correspondence with increased work participation and in addition to earlier reported health benefits. Part-time sick leave can be recommended from an economic perspective; however more consideration should be given to women with musculoskeletal diseases.

**Key terms** cost saving; disability; disability cost; mental disorder; musculoskeletal disease; musculoskeletal disorder; sickness benefit; sickness absence; sickness absence cost; sick leave cost.

Part-time sick leave is used in all Nordic and some European countries to enable continued working when health reasons prevent full-time work (1, 2). Part-time return to work (RTW) supported by partial sickness benefit has in many studies resulted in enhanced return to full-time work and reduced sickness absence (SA) in the long term (3–8). We are not aware of studies looking at the economic aspects of part-time sick leave.

The legislative framework for the use of part-time sick leave differs between countries. In most Nordic countries, part-time sick leave is used to compensate

for the proportional loss of work ability, and the sickness benefit will, respectively, range from a few to close to 100 percent of full sickness benefit. In Finland, however, part-time sick leave is a voluntary option for those who have been assessed by a physician to be incapable of performing their full duties, but are able to work 40–60% of full time. The partial sickness benefit is always set at 50% of full benefit. The application for the benefit is made jointly by the employer and the employee and granted by the Social Insurance Institution (SII). Part-time sick leave was introduced in Finland as

<sup>1</sup> Finnish Institute of Occupational Health, Helsinki, Finland.

<sup>2</sup> The Social Insurance Institution of Finland, Helsinki, Finland.

<sup>3</sup> National Institute for Health and Welfare, Helsinki, Finland.

<sup>4</sup> Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands.

late as 2007, and in 2010 the criteria for its use were widened.

Earlier we carried out a quasi-experiment looking at the effectiveness of the use of part-time sick leave at the early stage of work disability due to a musculoskeletal disease or mental disorder. During the two-year follow-up, the proportion of time at work was higher and sick leave and disability retirement lower among those who had had part-time sick leave compared with those on full-time sick leave. In the former group, the proportion of those with full disability retirement was about a third of those in the latter group (9).

In the current paper, we compare the accumulation of social security costs during two years between recipients of partial and full sickness benefit.

## Methods

We utilized a 70% random sample of the working age population in Finland on 31 December 2010. Data on compensated SA, with dates and diagnoses, were obtained from the SII. The Finnish Centre for Pensions (FCP) provided information on disability and other retirement, employment, and unemployment periods. Information on vocational rehabilitation periods was obtained from both SII and FCP. Demographic information and information of annual earnings, unemployment benefits and pension benefits were provided by Statistics Finland. Data were linked with social security numbers and provided to the researchers in unidentifiable form.

In Finland, the SII compensates SA after a so-called employer period of ten full SA days. Before 2010, part-time sick leave was available only for persons with a preceding full SA of 60 compensated days. The legislation was changed in 2010, enabling the use of part-time sick leave already from the first compensated day. Still, about 80% of recipients of the partial sick leave benefit at an early stage have a preceding compensated full-time sick leave period. We therefore chose the participants among those with a preceding full SA. Because about 70% of part-time sickness absences are prescribed due to a musculoskeletal disease or mental disorder (10), we further limited our study population to these two disease groups.

As described in our previous paper (9), there were a total of 1879 persons, who during a two-year recruitment period (1 January 2010–31 December 2011) had part-time sick leave due to a musculoskeletal disease or mental disorder, immediately following a continuous full-time SA of 1–59 compensated days. For each case with part-time sick leave, a control was selected among those who had a full-time sickness absence period continued with another full-time sickness absence. Propensity score matching was carried out within the four

diagnostic group  $\times$  gender strata to make the part-time and full-time groups comparable regarding sociodemographic factors (age, major region, employment sector, industrial sector, socioeconomic status, annual gross income in the year preceding the SA, number of days being employed or unemployed during the preceding two years) and the number of preceding SA days. This resulted in 1878 persons with part-time sick leave matched with 1878 controls with full-time sick leave. All participants were followed for two years. The complete register information enabled us to assess the “work participation status” for each participant for each follow-up day. Overall, we considered seven work participation statuses: (i) at work, (ii) full-time SA, (iii) partial work disability (including part-time sickness absence and partial disability retirement), (iv) receiving vocational rehabilitation benefit, (v) unemployed, (vi) full disability retirement, and (vii) other (outside labor force, status not known). Persons with partial work disability were considered to be working 50% of their working time.

The primary outcome for the present study was the difference in social security costs between the part-time (intervention) and full-time (control) sick leave group, expressed in absolute and relative values. We looked at the distribution of the costs of the different types of the benefits separately for the first and second follow-up year. We used percentile bootstrapping method with 5000 intervention–control pairs to estimate 95% confidence intervals (CI) for the relative difference in the costs (10). Due to a skewed distribution and a wide range of the costs, we also used the bootstrapping method to estimate the 95% CI for the difference in the mean costs per person per year.

The costs of the examined social security benefits were calculated as follows: the cost of a *compensated full-time SA day* was assessed for each person based on the person’s earned income total in state taxation for the preceding year in question using year-specific formulae provided by the SII (11–14). The cost of a *compensated part-time SA day* was calculated as 50% of a full-time sickness absence day. The cost of *vocational rehabilitation benefit day* was estimated as annual mean cost paid by the Finnish Centre for Pensions (15–18). The cost of an *unemployment day* was calculated based on the person’s unemployment income and number of unemployment days for the year in question. The cost of a *day on retirement* (including work disability, old-age, and other) was calculated based on the person’s pension income for the year in question (total pension income from all types of pensions for each group member, provided by Statistics Finland).

We looked at the cost difference in the entire study group and for the two diagnostic groups by gender. We further stratified by age group and selected industry. A preliminary scrutiny identified a few persons with very

high unemployment or retirement costs (one in part- and five in full-time sick leave group). We carried out a sensitivity analysis of the total costs where we removed these six persons and their matched pairs.

## Results

During the two-year follow-up, the number of full-time SA days was lower and both the part-time SA days and the overall SA costs were higher in the intervention than the control group (table 1). In total, about €9 million less social security costs were observed in the intervention compared with the control group, resulting in a cost ratio of 0.512 (95% CI 0.511–0.513) and a cost difference of -€2395 (95% CI -2890– -1899) per person per year. Apart from SA, the highest costs in both groups were due to retirement and vocational rehabilitation, however,

the absolute difference in costs was highest for retirement and the relative difference highest for vocational rehabilitation.

Costs were saved by the intervention in both diagnostic groups among both men and women, however, cost savings were lower for women compared with men, especially among those with musculoskeletal diseases (table 1). The largest relative savings emanated from vocational rehabilitation costs in each matched gender × diagnosis group, except men with a musculoskeletal disease, for whom the reduction in unemployment costs was the largest.

Social security costs were higher during the first compared with the second follow-up year in both intervention and control group (table 1). The share of full-time and part-time SA costs combined was the highest during the first follow-up year (23.8% + 26.2% in intervention and 23.8% + 0.9% in control group). During the second follow-up year, the share of SA costs continued

**Table 1.** Total number of days on ill-health-related, unemployment and retirement benefits, their social security costs (€) and relative cost differences (cost ratio) with 95% confidence intervals (CI) during two years of follow-up in the intervention (N=1878) and control group (N=1878). Cost distribution (%) of the benefits during the first and second follow-up year in the intervention and control group. [SA=sickness absence.]

	Number of pairs	Total number of days		Total costs (€)		Cost ratio (95% CI) <sup>a</sup>	Cost distribution (%) <sup>b</sup>		Cost distribution (%) <sup>b</sup>	
		Intervention	Control	Intervention	Control		Intervention		Control	
							1 <sup>st</sup> year	2 <sup>nd</sup> year	1 <sup>st</sup> year	2 <sup>nd</sup> year
All	1878									
Full-time SA		67 148	94 061	4 253 626	5 541 539	0.768 (0.767–0.769)	23.8	15.2	23.8	6.5
Part-time SA		61 897	1718	2 176 746	62 901	35.94 (35.74–36.15)	26.2	0.3	0.9	0.2
Vocational rehabilitation		9626	65 569	688 098	4 478 926	0.154 (0.153–0.155)	0.6	4.8	9.1	12.5
Unemployed		25 137	42 853	494 567	1 803 257	0.275 (0.274–0.279)	1.9	8.7	5.3	7.7
Retired <sup>c</sup>		62 651	133 778	1 753 386	6 474 449	0.272 (0.271–0.273)	4.0	14.3	13.9	20.0
Total				9 366 423	18 361 072	0.512 (0.511–0.513)	<b>56.6</b>	<b>43.4</b>	<b>53.1</b>	<b>46.9</b>
Musculoskeletal diseases (men)	233									
Full-time SA		7922	11 717	537 460	744 026	0.725 (0.722–0.727)	21.8	16.6	22.2	6.8
Part-time SA		6645	110	242 542	5113	50.75 (49.65–51.85)	25.0	0.0	0.1	0.2
Vocational rehabilitation		1141	6526	96 995	540 814	0.199 (0.194–0.203)	1.1	5.0	8.2	11.4
Unemployed		3384	6362	40 309	679 220	0.125 (0.119–0.130)	1.6	9.1	7.5	5.7
Retired <sup>c</sup>		8958	15 813	291 192	825 464	0.367 (0.364–0.370)	4.0	15.6	14.0	24.0
Total				1 208 498	2 794 637	0.444 (0.441–0.446)	<b>53.6</b>	<b>46.4</b>	<b>51.9</b>	<b>48.1</b>
Musculoskeletal diseases (women)	793									
Full-time SA		29 902	40 421	1 761 137	2 303 463	0.764 (0.763–0.766)	26.1	15.2	27.9	8.3
Part-time SA		26 651	619	876 106	20 745	46.98 (46.48–47.48)	24.2	0.3	0.9	0.3
Vocational rehabilitation		2756	19 921	179 930	1 274 655	0.146 (0.145–0.148)	0.2	3.2	7.7	9.8
Unemployed		9049	13 978	297 344	428 834	0.875 (0.848–0.902)	2.3	8.9	4.8	6.9
Retired <sup>c</sup>		30 425	51 629	691 045	2 171 120	0.323 (0.321–0.324)	4.3	15.5	13.0	20.5
Total				3 805 562	6 198 817	0.618 (0.616–0.620)	<b>57.0</b>	<b>43.0</b>	<b>54.2</b>	<b>45.8</b>
Mental disorders (men)	188									
Full-time SA		7087	8954	520 247	626 214	0.840 (0.837–0.844)	19.4	16.5	19.2	5.5
Part-time SA		5623	63	238 544	3486	67.83 (66.53–69.12)	24.7	0.1	0.7	0.1
Vocational rehabilitation		1921	8898	164 108	734 829	0.235 (0.231–0.238)	1.3	9.4	11.5	14.9
Unemployed		3949	6591	48 490	184 902	0.335 (0.328–0.342)	1.7	8.3	2.9	11.2
Retired <sup>c</sup>		5231	14 736	248 089	967 552	0.269 (0.266–0.272)	4.2	14.2	15.7	18.4
Total				1 219 478	2 516 983	0.491 (0.488–0.493)	<b>51.4</b>	<b>48.6</b>	<b>50.0</b>	<b>50.0</b>
Mental disorders (women)	664									
Full-time SA		22 237	32 969	1 434 782	1 867 836	0.770 (0.768–0.771)	21.8	16.6	22.2	6.8
Part-time SA		22 972	926	819 554	33 557	26.46 (26.22–26.70)	25.0	0.0	0.1	0.2
Vocational rehabilitation		3809	30 224	247 065	1 928 628	0.129 (0.128–0.130)	1.1	5.0	8.2	11.4
Unemployed		8755	15 922	108 424	510 301	0.185 (0.182–0.188)	1.6	9.1	7.5	5.7
Retired <sup>c</sup>		18 038	51 600	523 060	2 510 613	0.212 (0.211–0.213)	4.0	15.6	14.0	24.0
Total				3 132 885	6 850 635	0.459 (0.458–0.460)	<b>53.6</b>	<b>46.4</b>	<b>51.9</b>	<b>48.1</b>

<sup>a</sup> Cost ratio for the total costs for the first year of follow-up 0.545 (95% CI 0.544–0.546) and for the second year 0.473 (95% CI 0.472–0.474).

<sup>b</sup> Areas with italics and bolded italics each add up to 100%.

<sup>c</sup> Some persons were on part-time disability retirement.


to be the highest in the intervention group, whereas the share of retirement costs became predominant in the control group. Vocational rehabilitation contributed considerably less to the costs in the intervention than the control group both during the first and second follow-up year. Also unemployment contributed less to the costs in the intervention compared with the control group during the first follow-up year. Overall, relative savings were higher for the second compared with the first year (table 1, footnote a).

The average cost difference per person per year was around -€3000 in the matched gender × diagnosis groups, except in women with a musculoskeletal disease, for whom the difference was about -€1500 (figure 1a and appendix table A, [www.sjweh.fi/show\\_abstract.php?abstract\\_id=3780](http://www.sjweh.fi/show_abstract.php?abstract_id=3780)). Further subgroup analyses showed that in mental disorders, savings increased lin-


early with age in both genders. Overall, the savings were largest for 20–44 year-old men with a musculoskeletal disease and smallest for women in the same age and diagnostic group.

Among selected industrial sectors, cost savings were seen in manufacturing and in technical and scientific work in all gender × diagnosis groups. However, the difference in costs showed an inconsistent pattern in wholesale and retail trade as well as in human health and social work activities. In the latter sector, among men with musculoskeletal disease, the intervention resulted in an increase in social security costs (figure 1b and appendix table A).

The sensitivity analysis, excluding a few subjects with very high unemployment or retirement costs, showed total costs of €9.1 million in the intervention and €17.3 million in the control group, resulting in a cost ratio of 0.527 (95% CI 0.526–0.528).



**Figure 1a.** Mean differences in total social security costs per person per year (€) between the intervention and the control group by gender, diagnostic group and age group.



**Figure 1b.** Mean differences in total social security costs per person per year (€) between the intervention and the control group by gender, diagnostic group and selected industrial sector.

## Discussion

The current study provided an analysis of social security costs based on our previous quasi-experiment on the effects of part-time sick leave on work participation (9). We looked at what the social security cost difference would be during a two-year period if part-time sick leave was used instead of full-time sick leave to recover from a musculoskeletal disease or mental disorder. Our analysis within nationally representative register data showed that the use of part- instead of full-time sick leave at the early stage of work disability saved about half of all ill-health-related, unemployment and retirement costs during two years. The largest savings were attributable to differences in the costs of retirement and vocational rehabilitation. The savings were higher for the second compared with the first follow-up year.

Costs were saved in musculoskeletal diseases and mental disorders among both men and women, however, costs savings for women with a musculoskeletal disease were lowest. The savings increased linearly with age among both men and women with a mental disorder, however, there was no consistent pattern according to age among those with a musculoskeletal disease. The savings varied according to major industrial sector, being largest in manufacturing and technical and scientific work.

The strengths of this study are the nationally representative data where we had extensive information on social security benefits paid to the participants. A limitation was that, even though we could estimate the costs with a reasonable accuracy, the approximation may have affected the absolute values of the costs. Moreover, we assessed the costs from a social security perspective only and had no information on the costs of the employer in organizing part-time work and replacement for part-time or full-time SA that would have been necessary for a full economic analysis (19, 20). By using the propensity score, we attempted to minimize selection bias between the intervention and control group and achieved a good comparability between the groups with regard to the variables included in the propensity score (9). It is still possible that the groups were selected according to factors on which we had no information, eg, the individual's motivations regarding working (21).

In conclusion, we found that part- instead of full-time sick leave, at the early stage of work disability due to musculoskeletal diseases or mental disorders, leads to considerable social security cost savings during two years, in correspondence with increased work participation (9) and in addition to earlier reported health benefits (5). This study shows that part-time sick leave can be recommended also from an economic perspective; however, more consideration should be given to women

with musculoskeletal diseases as well as employees in human-health and social-work activities.

## Funding

The Academy of Finland (grant numbers 267589 and 303534) and NordForsk funded this work (grant number 76659).

## References

1. Kausto J, Miranda H, Martimo KP, Viikari-Juntura E. Partial sick leave--review of its use, effects and feasibility in the Nordic countries. *Scand J Work Environ Health.* 2008;34:239–49. <https://doi.org/10.5271/sjweh.1266>.
2. Kausto J. Effect of partial sick leave on work participation (Doctoral Thesis). People and Work Research Reports 102, Finnish Institute of Occupational Health 2013.
3. Høgelund J, Holm A, McIntosh J. Does graded return-to-work improve sick-listed workers' chance of returning to regular working hours? *J Health Econ.* 2010;29:158–69. <https://doi.org/10.1016/j.jhealeco.2009.11.009>.
4. Andrén D, Svensson M. Part-time sick leave as a treatment method for individuals with musculoskeletal disorders. *J Occup Rehabil.* 2012;22:418–26. <https://doi.org/10.1007/s10926-011-9348-7>.
5. Viikari-Juntura E, Kausto J, Shiri R, Kaila-Kangas L, Takala EP, Karppinen J, et al. Return to work after early part-time sick leave due to musculoskeletal disorders: a randomized controlled trial. *Scand J Work Environ Health.* 2012;38:134–43. <https://doi.org/10.5271/sjweh.3258>.
6. Kausto J, Viikari-Juntura E, Virta LJ, Gould R, Koskinen A, Solovieva S. Effectiveness of new legislation on partial sickness benefit on work participation: a quasi-experiment in Finland. *BMJ Open.* 2014 Dec 24;4(12):e006685. <https://doi.org/10.1136/bmjopen-2014-006685>.
7. Andrén D. Does part-time sick leave help individuals with mental disorders recover lost work capacity? *J Occup Rehabil.* 2014;24:344–60. <https://doi.org/10.1007/s10926-013-9467-4>.
8. Gran JM, Lie SA, Øyeflaten I, Borgan Ø, Aalen OO. Causal inference in multi-state models--sickness absence and work for 1145 participants after work rehabilitation. *BMC Public Health.* 2015 Oct 23;15:1082. <https://doi.org/10.1186/s12889-015-2408-8>.
9. Viikari-Juntura E, Virta LJ, Kausto J, Autti-Rämö I, Martimo KP, Laaksonen M, et al. Legislative change enabling use of early part-time sick leave enhanced return to work and work participation in Finland. *Scand J Work Environ Health.* 2017;43:447–56. <https://doi.org/10.5271/sjweh.3664>.
10. Briggs AH, O'Brien BJ, Blackhouse G. Thinking outside the box: recent advances in the analysis and presentation of uncertainty in cost-effectiveness studies. *Ann Rev Public Health.* 2002;23:377–401. <https://doi.org/10.1146/annurev>.

publhealth.23.100901.140534.

11. Social Insurance Institution of Finland Kela. Kelan tilastollinen vuosikirja 2010 (Statistical Yearbook of the Social Insurance Institution 2010). Official Statistics of Finland. Social Protection 2011. <https://helda.helsinki.fi/handle/10138/2264>.
12. Social Insurance Institution of Finland Kela. Kelan tilastollinen vuosikirja 2011 (Statistical Yearbook of the Social Insurance Institution 2011). Official Statistics of Finland. Social Protection 2012. <https://helda.helsinki.fi/handle/10138/2264>.
13. Social Insurance Institution of Finland Kela. Kelan tilastollinen vuosikirja 2012 (Statistical Yearbook of the Social Insurance Institution 2012). Official Statistics of Finland. Social Protection 2013. <https://helda.helsinki.fi/handle/10138/2264>.
14. Social Insurance Institution of Finland Kela. Kelan tilastollinen vuosikirja 2013 (Statistical Yearbook of the Social Insurance Institution 2013). Official Statistics of Finland. Social Protection 2014. <https://helda.helsinki.fi/handle/10138/2264>.
15. Saarnio L. Työeläkekuntoutus vuonna 2010 (Vocational rehabilitation in 2010, in Finnish). Eläketurvakeskuksen tilastoraportteja 03/2011 (Statistical reports of the Finnish Centre for Pensions 03/2011). Finnish Centre for Pensions, Helsinki, 2011.
16. Saarnio L. Työeläkekuntoutus vuonna 2011 (Vocational rehabilitation in 2011, in Finnish). Eläketurvakeskuksen tilastoraportteja 03/2012 (Statistical reports of the Finnish Centre for Pensions 03/2012). Finnish Centre for Pensions, Helsinki, 2012.
17. Saarnio L. Työeläkekuntoutus vuonna 2012 (Vocational rehabilitation in 2012, in Finnish). Eläketurvakeskuksen tilastoraportteja 03/2013 (Statistical reports of the Finnish Centre for Pensions 03/2013). Finnish Centre for Pensions, Helsinki, 2013.
18. Saarnio L. Työeläkekuntoutus vuonna 2013 (Vocational rehabilitation in 2013, in Finnish). Eläketurvakeskuksen tilastoraportteja 05/2014 (Statistical reports of the Finnish Centre for Pensions 05/2014). Finnish Centre for Pensions, Helsinki, 2014.
19. Uegaki K, de Bruijne MC, van der Beek AJ, van Mechelen W, van Tulder MW. Economic evaluations of occupational health interventions from a company's perspective: a systematic review of methods to estimate the cost of health-related productivity loss. *J Occup Rehabil*. 2011 Mar;21(1):90–9. <https://doi.org/10.1007/s10926-010-9258-0>.
20. van Dongen JM, van Wier MF, Tompa E, Bongers PM, van der Beek AJ, van Tulder MW, Bosmans JE. Trial-based economic evaluations in occupational health: principles, methods, and recommendations. *J Occup Environ Med*. 2014 Jun;56(6):563–72. <https://doi.org/10.1097/JOM.000000000000165>.
21. Sears JM, Rolle LR, Schulman BA, Wickizer TM. Vocational rehabilitation program evaluation: comparison group challenges and the role of unmeasured return-to-work expectations. *J Occup Rehabil*. 2014 Dec;24(4):777–89. <https://doi.org/10.1007/s10926-014-9509-6>.

Received for publication: 19 March 2018