Scand J Work Environ Health 2019;45(6):533-545

<https://doi.org/10.5271/sjweh.3832>

Published online: 23 May 2019, Issue date: 01 Nov 2019

Psychosocial effects of workplace exercise - A systematic review

by [Bordado Sköld M](#), [Bayattork M](#), [Andersen LL](#), [Schlünssen V](#)

This systematic review is the first to evaluate the effect of workplace exercise on psychosocial effects measured as both psychosocial work environment and mental health. The 22 randomized controlled trials included show limited psychosocial effects of workplace exercise. Variety in types of interventions as well as outcome measurements limit comparison across studies. More efficient and comparable study designs should be prioritized.

Affiliation: National Research Centre for the Working Environment, Lærsø Park Allé 105, 2100 Copenhagen East, Denmark. marhen60@gmail.com

Refers to the following texts of the Journal: [2006;32\(6\):443-462](#)
[2011;37\(1\):70-76](#) [2014;40\(4\):390-399](#) [2012;38\(2\):93-104](#)

Key terms: [mental health](#); [occupational health](#); [physical activity](#); [preventive medicine](#); [psychosocial work environment](#); [workplace intervention](#)

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/31120539

Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](#).

This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

Psychosocial effects of workplace exercise – a systematic review

by Margrethe Bordado Sköld, MD,^{1,2} Mohammad Bayattork, MSc,^{3,4} Lars L Andersen, PhD,^{1,5} Vivi Schlünssen, PhD^{1,6}

Bordado Sköld M, Bayattork M, Andersen LL, Schlünssen V. Psychosocial effects of workplace exercise – a systematic review. *Scand J Work Environ Health*. 2019;45(6):533–545. doi:10.5271/sjweh.3832

Objectives Workplace exercise interventions are known to reduce musculoskeletal complaints and are widely offered to promote employee's health. The psychosocial effects of such interventions are not clear. This review evaluates the effects of workplace exercise interventions on the psychosocial work environment and mental health.

Methods A systematic review was performed by screening four scientific databases (Pubmed, Cochrane Database, PsycINFO and Web of Science) for published randomized controlled trials in English from 1998–2018. The review followed the PRISMA statement guidelines using the Cochrane tool for assessing risk of bias and the GRADE approach for assessment of quality. Two researchers independently performed database search, study screening, and evaluation.

Results Twenty two randomized controlled trials met the inclusion criteria. Substantial heterogeneity in the studies included was found in study populations, types of intervention, and outcome measures performed. Quality assessment revealed a generally low level of quality in the studies included. This was partly the result of the limited possibilities for blinding and the self-reported outcome measures.

Conclusion Workplace exercise interventions seem to have limited effects on mental health and the psychosocial work environment. Future studies should incorporate blinding and independent outcome measures.

Key terms mental health; occupational health; physical activity; preventive medicine; psychosocial work environment; stress; workplace intervention.

Most people will spend up to one-third of their adult life at work, thus workplace health promotion interventions are considered to have great potential for improving health. For instance, occupational organizational changes have been initiated to help increase smoking cessation, reduce alcohol consumption and introduce healthier diets (1). The increasing proportion of people working in sedentary jobs is a contributing factor to the cumulated insufficient physical activity in the general population during the past decades (2–4) and insufficient physical activity is now globally one of the main causes of increased mortality and reduced ability (5). Occupational sedentary behavior is suspected to be associated with health risks (6) and many workplaces have focus on counteracting sedentary behavior through workplace exercise interventions.

Mental health

Mental health problems among workers is another major challenge and impacts workers themselves, employers, and society through reduced work ability, increased sick leave, and higher risk of disability retirement (7, 8). Work-related stressors are listed as the main cause of stress in 70% of workers with self-reported stress in the Danish workforce (9) and, despite political and occupational interventions to reduce work-related stressors, statistics show that the problem is still increasing (9).

Workplace exercise interventions

The positive effects on health of regular exercise are well known, such as reducing the risk for obesity,

¹ National Research Centre for the Working Environment, Copenhagen East, Denmark

² Department of Occupational and Social Medicine, Holbæk University Hospital, Holbæk, Denmark.

³ Sport Sciences and Physical Education, Faculty of Humanities Science, University of Hormozgan, Bandar Abbas, Iran.

⁴ Health and Sports Medicine Department, Faculty of Physical Education and Sport Sciences, University of Teheran, Teheran, Iran

⁵ Sport Sciences, Department of Health Science and Technology, Aalborg University, Aalborg, Denmark.

⁶ Department of Public Health, Environment, Occupation and Health, Danish Ramazzini Centre, Aarhus University, Aarhus C, Denmark.

Correspondence to: Margrethe Bordado Sköld, National Research Centre for the Working Environment, Lærso Park Allé 105, 2100 Copenhagen East, Denmark. [E-mail: marhen60@gmail.com]

cardiovascular diseases, diabetes, osteoporosis and some types of cancer (2, 10, 11). Regular exercise is also known to help prevent depression (12), reduce self-reported emotional stress (13–15), and improve quality of life and self-rated health (3, 10, 16).

Recently there has been an increasing number of studies examining the effects of workplace exercise interventions, showing that regular workplace exercise can reduce musculoskeletal complaints (17, 18), improve fitness, and reduce the risks of cardiovascular disease (19, 20). Effects on other parameters, such as mental health, quality of life, and the psychosocial work environment have also been suggested. For example, in their 2014 systematic review, Chu et al (21) evaluated the effects of work or home based exercise intervention on workers' mental health. The evaluated outcomes included depressive, anxiety, and stress symptoms, where 2 out of 3 studies showed significant effect on depressive symptoms, 2 of 7 showed evidence of reducing anxiety symptoms, and 7 of 13 showed possible improvement of stress relieve. In a meta-analysis from 2009 (19), Conn et al evaluated the effects of workplace exercise interventions on different health parameters and found a positive association on quality of life and mood measurements, though not reaching statistical significance. More recently, Danish studies have indicated that workplace exercise might influence workplace parameters such as work ability (22, 23) and sickness absence (22) as well as the psychosocial work environment, eg, social capital (24, 25). Other studies have shown uncertain effects of workplace exercise interventions (19, 26).

Thus, there might be a positive effect of workplace exercise intervention on both mental health as well as the psychosocial work environment.

Psychosocial work environment

Psychosocial work environment is a known predictor for common mental health problems (27), but a possible link between workplace exercise and how this impact the psychosocial work environment is not clear. Based on studies showing positive effects of workplace exercise on physical capacity, reduction of pain and employee wellbeing, one might hypothesize that this could lead to effects on factors impacting the psychosocial work environment. Furthermore, integration of exercise interventions will inevitably change the context and framework of a workday (eg, meeting colleagues and leaders in a different setting, changes in the distribution and organizing of employees' work assignments and break schedules, changes in work-life-balance (as exercise during work hours may influence leisure time activities, etc.) which also might influence parameters of the psychosocial work environment.

One of the models to operationalize the psychosocial

work environment is Karasek's job-strain model (28), which points to the essential relationship between job demands and decision latitude. According to the job-strain model, the amount of work demands together with the amount of work control arbitrate the amount of psychological strain. The model was later expanded to include occupational social support (29), also known as the iso-strain model, as social support affects how demands and control at work are perceived. The authors of the present review acknowledge the definition of psychosocial work environment based on the iso-strain model. However, when measured as an outcome in research, the difference between psychosocial work environment and mental health parameters are not always clearly separated. Often, the fields seem to interconnect in the measurement tools used. For instance, measurement outcomes could be social capital, (including cooperation, trust, vitality), quality of life, job strain (including psychological demands, work load, levels of influence, effort–reward balance), perceived job stress, vitality, fatigue, general wellbeing, burnout, as well as specific depression and anxiety scores.

As the two fields not always are clearly separated, we found it relevant to clarify a possible correlation between workplace exercise interventions to both mental health and psychosocial work environment.

We are not aware of any published reported or ongoing studies evaluating effects of workplace exercise on psychosocial work environment as well as mental health.

Objectives

The aim of this study was to evaluate peer-reviewed randomized controlled trials investigating the effects of workplace exercise on psychosocial working environment and mental health among employees. The study follows the PRISMA (Preferred Reporting Items for Systematic Review and Meta-Analysis) statement guideline for systematic reviews (30).

Methods

Eligibility criteria

Study design, setting and language. The following inclusion criteria were used for this systematic review: (i) Peer-reviewed publication (in press or accepted for publication) reported in English; (ii) Study method consisting of a randomized controlled trial; (iii) Exercise intervention for employees occurring at the workplace or during working hours; (iv) Intervention conducted minimum on a weekly basis; (v) Outcome including 'mental health' and/or 'psychosocial environment'.

As there are a great variety of different tools used to measure mental health and the psychosocial work environment – and as many of the measured outcomes seem to overlap – we found it necessary to use these broad outcome definitions. To ensure relevance of the studies included, each study's outcomes were therefore specifically addressed, sorted, and organized after best relevance, based on the studies own categorization.

In our protocol, we initially chose an inclusion criterion of a minimum of 12 weeks intervention duration as we expected a great amount of studies eligible for review inclusion. After performing the initial study searches however, we chose to remove this criterion. We realized that, overall, there were not many studies meeting all inclusion criteria, meaning we could potentially miss relevant studies for our purpose.

Studies measuring workability, absenteeism, and sick leave without other psychosocial or mental health outcomes were not included. In order to assess current workplace and labor market conditions, we included original, peer-reviewed published articles in English from 1 January 1998 to 15 August 2018.

Information sources

The following electronic databases were searched for eligible published studies: Medline, Cochrane database, PsycINFO and Web of Science. The search ended on 15 August 2018. No new studies were included after this date.

Search strategy

The search terms included words related to the eligible criteria of setting, intervention, and outcome. Separate search strings were formed for each of the databases, taking into account their different frameworks (table 1). The complete search strategy for all databases is available in the supplementary material (www.sjweh.fi/show_abstract.php?abstract_id=3832).

Study selection

Selection process, and data management. Two researchers independently screened the papers identified in the listed databases on title and abstract level for possible eligible studies meeting inclusion criteria. Systematic reviews in the research field were also screened for eligible studies. The software program Endnote was used for sorting and duplicate removal. Eligible full-text articles were then read and evaluated for possible inclusion. Two researchers independently extracted data and assessed risk of bias. Any disagreements in the selection and extraction process was discussed between the two researchers until consensus was reached. When necessary a third researcher offered assistance in the evaluation process.

Data extraction

The following data were extracted from the studies: Country where the study was conducted, year of publication, population (type of workplace or job task, recruitment process, number of participants, inclusion health status), description of the intervention (type of training, duration, group- or individual-based), information of statistical comparison analyses, outcomes measured (including type and validation of surveys used for measurement), information on follow up measurements.

Risk of bias in individual studies

The Cochrane tool for assessing risk of bias from the *Cochrane Handbook for Systematic Reviews of Interventions* (31) was used to assess potential biases in the included studies. This assessment of bias includes potential biases of selection, performance, detection, attrition, reporting, and other sources from seven risks of bias domains (random sequence generation, allocation concealment, blinding of participants and personnel, blinding of outcome assessment, incomplete outcome data, selective reporting, as well as other potential threats to validity).

Quality assessment

Quality was evaluated using the Grading of Recommendations, Assessment, Development and Evaluation (GRADE) approach (32), which grades studies as followed: very low, low, moderate, high. The quality assessment was based on the findings in data extraction, and follows the domains of quality evaluation in the GRADE approach: risk of bias, inconsistency, indirectness and imprecision (figure 1).

Table 1. Cochrane database. Results: N=204

#1	MeSH descriptor: [Workplace] explode all trees	711
#2	MeSH descriptor: [Mental Health] explode all trees	1142
#3	psychosocial work environment:ti,ab,kw (Word variations have been searched)	85
#4	MeSH descriptor: [Depression] explode all trees	9572
#5	MeSH descriptor: [Anxiety] explode all trees	6866
#6	MeSH descriptor: [Occupational Stress] explode all trees	198
#7	(occupational exercise or occupational physical activity or occupational workout or workplace exercise or workplace physical activity or workplace workout):ti,ab,kw (Word variations have been searched)	1248
#8	("occupational stress" or depression or anxiety or stress or "psychosocial work environment"):ti,ab,kw (Word variations have been searched)	104 151
#9	(exercise or "physical activity" or workout):ti,ab,kw (Word variations have been searched)	95 766
#10	#2 or #3 or #4 or #5 or #6 or #8	104 901
#11	#1 and #9	234
#12	#7 or #11	1248
#13	#10 and #12	204

Figure 1. PRISMA Flow chart of study selection process.

Results

Study selection

A total of 1560 records were identified by searching the databases PubMed, Web of Science, PsycINFO and the Cochrane database. After duplicate removal ($N=74$), review removal ($N=4$), and title abstract screening, 24 studies were evaluated for eligibility by full-text reading. The four reviews were screened for possible included eligible records, but no additional studies meeting the inclusion criteria of this review were identified. Two studies did not meet the inclusion criteria (one was not a randomized controlled trial and the other did not offer exercise as part of its intervention) and were excluded, leaving a final inclusion of 22 studies.

Study characteristics

The characteristics of the studies included are presented in table 2. Most studies ($N=17$) (24, 25, 33–47) were conducted in Europe, the rest in North America (48, 49), Japan (50, 51), and Australia (52). The workplace

settings varied from employees working in office/having administrative tasks ($N=10$) (34, 35, 37, 38, 41, 43, 47–49, 52), to health personnel in hospitals/nursing homes ($N=5$) (25, 33, 44, 46, 50), to laboratory technicians ($N=2$) (39, 42), to other manual job settings or workplaces with a mix of work tasks ($N=5$) (24, 36, 40, 45, 51). Sample size ranged from 24–860 participants, with a total of 4010 participants throughout the studies. The majority of studies ($N=12$) (25, 33, 34, 36, 38, 40, 42, 43, 46, 50–52) were based on healthy workers, three studies (24, 39, 48) examined workers with musculoskeletal problems, three studies (41, 45, 49) looked at sedentary/physical inactive workers, two studies (35, 47) included workers with light mental health complaints, one study (37) looked at workers with both mental and/or musculoskeletal complaints, and another (44) examined high aged workers. The exercise interventions consisted of a wide range of different exercises offered from 6–86 weeks (21 months), with most studies offering an intervention period ranging from 8–12 weeks (24, 25, 33–37, 39, 47, 48, 50–52) $N=13$. Interventions could be subdivided into cardio/aerobics ($N=3$) (35, 36, 46), strength/resistance ($N=6$) (24, 25, 33, 40, 42, 43), yoga

Table 2. Summary of findings. [C=control; I=intervention; N/A=not available; NW=nature walk; UW=urban walk.]

Study	Classic / cluster Waitlist/ cross-over	Outcome Primary / Secondary / Neither	Participants women: men ratio	Workplace	Duration	Type	Comparison (analysis method)	Dropout rate	Effects on mental health factors (endpoint & follow-up)	Effects on psychosocial work environment	Extended follow-up
Eriksen et al, 2002 Norway	Classic	Neither (among others)	N=860 Ca 1:1	Post office/ terminals	12 wks	Group I: - Aerobics - Aerobics with emphasis on work relevance + health program - Stress management	I: 3 C: 1 (one-way ANOVA and χ^2 tests)	C: 52% I: (3 types) 41%	No significant effect in stress level at endpoint or extended follow-up	N/A	1 yr
Van rehenen et al, 2005 The Netherlands	Classic	Primary	N= 130 1:9	Office/ admin (telecoms company)	8 wks	Individual I: Physical + relaxation exercises	I vs cognitive (MANOVA)	Cognitive I: 61% Exercise I: 55%	No significant effect on psychological complaints, burnout and fatigue at endpoint or extended follow-up	N/A	6 mths
Sjögren et al, 2006 Finland	Cluster (N=4) / cross over	Neither (among others)	N=90 4:1	Office/ admin	15 wks	Group and individual I: Light resistance training	Cross over after 15 wks (linear mixed models)	I 1:3.6% I 2: 5.7%	No significant effect on psychosocial functioning and general well-being at endpoint or extended follow-up	N/A	30 wks (ending cross over)
Tveito et al, 2009 Norway	Classic / waitlist	Secondary	N=40 1:0	Health personnel (nursing home)	36–37 wks (9 mths)	Group I: Aerobics based + health program	I vs C (one-way ANOVA and χ^2 tests)	I: 37% C: 19%	No significant effect on coping, job stress, subj. health complaints and health-related quality of life. Significant positive effect on health and well-being	No significant effect on work environment, effort-reward imbalance, demands or control	None (sick leave data 1 year after)
De zeeuw et al, 2010 The Netherlands	Classic / wait list	Primary	N=30 Ca. 1:1	Office/administrative (insurance company)	10 wks	Group I: Cardiovascular (power, cycling, stairs, situps etc)	I vs C (ANCOVA and χ^2)	I: 7% C 13%	No significant effect on depression score	N/A	None
Hartfiel et al, 2011 UK	Classic / wait list	Primary	N=48 Ca. 9:1	Office/administrative (university)	6 wks	Group I: Yoga	I vs C (two-way ANOVA)	I: 17% C: 17%	Significant positive effect on anxiousness, confusion, depressed, tired, and unsureness, greater sense of life purpose and satisfaction and more self-confidence	N/A	None
Hartfiel et al, 2012 UK	Classic / wait list	Neither (among others)	N=76 Ca. 9:1	Office / administrative	8 wks	Group I: Yoga	I vs C (ANOVA and multiple linear regression)	I:11% C: 30%	Significant positive effect on perceived stress and psychological well-being	N/A	None
Mesquita et al, 2012 Portugal	Classic / wait list	Neither (among others)	N= 229 0:1	Ware-house	86 wks (21 months)	Group I: Spinal strength and stability exercise	I vs C (Friedman ANOVA test)	I: 57% C: 60%	Significant positive effect on mental health	Significant positive effect on social functioning and role limitations due to emotional problems	None
Strijk et al, 2012 The Netherlands	Classic	Neither (among others)	N=730 Ca. 4:1	Hospital	24-25 wks (6 months)	Group and individual I: - Yoga session - Aerobic and strength session - Individual session	I vs C (linear regression analyses)	I:20% C: 22%	No significant effect on mental health	N/A	None
Cheema et al, 2013 Australia	Classic	Secondary	N=37 Ca.5:1	Office/adm (university)	10 wks	Group I: Yoga	I vs C (ANCOVA)	I: 6% C: 11%	No significant effect on anxiety, health related Quality of Life	No significant effect on job satisfaction)	None

Continued

Table 2. continued

Study	Classic / cluster Waitlist/ cross-over	Outcome Primary / Secondary /Neither	Participants women: men ratio	Workplace	Duration	Type	Comparison (analysis method)	Dropout rate	Effects on mental health factors (endpoint & follow-up)	Effects on psychosocial work environment	Extended follow-up
Roessler et al, 2013 Denmark	Classic (randomization thru clusters) / wait list	Secondary	N=427 Ca. 6:1	Laboratory (industry production)	20 wks	Group and individual I: Neck+ shoulder strength training	I vs C (MANOVA)	I:29% C: 11%	N/A	No significant effect on job satisfaction, influence, sense of community and time pressure	None
Brown et al, 201 UK 4	Classic	Secondary	N=94 1:4-1:5	Office /adm. (stock company)	8 wks	Group and individual I: - NW - UW	I nature vs I urban vs C (ANOVA)	I NW:16% I UW: 18% C: 34%	Significant positive effect mental health (low adherence in all walk groups)	N/A	None
Andersen et al, Denmark (N=18)	Cluster (N=18)	Secondary	N=200 1:0	Nurse/ nurses aid (hospital)	10 wks	Group I: Strength training	I vs home training (linear mixed model)	Work: 9% Home: 7%	N/A	Significant positive effect on social capital within workplace teams	None
Jay et al, 2015 Denmark	Classic	Secondary	N=112 1:0	Laboratory technicians (pharmacy)	10 wks	Group and individual I: Motor control + resistance + education + mindfulness	I vs C (linear mixed model)	I:5% C:5%	No significant effect on stress	N/A	None
Thøgersen et al, 2015 UK	Classic / waitlist	Primary	N=75 12:1	Non-academic employees (university)	16 wks (10 + 6 wks)	Group and individual I: Walking	I vs waitlist (mixed linear modeling)	I: 26% Waitlist: 25%	Significant positive effect on enthusiasm and relaxation	N/A	Unclear
Taylor et al, 2016 USA	Cluster (N=18)	Neither (among others)	N=185 5:1	Office / admin (not specified)	24-25 wks (6 months)	Group I: Stretch+ strength+ aerobic Individual I: walk brakes	I vs C (generalized mixed-effect, log normal)	Booster I: 37% PC I: 41% C: 44%	No significant effect on quality of life	N/A	None
Andersen et al, 2017 Denmark	Classic	Secondary	N=66 Ca. 1:4	Manual work (slaughter-house)	10 wks	Group I: Strength training	I vs C (linear mixed model)	I:1% Reference (ergo): 6%	No significant effect on mental health	Significant positive effect on social climate and vitality	none
Chopp-hurley et al, 2017 Canada	Classic/ wait list	Neither (among others)	N= 24 4:1-6:1	Academic+ non-academic (university)	12 wks	Group I: Yoga poses for lower limb strength	I vs C (one-way ANOVA)	I: 25% C: 8%	Significant positive effect on depressive symptoms No significant effect on self-efficacy	No significant effect on work resilience	None
Jakobsen et al, 2017 Denmark	Cluster (N=18)	Secondary	N=200 1:0	Nurses/ nurse aid (hospital)	10 wks	Group I: Strength training	I vs home trainin (linear mixed model)	Work: 9% Home: 7%	Significant positive effect vitality and concern	No significant effect on work-and leisure disability, emotional demands, influence at work, community sense, social support and mental health	None
Matsugaki et al, 2017 Japan	Classic	Secondary	N=30 1: 0	Nurses (Hospital)	12 wks	Group I: Resistance and aerobic training	I vs unsupervised training (Wilcoxon signed-rank test+ Mann-Whitney U test)	I: 0% C 7%	No significant effect on psychological status and depression level	N/A	None

Continued

Table 2. continued

Study	Classic / cluster Waitlist/ cross-over	Outcome Primary / Secondary / Neither	Participants women: men ratio	Workplace	Duration	Type	Comparison (analysis method)	Dropout rate	Effects on mental health factors (endpoint & follow-up)	Effects on psychosocial work environment	Extended follow-up
Michishita et al, 2017 Japan	Cluster randomization (N=11)	Neither (among others)	N=63 1:3	White collar workers (large company 31 workplaces)	10 wks	Group I: Resistance and aerobic training	I vs C (Mann-whitney U-test, Wilcoxon, chi ² and simple linear regression analysis)	I: 3% C: 1%	Significant positive effect fatigue, vigor, friendliness	Significant positive effect interpersonal stress, vigor, support, satisfaction	None
Puig-ribera et al, 2017 Spain	Cluster randomized (N=6)	Secondary	N=264 3:1	Office/adm (university)	19 wks (8+11)	Individual I: web-based encouragement to move more	I vs C (linear mixed model)	I: 32% C: 24%	No significant effect on mental well-being at end point or extended follow-up	N/A	2 months

(N=4) (37, 38, 48, 52), walks (N=2) (34, 45), web-based (N=1) (41) and mixed (N=6) (39, 44, 47, 49–51), though with wide individual differences between studies. Group training was most common with thirteen studies (24, 25, 33, 35–38, 40, 46, 48, 50–52) offering this, individual training occurred in two studies (41, 47), and a mix in seven (34, 39, 42–45, 49). Many studies were designed with an active reference group, offering an alternative exercise (24, 25, 33, 34, 36, 43, 49).

Only four studies (35, 38, 45, 47) had psychosocial outcomes as their primary outcomes. In ten (24, 25, 33, 34, 39, 41, 42, 46, 50, 52), it appeared as secondary outcome whereas it was one equal outcome among others in eight studies (36, 37, 40, 43, 44, 48, 49, 51). The same trial was used in two studies (25, 33) (intervention and study population) to conduct two different analyses with different outcome measures. Thirteen studies (34–39, 41, 43–45, 47, 49, 50) focused on mental health outcomes, while two (33, 42) focused on psychosocial environment, and seven (24, 25, 40, 46, 48, 51, 52) examined effects on both. All psychosocial outcomes were measured through self-reported questionnaires. Some questionnaires were recurring in the studies, the Copenhagen Psychosocial Questionnaire COPSOC was seen in two (25, 42), the survey SF-36 was used in five, with focus on quality of life in three studies (40, 46, 52) and mental health in the two latter. Other measures reported revealed a great variety in types of questionnaires used in the different studies. Some mentioned questionnaires were: the Maslach Burnout Inventory–General Survey for burn out; 4DSQ for psychological complaints; the Demand/Control Model and the Job Affect Scale (JAS) for psychological demands; Profile of Mood States Bipolar (POMS-Bi); Inventory of Positive Psychological Attitudes (IPPA); PHQ-9; Center for Epidemiologic Studies Depression Scale

(CES-D); Beck Depression Inventory-Second Edition (BDI-II) 9 for depressive symptoms; SF-8 for mental health; State-Trait Anxiety Inventory (STAI) for anxiety; Perceived Stress Survey (PSS) and Cohen's perceived stress questionnaire for stress symptoms; PANAS-X and Warwick-Edinburgh Mental Well-Being Scale for mental/ psychological well-being; the Arthritis Self efficacy scale (ASES) for self-efficacy; the Work Extrinsic and Intrinsic Motivation Scale for work motivation; the Copenhagen Psychosocial Questionnaire (COPSOC) for psychosocial work environment; and the General Nordic Questionnaire for Psychological and Social Factors at Work for social capital.

Furthermore, many measurements were based on self-reported 5-point Likert scales rating of perceived subjective symptoms as well as psychosocial work environment, developed for the studies.

All studies performed between group statistical analysis and included sufficient description of which methods were used in the study. Most studies measured outcomes two times: at baseline and at the end of the intervention period, a single study [Thøgersen et al (45)] measured continuously through mobile phones twice a week during the intervention period.

Quality assessment including risk of bias

The quality assessment of the included studies is presented in table 3. Risk of bias assessment was performed using the Cochrane Risk of Bias tool (31) and based on the evaluation the studies were graded into four categories: very high risk, high risk, moderate risk, low risk. Other parameters used to evaluate the quality of the studies were based on the GRADE approach (32) and included (i) description of statistics used for comparison, (ii) representative value of the included population, (iii)

Table 3. Quality assessment. [C=control without intervention; C+I=control with other intervention.]

Study	Risk of bias (4=very high 3=high 2=moderate 1=low)	Statistics ^a	Representative		Number of participants	Intervention		Comparison	Outcome		Extended follow-up >1 month	Quality of Evidence
			Population	Exclusion consequence ^b		Realistic	Easy integration		Common measure	Validated question- naire		
Eriksen et al, 2002	2	+	Healthy	Yes	>100	Yes	Yes	C and C+I	Yes	Yes	Yes	Moderate
Van rhenen et al, 2005	4	+	Specific	Yes	>100	Yes	Yes	C	Yes	Yes	Yes	Very low
Sjögren et al, 2006	2	+	Healthy	No	>100	Yes	Yes	C	Yes	Yes	Yes	Moderate
Tveito et al, 2009	3	+	Healthy	No	31–100	Yes	Yes	C	Yes	Yes	Yes	Low
De zeeuw et al, 2010	2	+	Specific	Yes	31–100	Yes	Yes	C	Yes	Yes	No	Moderate
Hartfiel et al, 2011	3	+	Healthy	Yes	<30	Yes	Yes	C	Yes	Yes	No	Low
Hartfiel et al, 2012	3	+	Specific	Yes	31–100	Yes	Yes	C	Yes	Yes	No	Low
Mesquita et al, 2012	4	+	Healthy	Yes	>100	Yes	Yes	C ^c	Yes	Yes	No	Very low
Strijk et al, 2012	2	+	Specific	No	>100	No	No	C	Yes	Yes	No	Moderate
Cheema et al, 2013	2	+	Healthy	Unclear	31–100	Yes	Yes	C	Yes	Yes	No	Moderate
Roessler et al, 2013	2	+	Healthy	No	>100	Yes	Yes	C	Yes	Yes	No	Moderate
Brown et al, 2014	3	+	Healthy	Yes	>100	Yes	Yes	C ^d	Yes	Yes	No	Low
Andersen et al, 2015	2	+	Healthy	Yes	>100	Yes	Yes	C+I	No	Yes	No	Moderate
Jay et al, 2015	2	+	Specific	Yes	>100	Yes	Yes	C	Yes	Yes	No	Moderate
Thøgersen et al, 2015	4	+	Specific	Yes	31–100	Yes	Yes	C	No	No (mobile) Yes	No	Low
Taylor et al, 2016	3	+	Specific	No	>100	Yes	Yes	C	Yes	Yes	No	Low
Andersen et al, 2017	2	+	Specific	Yes	31–100	Yes	Yes	C+I	Yes	Yes	No	Moderate
Chopp-hurley et al, 2017	3	+	Specific	Unclear	<30	Yes	Yes	C+I	No	Yes	No	Low
Jakobsen et al, 2017	2	+	Healthy	Yes	31–100	Yes	Yes	C+I	No	Yes	No	Moderate
Matsugaki et al, 2017	3	+	Healthy	Unclear	<30	Yes	Yes	C+I	Yes	Yes	No	Low
Michishita et al, 2017	3	+	Healthy	Yes	31–100	Yes	Yes	C	Yes	Yes	No	Low
Puig-ribera et al, 2017	2	+	Specific	Yes	>100	Yes	Yes	C	Yes	Yes	Yes	Moderate

^a Comparison calculations MANOVA / ANCOVA / ANOVA / Logistic regression.^b Mental/physical health issues were a population exclusion criteria in the study.^c Inside group only comparison, not valid due to large dropouts.^d Inside-group only comparison, not valid due to low adherence.

sample size, (iv) accessibility and transferability of the exercise intervention, (v) outcome comparison, (vi) outcome measurement validity, and transferability and (vii) presence of and duration of follow up

Risk of bias. No studies were graded as having *low risk of bias*, eleven (24, 25, 33, 35, 36, 39, 41–44, 52) met the criteria for having *moderate risk*, eight (34, 37, 38, 46, 48–51) were graded to have *high risk*, and three (40, 45, 47) were found to have a *very high risk of bias*. Blinding was declared not possible or not commented on in most studies, with the exception of two [Jay et al (39) and Puig-Ribera et al (41)] who reported to be single-blinded. All reported outcomes relevant for this

review were collected with self-reported questionnaires and thus not blinded. There were great differences in drop-out ranges. Ten of the studies (24, 25, 33, 35, 36, 39, 42–44, 52) reported that intention-to-treat analyses were performed, leaving the remainder (N=12) (34, 37, 38, 40, 41, 45–51) with high risks of attrition biases.

(i) *Statistics.* A sufficient description of a relevant performed statistical analysis for comparison was delivered in all studies, with a large variety of methods observed.

(ii) *Population.* Generally, the participants represented highly selective groups of workers, as many studies investigated workers with specific issues (eg, neck and shoulder pain), and/or because participants

were recruited through self-responsive invites where responsive rate was low, and/or many participants were excluded from the study due to health-related issues or other reasons.

(iii) *Sample size.* Ten of the studies (25, 33, 36, 39–42, 44, 47, 49) had a participant population >100, eight studies had 31–100 participants (24, 34, 37, 38, 43, 45, 46, 51), whereas three studies (35, 48, 50) were performed with very small sample sizes $N \leq 30$.

(iv) *Intervention design.* Most studies presented exercise interventions that, from the descriptions in the study designs appeared easy to implement and access at the work places

(v) *Comparison.* When presenting results, all studies included between-groups analyses, within-group analyses or subgroup analyses performed in many studies were not included in this review.

(vi) *Measurement.* Most measurement surveys were well-known validated questionnaires, however, studies also introduced uncommon and non-validated surveys. Use of single questions from domains was also seen in some studies.

(vii) *Extended follow-up.* Only four studies (36, 41, 43, 47) conducted extended follow-up measurements (ie, analyses a time period after intervention ended) with one year follow up being the longest duration of follow up.

The quality of evidence assessment revealed that twelve studies met the criteria for moderate quality, eight were rated with low quality and two very low quality of evidence. No studies met the criteria for high quality of evidence. Overall, positive effects on mental health and psychosocial work environment were typically found in only a single or a few parameters in the surveys used, the effects reported were found in different psychosocial parameters in the different studies, the reported effect sizes were found to be small. Six studies, Tveito et al (46), Hartfiel et al (38), Hartfiel et al 2012 (37), Brown et al (34), Thøgersen et al (45), Chopp-Hurley et al (48), and Jakobsen et al (25), reported a beneficial effect on mental health after intervention. Brown et al (34) reported that the positive effect found probably was not valid due to low adherence. Two studies by Andersen et al (24, 33) reported positive effects on the psychosocial work environment “social capital” (33) and “social climate” and “vitality” (24), respectively. Two studies, Mesquita et al (40) and Michishita et al (51), reported positive effects on both mental health and psychosocial environmental outcomes on a variety of different parameters. However, these two studies were graded with “very low” and “low” quality of evidence, which should be taken into account when evaluating effects. A total number of eleven studies [Eriksen et al (36), Van Rhenen et al (47), Sjögren et al (43), De Zeeuw et al (35), Strijk et al (44), Cheema et al (52), Roessler et al (42), Jay et al (39), Taylor et al (49), Matsugaki et al (50)

and Puig-Ribera et al (41)] reported no effects at all on psychosocial outcomes.

Discussion

Summary of evidence

The results of this review revealed that workplace exercise generally have limited positive effects on the psychosocial work environment and employers’ mental health. The majority of the included studies reported no or limited effects on a single or few parameters measured. None of the studies reported a negative effect on the psychosocial work environment. Thus, it is possible that exercise interventions may have effects on psychosocial parameters, both in mental health and psychosocial environment outcomes, however these effects seem to be small and dependent on the setting. The wide range of effects on different parameters probably reflects the great variety of design and methods of the studies.

Strengths and limitations

Value of results. To the best of our knowledge, this is the first study to review workplace exercise interventions and its effects on the psychosocial work environment as well as on mental health. By including a broad range of psychosocial outcomes, we are able to report a comprehensive evaluation of an emerging field of attention: workplace interventions to improve employers’ health. Thus, the results of this review might influence attitudes and actions for improving the work environment and potentially affect the overall working population. Workplaces today offer a great variety of types and integration of workplace intervention programs to improve health. Given the amount of time the working population spend at the worksite, workplace intervention programs are seen as possible efficient ways to enhance employees’ health (17). The results of this review indicate, however, that the effects of these programs on psychosocial parameters is limited. Nevertheless, workplace interventions are today acknowledged and encouraged by both governments and health organizations as a part of a general occupational health promotion (53–55), and workplace wellness, including fitness or other types of exercise at work, are commonly offered by employers (17, 56, 57).

Naturally, developing and implementing convenient programs and settings for exercise enhancement have concurrently been in focus, which was reflected in the studies of this review as most of the intervention designs chosen appeared to be easy to integrate and would transfer to other settings.

Strengths

This review included searches in many different academic databases, hereby reducing the risk of missing relevant studies for inclusion. We did not include grey literature or unpublished materials as we judged the academic literature to be the best source to answer our research question. It is possible that we have not included all trials in the area, but we do not think this influenced our conclusion. By following the PRISMA guidelines for systematic reviews, we ensured a systematic research process, which included a parallel independent screening, data extraction, and risk of bias assessment by two researchers, a method that reduces errors in the evaluation across studies.

Limitations

Still, the results of this review must be interpreted with caution and in accordance with its limitations. Of the 22 included studies, none were graded as having high quality evidence. A main reason for this is the weight of selection, performance, and detection bias (blinding) applied in the quality assessment method (Cochrane and GRADE). Blinding of participants in behavioral interventions is not possible, as participants are aware of what they are doing. Furthermore, psychosocial outcomes are difficult to measure without self-reported questionnaires. Still, we chose to evaluate studies with these risk of bias assessment tools as blinding increases credibility and validity of study results, and using the original, non-modified tool for risk of bias ensured transparency in our methods.

The inclusion of study results from the two studies graded as very low (40, 47) did not change the overall findings in this review.

Due to the great variety of study designs (population, types and duration of exercise interventions, tools used for self-reported measures of psychosocial outcomes), it is difficult to perform a synthesis across studies, which affects the possibility of drawing overall conclusions. The study populations were typically highly selected groups of participants, which lowers the external validity of the outcomes. Finally, we do not know if a longer period of intervention would have changed outcomes for some of the trials, but there was no consensus in positive results in the eight studies offering the intervention for >12 weeks.

Perspectives

Based on connections between physical and mental health, and mental health and the psychosocial work environment, we hypothesized that there might be positive effects of workplace exercise on psychosocial

parameters. This was not supported in the results of this review. However, our findings can also illustrate the complexity in evaluating the wide range of workplace health promotion interventions offered, as physical and mental health at work might, as mentioned, be interconnected, and context-dependent local culture might affect outcome together with the different variables as type of workplace, population, setting, intervention, motives for implementation etc. Overall, the large heterogeneity of the studies strongly implies the need for more corresponding intervention designs and use of measurement tools in the future to enable better comparison.

Better methods for studying workplace interventions are desirable, but difficult as blinding is not possible. Some of the studies included in this review provided an option to blinding by offering two different interventions, where the participants are likely to believe that both may have benefits. This was done in the studies by Eriksen et al (36), Brown et al (34), Andersen et al (24, 33), Taylor et al (49), and Jakobsen et al (25), where there were active comparison groups. Examples of such were comparison of exercise at home versus work (25, 33), instructor- versus computer-based exercise (49), instructor-based exercise versus ergonomic instructions at work (24), and comparison of different types of exercises between intervention groups (34, 36). This procedure is likely to minimize the effects of placebo on the outcomes, as all groups receive an intervention that possibly may be beneficial. Another method used for implementation and evaluation of interventions which cannot be blinded is stepped wedged design (58). Here, cluster randomizing of teams or workplaces as a whole together with gradual offering of interventions to the included clusters enables comparison with the participants as their own controls. Thøgersen et al (45) conducts a small-scale similar method by using delayed start of intervention for their reference group.

Concluding remarks

Workplace exercise seem to have limited effects on mental health and the psychosocial work environment. The great variety across populations, types of interventions, and outcome measures in the included trials in this review hampers possibilities to draw conclusions across studies. Evaluating effects of behavioral workplace interventions is challenging due to difficulties with blinding and independent outcome measure. Alternative designs should be prioritized in the future.

Conflict of interest, funding, and registration

The authors declare no conflicts of interest. This study received no external funding. The protocol of this review was registered and is accessible in the Prospero register (www.crd.york.ac.uk/prospero) ID: CRD42018099556.

Acknowledgements

Thanks to Elizabeth Bengtsen, librarian at the National Research Centre for the Working Environment for providing valuable feedback on the search strings and help and guidance on the project overall.

References

- Andreasen M, Falk J, Nielsen NS. Sundhed og trivsel på arbejdspladsen– Inspiration til systematisk og strategisk arbejde med sundhedsfremme i virksomheden [Health and well-being among workers – Inspiration to systematic and strategic health promotion in workplaces]. Copenhagen: Danish Health Authority; 2010. Report No.: 978-87-7104-027-2.
- Sjøl A, Thomsen KK, Schroll M, Andersen LB. Secular trends in acute myocardial infarction in relation to physical activity in the general Danish population. *Scand J Med Sci Sports* 2003 Aug;13(4):224–30. <https://doi.org/10.1034/j.1600-0838.2003.00310.x>.
- Christensen AI, Severin M, Holmberg T, Eriksen L, Toftager M, Zachariassen A et al. KRAM-undersøgelsen i tal og billeder, Statens Institut for Folkesundhed [The Survey of Lifestyle Factors – in numbers and graphics]. Copenhagen: The National Institute of Public Health, the University of Southern Denmark; 2009. Report No.: 978-87-992261-2-2
- van der Ploeg HP, Møller SV, Hannerz H, van der Beek AJ, Holtermann A. Temporal changes in occupational sitting time in the Danish workforce and associations with all-cause mortality: results from the Danish work environment cohort study. *Int J Behav Nutr Phys Act* 2015 Jun;12:71. <https://doi.org/10.1186/s12966-015-0233-1>.
- World Health Organization W. Obesity. preventing and managing the global epidemic. Geneva: WHO; 2000. Report No.: 92 4 120894 5.
- van Uffelen JG, Wong J, Chau JY, van der Ploeg HP, Riphagen I, Gilson ND et al. Occupational sitting and health risks: a systematic review. *Am J Prev Med* 2010 Oct;39(4):379–88. <https://doi.org/10.1016/j.amepre.2010.05.024>.
- Andersen MF, Nielsen KM, Brinkmann S. Meta-synthesis of qualitative research on return to work among employees with common mental disorders. *Scand J Work Environ Health* 2012 Mar;38(2):93–104. <https://doi.org/10.5271/sjweh.3257>.
- Broughton A. Work-related stress Dublin: European Foundation for the Improvement of Living and Working Conditions; 2010.
- Arbejdsmiljø DNFF. Fakta om Arbejdsmiljø & Helbred 2016 [The Working Environment in Denmark 2016]. Copenhagen: The National Research Centre for the Working Environment; 2016.
- US Department of Health and Human Services. Physical Activity and Health: A Report of the Surgeon General. Atlanta, GA, U.S.: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion; 1996.
- Bauman AE. Updating the evidence that physical activity is good for health: an epidemiological review 2000-2003. *J Sci Med Sport* 2004 Apr;7(1 Suppl):6–19. [https://doi.org/10.1016/S1440-2440\(04\)80273-1](https://doi.org/10.1016/S1440-2440(04)80273-1).
- Mammen G, Faulkner G. Physical activity and the prevention of depression: a systematic review of prospective studies. *Am J Prev Med* 2013 Nov;45(5):649–57. <https://doi.org/10.1016/j.amepre.2013.08.001>.
- Salmon P. Effects of physical exercise on anxiety, depression, and sensitivity to stress: a unifying theory. *Clin Psychol Rev* 2001 Feb;21(1):33–61. [https://doi.org/10.1016/S0272-7358\(99\)00032-X](https://doi.org/10.1016/S0272-7358(99)00032-X).
- Hassmén P, Koivula N, Uutela A. Physical exercise and psychological well-being: a population study in Finland. *Prev Med* 2000 Jan;30(1):17–25. <https://doi.org/10.1006/pmed.1999.0597>.
- De Mello MT, Lemos VA, Antunes HK, Bittencourt L, Santos-Silva R, Tufik S. Relationship between physical activity and depression and anxiety symptoms: a population study. *J Affect Disord* 2013 Jul;149(1-3):241–6. <https://doi.org/10.1016/j.jad.2013.01.035>.
- Schnohr P, Kristensen TS, Prescott E, Scharling H. Stress and life dissatisfaction are inversely associated with jogging and other types of physical activity in leisure time--The Copenhagen City Heart Study. *Scand J Med Sci Sports* 2005 Apr;15(2):107–12. <https://doi.org/10.1111/j.1600-0838.2004.00394.x>.
- Proper KI, Koning M, van der Beek AJ, Hildebrandt VH, Bosscher RJ, van Mechelen W. The effectiveness of worksite physical activity programs on physical activity, physical fitness, and health. *Clinical journal of sport medicine: official journal of the Canadian Academy of Sport Medicine*. 2003;13(2):106–17.
- Dias HJ. Evaluation of the effectiveness of workplace exercise in controlling neck, shoulder and low back pain: a systematic review. *Braz J Phys Ther* 2009;13(6). <http://dx.doi.org/10.1590/S1413-35552009000600002>.
- Conn VS, Hafdahl AR, Cooper PS, Brown LM, Lusk SL. Meta-analysis of workplace physical activity interventions. *Am J Prev Med* 2009 Oct;37(4):330–9. <https://doi.org/10.1016/j.amepre.2009.06.008>.
- Korshej M, Holtermann A. Fysiske Arbejdskrav og Fitness – Betydning for hjertekarsygdom [Physical Work demands and Fitness – Effects on risk factors for cardiovascular disease]. Copenhagen: The National Research Centre for the Working Environment; 2015. Report No.: 978-87-7904-289-6 20.
- Chu AH, Koh D, Moy FM, Müller-Riemenschneider F. Do workplace physical activity interventions improve mental health outcomes? *Occup Med (Lond)* 2014 Jun;64(4):235–45. <https://doi.org/10.1093/occmed/kqu045>.
- Justesen JB, Søgaard K, Dalager T, Christensen JR, Sjøgaard

- G. The Effect of Intelligent Physical Exercise Training on Sickness Presenteeism and Absenteeism Among Office Workers. *J Occup Environ Med* 2017 Oct;59(10):942–8. <https://doi.org/10.1097/JOM.0000000000001101>.
23. Lidegaard M, Søgaard K, Krusturup P, Holtermann A, Korshøj M. Effects of 12 months aerobic exercise intervention on work ability, need for recovery, productivity and rating of exertion among cleaners: a worksite RCT. *Int Arch Occup Environ Health* 2018 Feb;91(2):225–35. <https://doi.org/10.1007/s00420-017-1274-3>.
24. Andersen LL, Persson R, Jakobsen MD, Sundstrup E. Psychosocial effects of workplace physical exercise among workers with chronic pain: randomized controlled trial. *Medicine (Baltimore)* 2017 Jan;96(1):e5709. <https://doi.org/10.1097/MD.0000000000005709>.
25. Jakobsen MD, Sundstrup E, Brandt M, Andersen LL. Psychosocial benefits of workplace physical exercise: cluster randomized controlled trial. *BMC Public Health* 2017 Oct;17(1):798. <https://doi.org/10.1186/s12889-017-4728-3>.
26. Aas RW, Tuntland H, Holte KA, Røe C, Lund T, Marklund S et al. Workplace interventions for neck pain in workers. *Cochrane Database Syst Rev* 2011 Apr;(4):CD008160.
27. Stansfeld S, Candy B. Psychosocial work environment and mental health—a meta-analytic review. *Scand J Work Environ Health* 2006 Dec;32(6):443–62. <https://doi.org/10.5271/sjweh.1050>.
28. Johnson JV, Hall EM. Job strain, work place social support, and cardiovascular disease: a cross-sectional study of a random sample of the Swedish working population. *Am J Public Health* 1988 Oct;78(10):1336–42. <https://doi.org/10.2105/AJPH.78.10.1336>.
29. Karasek RA. Job Demands, Job Decision Latitude, and Mental Strain: Implications for Job Redesign. *Adm Sci Q* 1979;24(2):285–308. <https://doi.org/10.2307/2392498>.
30. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009 Jul;6(7):e1000097. <https://doi.org/10.1371/journal.pmed.1000097>.
31. Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD et al.; Cochrane Bias Methods Group; Cochrane Statistical Methods Group. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011 Oct;343:d5928. <https://doi.org/10.1136/bmj.d5928>.
32. Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011 Apr;64(4):383–94. <https://doi.org/10.1016/j.jclinepi.2010.04.026>.
33. Andersen LL, Poulsen OM, Sundstrup E, Brandt M, Jay K, Clausen T et al. Effect of physical exercise on workplace social capital: cluster randomized controlled trial. *Scand J Public Health* 2015 Dec;43(8):810–8. <https://doi.org/10.1177/1403494815598404>.
34. Brown DK, Barton JL, Pretty J, Gladwell VF. Walks4Work: assessing the role of the natural environment in a workplace physical activity intervention. *Scand J Work Environ Health* 2014 Jul;40(4):390–9. <https://doi.org/10.5271/sjweh.3421>.
35. De Zeeuw E, Tak E, Dusseldorp E, Hendriksen IJ. Workplace exercise intervention to prevent depression: a pilot randomized controlled trial. *Ment Health Phys Act* 2010;3(2):72–7. <https://doi.org/10.1016/j.mhpa.2010.09.002>.
36. Eriksen HR, Ihlebaek C, Mikkelsen A, Grønningsaeter H, Sandal GM, Ursin H. Improving subjective health at the worksite: a randomized controlled trial of stress management training, physical exercise and an integrated health programme. *Occup Med (Lond)*. 2002 Oct;52(7):383–91. <https://doi.org/10.1093/occmed/52.7.383>.
37. Hartfiel N, Burton C, Rycroft-Malone J, Clarke G, Havenhand J, Khalsa SB et al. Yoga for reducing perceived stress and back pain at work. *Occup Med (Lond)* 2012 Dec;62(8):606–12. <https://doi.org/10.1093/occmed/kqs168>.
38. Hartfiel N, Havenhand J, Khalsa SB, Clarke G, Krayner A. The effectiveness of yoga for the improvement of well-being and resilience to stress in the workplace. *Scand J Work Environ Health*. 2011 Jan;37(1):70–6. <https://doi.org/10.5271/sjweh.2916>.
39. Jay K, Brandt M, Hansen K, Sundstrup E, Jakobsen MD, Schraefel MC et al. Effect of Individually Tailored Biopsychosocial Workplace Interventions on Chronic Musculoskeletal Pain and Stress Among Laboratory Technicians: Randomized Controlled Trial. *Pain Physician* 2015;18(5):459–71.
40. Mesquita CC, Ribeiro JC, Moreira P. An Exercise Program Improves Health-Related Quality of Life of Workers. *Appl Res Qual Life* 2012;7(3):295–307. <https://doi.org/10.1007/s11482-011-9161-7>.
41. Puig-Ribera A, Bort-Roig J, Giné-Garriga M, González-Suárez AM, Martínez-Lemos I, Fortuño J et al. Impact of a workplace 'sit less, move more' program on efficiency-related outcomes of office employees. *BMC Public Health* 2017 May;17(1):455. <https://doi.org/10.1186/s12889-017-4367-8>.
42. Roessler KK, Rugulies R, Bilberg R, Andersen LL, Zebis MK, Sjøgaard G. Does work-site physical activity improve self-reported psychosocial workplace factors and job satisfaction? A randomized controlled intervention study. *Int Arch Occup Environ Health* 2013 Nov;86(8):861–4. <https://doi.org/10.1007/s00420-012-0823-z>.
43. Sjögren T, Nissinen KJ, Järvenpää SK, Ojanen MT, Vanharanta H, Mäkiä EA. Effects of a physical exercise intervention on subjective physical well-being, psychosocial functioning and general well-being among office workers: a cluster randomized-controlled cross-over design. *Scand J Med Sci Sports* 2006 Dec;16(6):381–90. <https://doi.org/10.1111/j.1600-0838.2005.00516.x>.
44. Strijk JE, Proper KI, van der Beek AJ, van Mechelen W. A worksite vitality intervention to improve older

- workers' lifestyle and vitality-related outcomes: results of a randomised controlled trial. *J Epidemiol Community Health* 2012 Nov;66(11):1071–8. <https://doi.org/10.1136/jech-2011-200626>.
45. Thøgersen-Ntoumani C, Loughren EA, Kinnafick FE, Taylor IM, Duda JL, Fox KR. Changes in work affect in response to lunchtime walking in previously physically inactive employees: A randomized trial. *Scand J Med Sci Sports* 2015 Dec;25(6):778–87. <https://doi.org/10.1111/sms.12398>.
 46. Tveito TH, Eriksen HR. Integrated health programme: a workplace randomized controlled trial. *J Adv Nurs* 2009 Jan;65(1):110–9. <https://doi.org/10.1111/j.1365-2648.2008.04846.x>.
 47. Van Rhenen W, Blonk RW, van der Klink JJ, van Dijk FJ, Schaufeli WB. The effect of a cognitive and a physical stress-reducing programme on psychological complaints. *Int Arch Occup Environ Health* 2005 Mar;78(2):139–48. <https://doi.org/10.1007/s00420-004-0566-6>.
 48. Chopp-Hurley JN, Brennenman EC, Wiebenga EG, Bulbrook B, Keir PJ, Maly MR. Randomized Controlled Trial Investigating the Role of Exercise in the Workplace to Improve Work Ability, Performance, and Patient-Reported Symptoms Among Older Workers With Osteoarthritis. *J Occup Environ Med* 2017 Jun;59(6):550–6. <https://doi.org/10.1097/JOM.0000000000001020>.
 49. Taylor WC, Paxton RJ, Shegog R, Coan SP, Dubin A, Page TF et al. Impact of Booster Breaks and Computer Prompts on Physical Activity and Sedentary Behavior Among Desk-Based Workers: A Cluster-Randomized Controlled Trial. *Prev Chronic Dis* 2016 Nov;13:E155. <https://doi.org/10.5888/pcd13.160231>.
 50. Matsugaki R, Kuhara S, Saeki S, Jiang Y, Michishita R, Ohta M et al. Effectiveness of workplace exercise supervised by a physical therapist among nurses conducting shift work: A randomized controlled trial. *J Occup Health* 2017 Jul;59(4):327–35. <https://doi.org/10.1539/joh.16-0125-OA>.
 51. Michishita R, Jiang Y, Ariyoshi D, Yoshida M, Moriyama H, Yamato H. The practice of active rest by workplace units improves personal relationships, mental health, and physical activity among workers. *J Occup Health* 2017 Mar;59(2):122–30. <https://doi.org/10.1539/joh.16-0182-OA>.
 52. Cheema BS, Houridis A, Busch L, Raschke-Cheema V, Melville GW, Marshall PW et al. Effect of an office worksite-based yoga program on heart rate variability: outcomes of a randomized controlled trial. *BMC Complement Altern Med* 2013 Apr;13:82. <https://doi.org/10.1186/1472-6882-13-82>.
 53. World Health Organization W. Workplace health promotion [Available from: http://www.who.int/occupational_health/topics/workplace/en/index1.html].
 54. Sundhedsstyrelsen S. Muskelskelettlidelser 2017 [Available from: <https://www.sst.dk/da/sygdom-og-behandling/kronisk-sygdom/muskelskelettlidelser>].
 55. Arbejdsmiljø V. Daglige øvelser skaber sundhed og arbejdsglæde 2018 [Available from: <https://amid.dk/inspiration-fra-andre/msb/motion-i-arbejdstiden-en-investering/>].
 56. Mattke S, Liu H, Caloyeras J, Huang CY, Van Busum KR, Khodyakov D et al. Workplace Wellness Programs Study: final Report. *Rand Health Q* 2013 Jun;3(2):7.
 57. Meng L, Wolff MB, Mattick KA, DeJoy DM, Wilson MG, Smith ML. Strategies for Worksite Health Interventions to Employees with Elevated Risk of Chronic Diseases. *Saf Health Work* 2017 Jun;8(2):117–29. <https://doi.org/10.1016/j.shaw.2016.11.004>.
 58. Mdege ND, Man MS, Taylor Nee Brown CA, Torgerson DJ. Systematic review of stepped wedge cluster randomized trials shows that design is particularly used to evaluate interventions during routine implementation. *J Clin Epidemiol* 2011 Sep;64(9):936–48. <https://doi.org/10.1016/j.jclinepi.2010.12.003>.

Received for publication: 2 January 2019