Scand J Work Environ Health 2021;47(8):609-618

<https://doi.org/10.5271/sjweh.3979>

Published online: 16 Aug 2021, Issue date: 01 Nov 2021

Are resident handlings in eldercare wards associated with musculoskeletal pain and sickness absence among the workers? A prospective study based on onsite observations

by [Januario LB](#), [Mathiassen SE](#), [Stevens ML](#), [Holtermann A](#), [Bergström G](#), [Rugulies R](#), [Karstad K](#), [Hallman DM](#)

This is the first multi-level study to use onsite observations of resident handlings in a latent profile analysis, interpret those in terms of ward 'phenotypes', and address the prospective association of phenotypes with musculoskeletal pain and sickness absence among eldercare workers. Our findings indicate that combined organizational factors, e.g. frequent handlings and use of devices, are important determinants of musculoskeletal health.

Affiliation: Department of Occupational Health Sciences and Psychology, Centre for Musculoskeletal Research, University of Gävle Kungsbäcksvägen 47, 801 76, Gävle, Sweden. leticia.januario@hig.se

Refers to the following texts of the Journal: [2017;43\(6\):526-539](#)
[2013;39\(2\):164-169](#) [2012;38\(6\):582-589](#)

The following article refers to this text: [0;0 Special issue:0](#)

Key terms: [eldercare](#); [healthcare](#); [latent profile analysis](#); [longitudinal study](#); [musculoskeletal pain](#); [patient handling](#); [prospective study](#); [resident](#); [sickness absence](#)

This article in PubMed: www.ncbi.nlm.nih.gov/pubmed/34397097

Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](#).

This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Are resident handlings in eldercare wards associated with musculoskeletal pain and sickness absence among the workers? A prospective study based on onsite observations

by Leticia Bergamin Januario, PhD,¹ Svend Erik Mathiassen PhD,¹ Matthew L Stevens, PhD,² Andreas Holtermann, PhD,² Gunnar Bergström, PhD,^{1,3} Reiner Rugulies, PhD,^{2,4,5} Kristina Karstad, PhD,² David M Hallman, PhD¹

Januario LB, Mathiassen SE, Stevens ML, Holtermann A, Bergström G, Rugulies R, Karstad K, Hallman DM. Are resident handlings in eldercare wards associated with musculoskeletal pain and sickness absence among the workers? A prospective study based on onsite observations. *Scand J Work Environ Health*. 2021;47(8):609–618. doi:10.5271/sjweh.3979

Objectives We aimed to identify eldercare wards with different types of resident handling characteristics ('phenotypes') and determine the prospective association between these characteristics and musculoskeletal pain and sickness absence among workers during a one-year follow-up.

Methods Our study was based on the DOSES cohort, including 467 workers at 103 eldercare wards. At baseline, resident handlings were assessed using onsite observations. Workers' self-reported musculoskeletal pain and sickness absence were assessed during the following year using text messages. Observations of the frequency of handlings per shift, use of assistive devices, assistance from others, and barriers (interruptions and impediments) were estimated for each worker, aggregated at ward level, and entered into a latent profile analysis, identifying ward phenotypes. We then used generalized estimating equations to determine associations between ward phenotypes, musculoskeletal pain and sickness absence.

Results We identified four ward phenotypes: 'turbulent' (many handlings with devices and assistance, many barriers), 'strained' (many handlings without devices or assistance, some barriers), 'unpressured' (few handlings, yet without devices or assistance, few barriers) and 'balanced' (some handlings with devices and assistance, some barriers). Compared to workers in balanced wards, workers in turbulent wards had more days with neck-shoulder and low-back pain (LBP); and those working in strained wards had more days with LBP and higher pain intensities.

Conclusion We found that ward phenotypes based on resident handling characteristics were predictive of musculoskeletal pain and sickness absence over one year. This shows that organizational factors related to resident handling are important determinants of musculoskeletal health among eldercare workers.

Key terms healthcare; latent profile analysis; longitudinal study; patient handling.

Eldercare workers often report low-back pain (LBP), neck-shoulder pain (NSP) and musculoskeletal sickness absence, likely due to high physical workload and poor psychosocial working conditions (1–3). This is not only an issue for current workers, it also impedes recruitment of a new workforce required to look after the increased number of elderly in need of care in Europe (4, 5).

As proposed in previous exposure–effect models (6, 7), musculoskeletal health at work is determined in a complex interplay between factors pertaining to tasks, jobs, and the work environment (external exposures),

and other factors influencing the eventual mechanical load on individual workers (internal exposures such as postures, movement and forces). In eldercare work, resident handlings are a key external exposure in the development of musculoskeletal disorders (8, 9) since they lead to complex and high biomechanical demands, including pronounced arm elevation, trunk flexion and trunk rotation, often while handling large loads (10, 11). This external exposure to handlings is, in turn, influenced by surrounding organizational factors setting conditions for performing the handlings (6). Examples

¹ Department of Occupational Health Sciences and Psychology, Centre for Musculoskeletal Research, University of Gävle, Gävle, Sweden.

² National Research Centre for the Working Environment, Copenhagen, Denmark.

³ Unit of Intervention and Implementation Research for Worker Health, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden.

⁴ Department of Public Health, University of Copenhagen, Copenhagen, Denmark.

⁵ Department of Psychology, University of Copenhagen, Copenhagen, Denmark.

Correspondence to: Leticia Bergamin Januario, Department of Occupational Health Sciences and Psychology, Centre for Musculoskeletal Research, University of Gävle, Kungsbäcksvägen 47, 801 76, Gävle, Sweden. [E-mail: leticia.januario@hig.se]

are under-staffed wards, shortage of time to perform tasks as prescribed, and broken assistive devices, all of which can result in compromised resident handling. The resulting internal exposure on the individual worker gives rise to acute responses (eg, increases in oxygen consumption, heart rate, and perceived exertion), which may, in the long term, have effects on musculoskeletal health (6). Several aspects of handlings – such as their frequency, possible interruptions, and use (or not) of assistive devices – may influence the eventual internal exposure and thus musculoskeletal health (11). For instance, failure to follow handling guidelines can negatively influence musculoskeletal health (12–14). Since handling aspects occur in parallel, they should be addressed together, instead of one at a time, as in previous eldercare studies (15, 16).

To a major extent, previous studies in eldercare have also been based on self-reported data on physical workload, which are prone to bias (17). As an alternative, observational exposure assessment tools have been developed to measure physical workloads during caretaking (18–21). However, to our knowledge, onsite observations have not been applied in any study addressing effects on health of combined handlings characteristics in eldercare.

Eldercare homes are often organized in separate wards, differing in organizational, managerial and structural characteristics that likely influence resident handlings and consequently the musculoskeletal health of workers at the wards. Therefore, examining handlings at a ward level would allow a better understanding of the organizational determinants of exposures, which, in turn, could lead to evidence-based interventions at that level, eg, addressing staff ratio or the availability of assistive devices. Such interventions would likely be more effective and sustainable than initiatives focusing solely on the behaviors of the individual worker (eg, personal training programs), which are known to have limited effectiveness (22). In the present study, we combine the need to address several handling aspects in parallel with the idea of acting at the ward level by determining latent sub-populations of wards sharing similar resident handling characteristics, using latent profile analysis. We label these sub-populations as ward ‘phenotypes’, referring to the term used in genetics for *a set of observable characteristics of an organism* (23).

This study aims to identify phenotypes of wards sharing similar physical and psychosocial conditions for resident handlings and then determine the extent to which they are associated with musculoskeletal pain and sickness absence among workers at the wards over the course of one year.

Methods

Design and study population

This prospective study uses data from the Danish Observational Study of Eldercare work and musculoskeletal disorders (DOSES), collected between September 2013 and January 2016 and previously described (24). Briefly, DOSES is a prospective study, in which data at multiple organizational levels were considered. In total 553 eldercare workers, employed in 126 wards nested in 20 eldercare homes, were included with the following inclusion criteria: aged 18–65 years; employed >15 hours/week; working on day, evening or rotating shifts (ie, not exclusively at night); and spending $\geq 25\%$ of their working time in tasks associated with actual resident care. The Danish Data Protection Agency and the regional Ethics Committee in Copenhagen, Denmark, (H-4-2013-028) approved the study and all participants provided written informed consent to participate. Baseline data collection included questionnaires and onsite observations. During a one-year follow-up, data on self-reported musculoskeletal pain and sickness absence were collected. In our subsequent analyses, we excluded workers who refused to be observed, did not report their work schedules and/or did not answer the questions related to musculoskeletal pain and/or sickness absence at baseline and/or follow-up. To ensure that handlings were representative of wards, we only included wards with ≥ 2 participating workers. Eventually, the present study included 467 workers distributed across 103 wards in 20 eldercare homes.

Measurements

Descriptive characteristics. At baseline, ward managers were asked to complete a questionnaire addressing conditions at their ward. These ward managers were, in general, responsible for communicating with higher-level management, taking decisions at their own ward, and managing the staff there. The questionnaire asked about type of ward (somatic, dementia, rehabilitation or psychiatric unit), resident information (ie, number of residents in the wards; weight, physical and psychosocial function level of each resident), staff ratio, and how often assistive devices were not in place (daily, 1–3 times/week, 2 times/month or ≤ 1 time/month). Physical function level was categorized based on the resident’s need for physical assistance (light, moderate, extensive or complete need). Psychosocial function level was categorized based on the resident’s usual behavior (mostly neutral, mostly positive/appreciative, mostly resistant, and mostly aggressive).

The eldercare workers answered a baseline questionnaire asking about age, smoking habits (smoker or

non-smoker) and musculoskeletal pain and sickness absence due to NSP and/or LBP, as described in detail below. Body mass index (BMI) was calculated based on measured height and weight (kg/m^2).

Observations – resident handling characteristics. Handling tasks were observed using the DOSES observation instrument, shown previously to have a good reliability (20). At any particular ward, trained research assistants followed workers for 4 hours during day shifts and 4–5 hours during evening shifts, recording, to the extent possible, all handlings of all residents on a tablet, using the Noldus Observer XT 11 software (Noldus, Wageningen, The Netherlands). A previous observational study in Danish eldercare homes (21) showed that more than 70% of all handlings occurred during these periods and thus, to save resources, observations in DOSES were limited to these periods.

The observation instrument was composed of 26 items in four sections: setting and surroundings, manual handling activities, psychosocial interactions with the residents, and barriers (20). Observations of manual handling activities and of barriers were included in the subsequent identification of ward phenotypes. Manual handling activities were classified in terms of whether the worker used an assistive device or not (including the type of device used) and whether the handling was performed by the worker alone, or with the assistance of another person (typically a co-worker or a visitor). Barriers included interruptions and impediments (25) during the care of the resident. An interruption was defined as an event that significantly interrupted the eldercare worker in performing a task (eg, an urgent request for assistance from a co-worker); and an impediment was defined as an obstacle for completing a task requiring some effort (eg, broken or missing assistive device).

Each single observation was registered as an exposure associated with the observed resident. Thus, some residents had more registered observations than others because they were in need of more handlings during a shift. A majority of residents in the 20 eldercare homes were observed (88% in day shifts and 79% in evening shifts). Reasons for not observing a resident could be that (i) the resident was not in need of care or was hospitalized, (ii) the eldercare worker taking care of that resident declined to be followed, and (iii) there was a shortage of research assistants at that point in time. Estimates of handling characteristics for non-observed residents were obtained by imputation. Next, handlings at the level of individual eldercare workers were estimated based on work schedules, which were obtained from the ward managers for all workers, irrespective of whether they participated in observations or not. Schedules were then combined with the data based on handlings at the resident level to give a task-based estimate of the num-

ber of handlings per shift for each worker, as well as the associated handling characteristics. In our eventual analysis of phenotypes, we considered (i) the frequency of handlings per shift, and calculated (ii), the proportion of those handlings done without any assistive device and (iii), the proportion of handlings done without any assistance from others. We also considered the number of (iv) interruptions per shift and (v) impediments per shift. We aggregated observations of these five variables at the ward level by averaging the handling exposure estimates of all eldercare workers in each specific ward. In this study, observations during day and evening shifts were summarized in terms of values per shift, disregarding time of day. Then, we used these ward-level scores in a latent profile analysis of handling characteristics.

Musculoskeletal pain and sickness absence. The eldercare workers reported musculoskeletal pain and sickness absence at baseline by answering a questionnaire, and during the one-year follow-up by answering, at the most, six questions in text messages (www.sms-track.com) (24, 26). At baseline and every four weeks during follow-up, the workers were asked to report the number of days they experienced NSP and LBP during the preceding four weeks (0–28 days). If participants reported any pain in one or both of these regions, they received a follow-up question about the intensity of their NSP and/or LBP (scale 0–10), and the number of days (0–28) when pain interfered with work (“Within the past four weeks, how many days did pain in your low-back and/or neck-shoulder made it difficult to perform your normal work, ie, interfered/limited your performance at work?”).

At baseline, and every 12 weeks in follow-up, the workers were also asked to report the number of days on sickness absence due to NSP and/or LBP in the past three months (0–84 days). Answers to pain-related work interference were dichotomized into 0 or ≥ 1 days, while the other two variables describing number of days were kept continuous.

Statistical analysis

Based on the handling characteristics of each of the 103 wards (frequency of handlings, proportion of handlings done without assistive devices, proportion of handlings done without assistance from others, number of interruptions, and number of impediments), we performed a latent profile analysis, using the Latent Gold software (version 5.1, Statistical innovations, Belmont, MA, USA). This technique allows multiple variables to be combined in an analysis identifying latent profiles, ie, subpopulations within a population sharing similar properties with respect to the set of observed variables (27–29). In our case, the latent profile analysis was used

to identify wards sharing handling characteristics, as a basis for identifying ‘phenotypes’ of wards (see below), which could then be examined for their ability to predict musculoskeletal pain and sickness absence.

The latent profile analysis requires the user to decide the number of profiles that suits data the best. In this process, we considered model fit indices as well as the interpretability of identified profiles. Model fit statistics included the Akaike information criterion (AIC) and the Bayesian information criterion (BIC), showing the trade-off between the simplicity of the model and the goodness of fit, and entropy, describing the degree of certainty of the classification. We also required each profile to contain at least five wards so that even the smallest profile would be generalizable. To better understand the distinct handling characteristics of the eventually identified profiles, we compared the characteristics of wards, workers and residents in each profile, using analysis of variance (ANOVA) for descriptive continuous variables and Chi-squared tests for descriptive categorical variables. At this point, and after further examining the specific ward characteristics expressed in each profile, we labeled the ward profiles as phenotypes, in the sense that they reflect a set of observable and interpretable characteristics pertaining to the ward.

To estimate the effects of working at a certain ward phenotype (ward-level predictor) on reported musculoskeletal pain and sickness absence during the one-year follow-up (individual-level outcome) we used generalized estimating equations (GEE) (30). In these analyses, each worker was assigned the ward phenotype in which they worked as a predictor, and their self-reported musculoskeletal pain and sickness absence at each time point during follow-up as the outcome (14 time points for pain and 5 time points for sickness absence). We included time as a within-subject covariate, and workers and wards as random effects in an unstructured correlation matrix. For the continuous outcome variables (days with NSP, intensity NSP, days with LBP, intensity LBP, and sickness absence due to NSP/LBP), we used a Poisson distribution model, with a log link function. For the dichotomous variable (pain-related work interference), we used a binary logistic model.

We adjusted each of the six models for their corresponding outcome values obtained at baseline. We further adjusted the models for age, BMI and smoking habits. We also did a sensitivity analysis by re-running each of the six primary models solely with participants reporting ‘little’ pain at baseline, defined as pain ≤ 5 days/month, pain intensity ≤ 2 (0–10 scale), 0 days with pain-related work interference, and 0 days with sickness absence. We performed all tests in SPSS (Statistical Package for Social Science, v. 20, IBM Corp, Armonk, NY, USA) and set the significance level at 0.05.

Results

Identification and characterization of ward phenotypes

According to the model fit statistics (supplementary material, <https://www.sjweh.fi/article/3979>, table S1), models with four and five profiles were the more certain (higher entropy values), while still being sufficiently parsimonious according to BIC and AIC values. These two models also fulfilled the criterion of the smallest profile containing at least five wards. Taking into account that the model with four profiles showed better parsimony and had more workers in the smallest profile than the model with five profiles, and that the differences in the handling characteristics between the four ward profiles were more intuitive than those in the five-profiles model, we settled on the four-profile solution.

We then examined the four ward profiles (see figure 1) and described their properties in terms of four phenotypes with the following labels:

‘Turbulent’ wards: many handling operations (mean 11.8, standard deviation (SD) 7.1 per shift), some of which are done without assistive devices [mean 41 (SD 18) %] or assistance from others [mean 20 (SD 9) %], and with many barriers [mean 6.2 (SD 2.5) interruptions per shift; mean 2.7 (SD 1.8) impediments per shift].

‘Strained’ wards: many handling operations [mean 11.4 (SD 4.3) per shift], many of which are done without assistive devices [mean 52 (SD 18) %] or assistance from others [mean 67 (SD 10) %], with some barriers [mean 3.1 (SD 1.8) interruptions per shift; mean 1.9 (SD 1.3) impediments per shift].

‘Unpressured’ wards: few handling operations [mean 4.9 (SD 1.8) per shift], even if many of which are done without assistive devices [mean 53 (SD 21) %] or assistance from others [mean 61 (SD 24) %], and few barriers [mean 1.7 (SD 0.8) interruptions per shift; mean 1.5 (SD 0.7) impediments per shift].

‘Balanced’ wards: some handling operations [mean 9.2 (SD 1.6) per shift], some of which without assistive devices [mean 47 (SD 12) %] or assistance from others [mean 34 (SD 11) %], and with some barriers [mean 2.9 (SD 1.0) interruption per shift; mean 2.1 (SD 0.7) impediments per shift].

Based on the latent profile analysis, mean 7 (6.8%), 21 (20.4%), 52 (50.5%) and 23 (22.3%) of the 103 wards were categorized as turbulent, strained, unpressured, and balanced, respectively. Table 1 presents descriptive statistics on wards, workers and residents, stratified by ward phenotype.

At turbulent wards, the staff ratio was smaller than at the other ward phenotypes and a higher proportion of residents were classified as needing complete physical assistance. At strained wards, devices were reported to be misplaced daily more often than in the other ward

Figure 1. Standardized values (Z-scores) of number of handlings per shift, proportion of handlings without assistive devices (proportion w/o devices), proportion of handlings without assistance from co-workers or others (prop. w/o assistance), number of interruptions per shift and number of impediments per shift, in each ward phenotype. Each phenotype was identified using a latent profile analysis based on observed handling characteristics at the ward level. Positive values in the y-axis represent negative attributes, ie, higher number of handlings, higher proportion of handlings done without devices, higher proportion of handlings done without help and higher rate of interruptions and impediments per shift.

Table 1. Descriptive characteristics at ward, worker and resident level for each ward phenotype (N=103), as identified using latent profile analysis of observed handling characteristics, and P-values for differences between ward phenotypes. Data based on the baseline questionnaire. [SD=standard deviation.]

	Total		Turbulent wards		Strained wards		Unpressured wards		Balanced wards		P-value
	Mean (SD)	N (%)	Mean (SD)	N (%)	Mean (SD)	N (%)	Mean (SD)	N (%)	Mean (SD)	N (%)	
Ward characteristics											
Staff ratio	0.46 (0.10)		0.42 (0.16)		0.44 (0.08)		0.47 (0.12)		0.47 (0.07)		0.01
Type of ward											<0.01
Somatic		77 (76.2)		6 (85.7)		21 (100.0)		36 (70.6)		14 (63.6)	
Dementia		20 (19.6)		1 (14.3)		0 (0.0)		13 (25.5)		6 (27.3)	
Rehabilitation		2 (1.9)		0 (0.0)		0 (0.0)		2 (3.9)		0 (0.0)	
Psychiatric		2 (1.9)		0 (0.0)		0 (0.0)		0 (0.0)		2 (9.1)	
Devices not in place											<0.01
Daily		9 (11.8)		0 (0.0)		6 (31.6)		1 (2.7)		2 (11.8)	
1–3 times/week		2 (2.6)		0 (0.0)		0 (0.0)		2 (5.4)		0 (0.0)	
2 times/month		13 (17.1)		1 (33.3)		3 (15.8)		8 (21.6)		1 (5.9)	
≤1 time/month		51 (68.0)		2 (66.7)		10 (52.6)		26 (70.3)		14 (82.4)	
Worker characteristics											
Number of workers		467 (100.0)		23 (4.9)		74 (15.8)		251 (53.7)		119 (25.5)	
Age (years)	45.5 (10.7)		45.0 (10.6)		44.5 (11.5)		45.1 (10.3)		46.8 (11.3)		0.38
Body mass index (kg/m ²)	26.4 (5.3)		25.0 (5.1)		27.8 (5.7)		26.5 (5.3)		25.5 (4.7)		0.01
Smoking habits											0.46
Smokers		135 (28.9)		8 (34.8)		22 (29.7)		71 (28.3)		34 (28.6)	
Non-smokers		332 (71.1)		15 (65.2)		52 (70.3)		180 (71.7)		85 (71.4)	
Resident characteristics											
Number of residents		1112 (100.0)		76 (6.8)		212 (19.1)		585 (52.6)		239 (21.5)	
Weight (kg) ^a	65.6 (5.2)		64.1 (2.3)		65.2 (5.0)		65.5 (4.1)		66.7 (7.9)		0.87
Physical function level ^a											<0.01
Light		243 (21.9)		18 (23.7)		36 (17.0)		140 (23.9)		49 (20.5)	
Moderate		379 (34.1)		21 (27.6)		55 (25.9)		218 (37.3)		85 (35.6)	
Extensive		222 (20.0)		19 (25.0)		53 (25.0)		106 (18.1)		44 (18.4)	
Completely		268 (24.1)		18 (23.7)		68 (32.1)		121 (20.7)		61 (25.5)	
Psychosocial function level ^a											0.30
Mostly neutral		313 (28.3)		22 (28.9)		73 (35.1)		162 (27.7)		56 (23.4)	
Mostly appreciative		429 (38.8)		26 (34.2)		78 (37.5)		224 (38.4)		101 (42.3)	
Mostly resistant		275 (24.8)		22 (28.9)		42 (20.2)		146 (25.0)		65 (27.2)	
Mostly aggressive		90 (8.1)		6 (7.9)		15 (7.2)		52 (8.9)		17 (7.1)	

^a Information provided by team managers. Bold values show significant differences between phenotypes.

phenotypes, and the workers had higher BMI. Unpressured and balanced wards were not extreme in any respect. Descriptive characteristics of the eldercare workers' musculoskeletal pain and sickness absence at baseline and during the one-year follow-up, stratified by ward phenotype, are shown in table 2.

Associations between ward phenotypes, musculoskeletal pain and sickness absence

During the one-year follow-up, turbulent wards were significantly associated with more days per month with NSP ($\beta=0.54$; $P<0.01$) and LBP ($\beta=0.37$; $P<0.01$) among the workers than balanced wards (table 3). Similar associations were found after adjustments for age, BMI and smoking habits, although only statistically significant for LBP ($\beta=0.35$; $P=0.01$). Thus, the adjusted model predicted that the number of days with LBP per month at follow-up would be 0.35 days larger at turbulent versus balanced wards.

Strained wards were associated with higher intensity of NSP pain ($\beta=0.14$; $P=0.03$), more days with LBP ($\beta=0.21$; $P=0.02$), and higher intensity of LBP ($\beta=0.15$; $P=0.04$) and were more likely to report pain-related work interference [odds ratio (OR) 1.67; $P=0.01$] compared with balanced wards. Strained wards were also associated with more days with musculoskeletal sickness absence ($\beta=0.83$; $P=0.07$) than balanced wards. After adjustments these results remained significant and with slightly higher effect estimates (table 3). Unpressured wards did not differ significantly from balanced wards in pain and sickness absence.

The sensitivity analysis of workers with low pain and no sickness absence at baseline showed, in general, stronger and more significant associations at follow-up in this sub-population than in the total population (supplementary table S2).

Discussion

Summary of findings

To our knowledge, our study is the first to collect onsite observations of resident handlings in eldercare, interpret those in terms of ward phenotypes, and address the prospective association of these phenotypes with musculoskeletal pain and sickness absence among eldercare workers. We identified four distinct ward phenotypes, labeled as turbulent, strained, unpressured, and balanced, representing distinctly different combinations of various handling characteristics. After adjusting for age, BMI and smoking habits, we found that workers at turbulent and strained wards reported a larger increase

Table 2. Musculoskeletal pain and sickness absence at baseline and over the one-year follow-up for the four ward phenotypes. Each phenotype was identified using a latent profile analysis based on observed handling characteristics at ward level. Values are presented as mean (standard deviation) between workers within a specific phenotype. [SD=standard deviation; LBP=low-back pain; NSP=neck-shoulder pain].

	Turbulent wards Mean (SD)	Strained wards Mean (SD)	Unpressured wards Mean (SD)	Balanced wards Mean (SD)
Days w/ NSP/month (0–28 days)				
Baseline	4.9 (8.3)	9.1 (9.8)	8.4 (10.1)	7.8 (9.8)
During follow-up	8.2 (8.0)	8.9 (7.8)	7.3 (7.6)	6.7 (6.6)
Intensity NSP/month (0–10 scale)				
Baseline values	3.4 (3.4)	4.4 (4.4)	4.1 (3.2)	3.8 (3.1)
During follow-up	3.5 (2.1)	4.2 (2.4)	3.5 (2.5)	3.2 (2.5)
Days w/ LBP/month (0–28 days)				
Baseline values	5.3 (9.2)	8.0 (8.5)	7.9 (9.5)	7.8 (9.7)
During follow-up	8.6 (8.5)	7.9 (7.0)	7.0 (7.3)	5.9 (6.5)
Intensity LBP/month (0–10 scale)				
Baseline values	3.5 (3.4)	4.3 (3.2)	4.0 (3.2)	3.8 (3.2)
During follow-up	3.8 (2.4)	4.1 (2.5)	3.4 (2.4)	3.1 (2.5)
Days w/ pain-related work interference (0–28 days)				
Baseline values	2.9 (6.3)	3.5 (5.8)	4.1 (7.8)	3.8 (7.4)
During follow-up	6.7 (6.3)	5.7 (6.3)	4.8 (6.3)	3.9 (4.9)
Days w/ musculoskeletal sickness absence (0–84 days)				
Baseline values	0.2 (0.7)	0.6 (2.1)	1.1 (6.4)	1.1 (5.5)
During follow-up	3.4 (9.3)	2.1 (5.7)	1.8 (5.8)	0.9 (3.0)

in pain during the one-year follow up than workers at balanced wards.

Identified ward phenotypes

Among the four ward phenotypes, two (turbulent and strained) appeared to have a less favorable combination of handling characteristics, while the two others (unpressured and balanced) had more beneficial profiles. To our knowledge, this is the first study to evaluate different aspects of resident handlings concomitantly.

At turbulent wards, workers had high physical demands reflected by a high frequency of handlings, but they also had resources to perform these handlings, as shown by the high proportion of handlings done with assistive devices and assistance from others. Handling assistance has been shown to reduce physical load during handling tasks (15, 31). On the other hand, handlings at turbulent wards also suffered many barriers, which has been associated with poor health (21, 25). Turbulent wards had a low staff ratio compared to other ward phenotypes (table 1), and the high number of interruptions may be explained by workers being disturbed more often because their colleagues needed assistance.

The strained wards also appear to have less favorable handlings characteristics; handlings were both frequent and, to a considerable extent, performed without assistive devices or assistance from others, likely have increasing the physical load on workers (15, 16, 31). Unpressured and balanced wards showed more favor-

Table 3. Associations between ward phenotypes, musculoskeletal pain and sickness absence over the one-year follow-up (14 time points for pain and 5 for sickness absence). Each phenotype (turbulent, strained, unpressured, and balanced) was identified using a latent profile analysis based on observed handling characteristics at ward level. Balanced wards was used as reference in all analyses. **Bold indicates a statistically significant difference.** [CI=confidence interval].

Ward phenotypes	Model 1 ^a			Model 2 ^b		
	Coefficient	95% CI	P-value	Coefficient	95% CI	P-value
Days with NSP per month (0–28 days) ^c						
Turbulent	0.54	0.26–0.82	<0.01	0.46	-0.21–1.13	0.18
Strained	0.17	-0.05–0.38	0.12	0.17	-0.30–0.64	0.48
Unpressured	0.04	-0.13–0.22	0.65	0.14	-0.24–0.52	0.47
Balanced	0.00			0.00		
Intensity NSP per month (0–10 scale) ^c						
Turbulent	0.12	-0.07–0.30	0.21	0.15	-0.04–0.34	0.13
Strained	0.14	0.01–0.28	0.03	0.18	0.04–0.31	0.01
Unpressured	0.04	-0.07–0.16	0.46	0.06	-0.06–0.19	0.32
Balanced	0.00			0.00		
Days with LBP per month (0–28 days) ^c						
Turbulent	0.37	0.16–0.58	<0.01	0.35	0.08–0.63	0.01
Strained	0.21	0.04–0.39	0.02	0.22	0.03–0.42	0.03
Unpressured	0.08	-0.07–0.22	0.29	0.11	-0.05–0.26	0.17
Balanced	0.00			0.00		
Intensity LBP per month (0–10 scale) ^c						
Turbulent	0.17	-0.05–0.40	0.13	0.11	-0.13–0.35	0.35
Strained	0.15	0.01–0.29	0.04	0.17	0.02–0.31	0.03
Unpressured	0.03	-0.08–0.13	0.62	0.05	-0.06–0.16	0.41
Balanced	0.00			0.00		
Days musculoskeletal sickness absence (0–84 days) ^c						
Turbulent	0.86	-0.28–2.01	0.14	0.90	-0.21–2.02	0.11
Strained	0.83	-0.05–1.71	0.07	0.67	-0.25–1.60	0.15
Unpressured	0.40	-0.30–1.11	0.26	0.34	-0.38–1.06	0.35
Balanced	0.00			0.00		
Days with pain-related work interference – categorical (% with ≥1 days/month) ^d						
Turbulent	1.53	0.83–2.83	0.17	1.33	0.67–2.63	0.40
Strained	1.67	1.14–2.44	0.01	1.79	1.20–2.66	<0.01
Unpressured	1.16	0.88–1.54	0.29	1.20	0.89–1.61	0.22
Balanced	1.00			1.00		

^a Adjusted for the baseline values of the outcomes.

^b Further adjusted by age, body mass index, smoking habits.

^c Model coefficient value expressed as β .

^d Model coefficient value expressed as odds ratio.

able handling characteristics, even though they were not ideal. Based on results from previous studies, ‘ideal’ wards would have <10 handlings per shift (11), most of these performed with assistive devices or assistance from others (15, 16, 31), and only few barriers (21, 25). Our current sample of wards did not include any such ideal ward phenotype. While interesting, a further examination of the determinants of ward-level handling characteristics, and hence of the handling phenotypes, was beyond the aims of the present study.

Association of ward phenotypes with musculoskeletal pain and sickness absence

Compared with balanced wards, working in a turbulent ward was associated with more frequent NSP and LBP, while working in a strained ward was associated with more days of LBP, higher intensity of NSP and LBP and a higher likelihood of pain interfering with work. This suggests that the risk of musculoskeletal pain increases with the frequency of handlings, particularly when handlings are more often performed without the use

of assistive devices or assistance from others. This is consistent with previous studies reporting that a higher frequency of handlings is associated with persistent LBP in eldercare workers (11) and a higher risk of back injuries among hospital healthcare workers (32). Notably, these previous studies, like several others (12–14, 31, 33), considered just one aspect of handlings at the time and were focused on individual-based exposure assessment. Previous studies suggest that pain intensity and pain-related work interference have clear associations with sickness absence among healthcare workers (34). But we did not find a statistically significant association between sickness absence and the ward phenotypes, even though we observed a tendency for more absence days due to musculoskeletal pain among workers at strained wards.

Performing resident handlings without assistive devices, as observed to a larger extent in strained wards, have been associated with LBP among nurses (35), but evidence supporting the efficacy of using assistive devices in reducing musculoskeletal pain among healthcare workers is still limited (36). A possible explanation is that

studies often consider the use of assistive devices as an isolated factor and ignore other important organizational factors that may act in parallel and correlate with the use of devices. By including job barriers (interruptions and impediments), defined as poor organizational, psychosocial and/or physical work conditions impeding the performance of work tasks (20, 37, 38), we addressed factors beyond the mere physical aspects of handlings. Thus, our ward phenotypes reflect a set of important external exposures associated with resident handlings, and our study considers the eventual results of these exposures in terms of individual outcomes, ie, pain and sickness absence, as illustrated in the exposure-effect model (6). A previous study showed that interruptions and impediments assessed using onsite observations were associated with musculoskeletal disorders in urban transit operators (25), but to our knowledge no previous study has examined this association among eldercare workers.

We found a large variability in reported pain and sickness absence at baseline, and so we also performed a sensitivity analysis including only those workers with little pain and no sickness absence. Ward phenotype predicted musculoskeletal pain and sickness absence even stronger in this sub-population, suggesting that the handling characteristics at turbulent and strained wards could, indeed, increase the risk of workers developing musculoskeletal disorders.

Methodological considerations, implications and future research

The use of a systematic and reliable onsite observation tool for assessing resident handlings characteristics (20, 24) is a strength of this study as it provided information that will likely be more accurate than that obtained, eg, by self-reports from the workers. Also, this approach eliminates the risk of common method bias, which is a likely fallacy in studies where both exposures and outcomes are based on self-report (17). In aggregating observations at the ward level, thus obtaining 'non-individualized' information about general handling characteristics at a specific ward, we further reduced the risk of common method bias. Another strength of the study is the novel use of latent profile analysis, which allowed us to identify clusters of wards sharing similar combinations of handling characteristics, which we could then interpret and label in terms of ward phenotypes. The use of latent profile analysis also allowed us to integrate physical and psychosocial aspects of handlings, and thus to examine their combined impact on musculoskeletal health. We believe that 'clustered' exposure assessment approaches like this offer an appealing alternative to standard multivariate regression, where exposure variables are entered in parallel, and correlations may severely challenge the interpretation of results. The

extensive assessment of pain and sickness absence, using 14 and 5 repeated measurements, respectively, may also be considered a strength of the study.

Still, the study also suffers some limitations. Even though the use of an observational tool can be considered an advantage compared to self-reports, it should ideally be combined with direct measurements of exposures (39), which we did not. Observations at a particular ward were performed during a short period of time, and the estimated handling variables at the ward level are subject to some inaccuracy due to variability between days, both on the short term, for instance because residents have different needs on different days, and on the longer term, because residents may be replaced by other residents with different needs, or staffing may change. Also, handlings for, on average, about 15% of the residents at a ward were obtained by imputation, which may have introduced additional variance. However, based on our experiences in eldercare, we believe these sources of exposure variability to be moderate, and thus having only a minor impact on the accuracy of the average ward-level exposure. Notably, our exposure assessment strategy was not sensitive to the proportion of participating workers, since work schedules were obtained for everybody, directly from the ward manager (24). Ward phenotypes were determined in a data-driven process, and even though our phenotypes appear representative of eldercare conditions, it is possible that the ward classification could be different in other populations. Future studies are needed to verify the stability of the phenotyping across wards in general.

The assessment of pain and sickness absence during the one-year follow-up relied solely on self-reports, which could introduce bias. However, we trust this risk to be reduced by the frequent assessment of outcomes, reducing recall bias, and we emphasize that common method variance of exposures and outcomes is likely small, due to the use of observational-based, aggregated exposures. Finally, while we believe each of the six selected outcomes to be important in reflecting different aspects of pain (severity, duration) and its consequences (interference with work, sickness absence due to musculoskeletal pain), we acknowledge the risk of obtaining significant results by chance in multiple tests.

Our results, in illustrating the importance of ward-level characteristics to health can be used in future studies as a basis for designing ward-level tools for preventing musculoskeletal symptoms in eldercare workers, as an alternative to, or complementing, individualized interventions, such as having workers attend courses in lifting techniques. While we believe that intervening on organizational and structural determinants of handlings would lead to more effective and sustainable results than interventions directed towards the behaviors of individual workers, future studies should verify the effectiveness of ward-level organizational tools, for

example in randomized control trials, before they are generally recommended for practice (40).

Concluding remarks

Based on onsite observations of resident handlings in eldercare, we were able to identify four handling phenotypes at the ward level. Our results show that turbulent and strained ward phenotypes were associated with some indicators of musculoskeletal pain during the one-year follow-up. Thus, our study suggests that ward phenotypes with frequent handlings, either combined with limited use of assistive devices or with many barriers, may impose an increased risk of musculoskeletal pain among the workers. Our findings encourage future studies devoted to the development of ward-level, organizational interventions in eldercare homes.

Acknowledgements

AFA Insurance (grant number 180076) funded this work at the University of Gävle, and the Danish Working Environment Fund (22-2019-09) funded the work performed by researchers from the National Research Centre for the Working Environment.

The authors declare no conflict of interest.

References

- Andersen LL, Clausen T, Persson R, Holtermann A. Dose-response relation between perceived physical exertion during healthcare work and risk of long-term sickness absence. *Scand J Work Environ Health* 2012 Nov;38(6):582–9. <https://doi.org/10.5271/sjweh.3310>.
- Davis KG, Kotowski SE. Prevalence of Musculoskeletal Disorders for Nurses in Hospitals, Long-Term Care Facilities, and Home Health Care: A Comprehensive Review. *Hum Factors* 2015 Aug;57(5):754–92. <https://doi.org/10.1177/0018720815581933>.
- Lund T, Labriola M, Villadsen E. Who is at risk for long-term sickness absence? A prospective cohort study of Danish employees. *Work* 2007;28(3):225–30.
- Hasselhorn HM, Conway PM, Widerszal-Bazyl M, Simon M, Tackenberg P, Schmidt S et al. Contribution of job strain to nurses' consideration of leaving the profession—results from the longitudinal European nurses' early exit study. *SJWEH Suppl.* 2008;34(6):75–82.
- Genet N, Boerma WG, Kroneman M, Hutchinson A, Saltman RB. Home Care across Europe - Current structure and future challenges. 1 st editi. World Heal. Organ. European Observatory on Health Systems and Policies; 2012.
- Winkel J, Mathiassen SE. Assessment of physical work load in epidemiologic studies: concepts, issues and operational considerations. *Ergonomics* 1994 Jun;37(6):979–88. <https://doi.org/10.1080/00140139408963711>.
- Westgaard RH, Winkel J. Guidelines for occupational musculoskeletal load as a basis for intervention: a critical review. *Appl Ergon* 1996 Apr;27(2):79–88. [https://doi.org/10.1016/0003-6870\(95\)00062-3](https://doi.org/10.1016/0003-6870(95)00062-3).
- Yassi A, Lockhart K. Work-relatedness of low back pain in nursing personnel: a systematic review. *Int J Occup Environ Health* 2013 Jul-Sep;19(3):223–44. <https://doi.org/10.1179/2049396713Y.0000000027>.
- da Costa BR, Vieira ER. Risk factors for work-related musculoskeletal disorders: A systematic review of recent longitudinal studies. *Am J Ind Med* 2010 Mar;53(3):285–323.
- Elford W, Straker L, Strauss G. Patient handling with and without slings: an analysis of the risk of injury to the lumbar spine. *Appl Ergon* 2000 Apr;31(2):185–200. [https://doi.org/10.1016/S0003-6870\(99\)00026-5](https://doi.org/10.1016/S0003-6870(99)00026-5).
- Holtermann A, Clausen T, Jørgensen MB, Burdorf A, Andersen LL. Patient handling and risk for developing persistent low-back pain among female healthcare workers. *Scand J Work Environ Health* 2013 Mar;39(2):164–9. <https://doi.org/10.5271/sjweh.3329>.
- Noble NL, Sweeney NL. Barriers to the Use of Assistive Devices in Patient Handling. *Workplace Health Saf* 2018 Jan;66(1):41–8. <https://doi.org/10.1177/2165079917697216>.
- Koppelaar E, Knibbe JJ, Miedema HS, Burdorf A. Determinants of implementation of primary preventive interventions on patient handling in healthcare: a systematic review. *Occup Environ Med* 2009 Jun;66(6):353–60. <https://doi.org/10.1136/oem.2008.042481>.
- Koppelaar E, Knibbe JJ, Miedema HS, Burdorf A. Individual and organisational determinants of use of ergonomic devices in healthcare. *Occup Environ Med* 2011 Sep;68(9):659–65. <https://doi.org/10.1136/oem.2010.055939>.
- Vinstrup J, Jakobsen MD, Madeleine P, Andersen LL. Physical exposure during patient transfer and risk of back injury & low-back pain: prospective cohort study. *BMC Musculoskelet Disord* 2020 Oct;21(1):715. <https://doi.org/10.1186/s12891-020-03731-2>.
- Koppelaar E, Knibbe HJ, Miedema HS, Burdorf A. The influence of ergonomic devices on mechanical load during patient handling activities in nursing homes. *Ann Occup Hyg* 2012 Jul;56(6):708–18.
- Podsakoff PM, MacKenzie SB, Lee JY, Podsakoff NP. Common method biases in behavioral research: a critical review of the literature and recommended remedies. *J Appl Psychol* 2003 Oct;88(5):879–903. <https://doi.org/10.1037/0021-9010.88.5.879>.
- Jansen JP, Burdorf A. Effects of measurement strategy and statistical analysis on dose-response relations between physical workload and low back pain. *Occup Environ Med* 2003 Dec;60(12):942–7. <https://doi.org/10.1136>

[oem.60.12.942](https://doi.org/10.1016/j.ajep.2018.01.000).

19. Jansen JP, Morgenstern H, Burdorf A. Dose-response relations between occupational exposures to physical and psychosocial factors and the risk of low back pain. *Occup Environ Med* 2004 Dec;61(12):972–9. <https://doi.org/10.1136/oem.2003.012245>.
20. Karstad K, Rugulies R, Skotte J, Munch PK, Greiner BA, Burdorf A, et al. Inter-rater reliability of direct observations of the physical and psychosocial working conditions in eldercare: An evaluation in the DOSES project. *Appl Ergon*. 2018 May;69:93–103. <https://doi.org/10.1016/j.apergo.2018.01.000>.
21. Jakobsen LM, Jorgensen AF, Thomsen BL, Greiner BA, Rugulies R. A multilevel study on the association of observer-assessed working conditions with depressive symptoms among female eldercare workers from 56 work units in 10 care homes in Denmark. *BMJ Open* 2015 Nov;5(11):e008713–008713. <https://doi.org/10.1136/bmjopen-2015-008713>.
22. Koppelaar E, Knibbe JJ, Miedema HS, Burdorf A. The influence of individual and organisational factors on nurses' behaviour to use lifting devices in healthcare. *Appl Ergon* 2013 Jul;44(4):532–7. <https://doi.org/10.1016/j.apergo.2012.11.005>.
23. Austin CP. Talking Glossary of Genetic Terms [Internet]. Natl. Hum. Genome Res. Inst. [cited 2021 May 27]. Available from: <https://www.genome.gov/genetics-glossary/Phenotype>.
24. Karstad K, Jørgensen AF, Greiner BA, Burdorf A, Søgaard K, Rugulies R et al. Danish Observational Study of Eldercare work and musculoskeletal disorderS (DOSES): a prospective study at 20 nursing homes in Denmark. *BMJ Open* 2018 Feb;8(2):e019670. <https://doi.org/10.1136/bmjopen-2017-019670>.
25. Greiner BA, Krause N. Observational stress factors and musculoskeletal disorders in urban transit operators. *J Occup Health Psychol* 2006 Jan;11(1):38–51. <https://doi.org/10.1037/1076-8998.11.1.38>.
26. Johansen B, Wedderkopp N. Comparison between data obtained through real-time data capture by SMS and a retrospective telephone interview. *Chiropr Osteopat* 2010 May;18(1):10. <https://doi.org/10.1186/1746-1340-18-10>.
27. Oberski D. Mixture Models: Latent Profile and Latent Class Analysis. In: Robertson J, Kaptein M, editors. Cham: Springer International Publishing; 2016. p. 275–87.
28. Vermunt JK. Multilevel Latent Class Models. *Sociol Methodol* 2003 Aug;33(1):213–39. <https://doi.org/10.1111/j.0081-1750.2003.t01-1-00131.x>.
29. Mäkitangas A, Tolvanen A, Aunola K, Feldt T, Mauno S, Kinnunen U. Multilevel Latent Profile Analysis With Covariates. *Organ Res Methods* 2018 Oct;21(4):931–54. <https://doi.org/10.1177/1094428118760690>.
30. Hubbard AE, Ahern J, Fleischer NL, Van der Laan M, Lippman SA, Jewell N et al. To GEE or not to GEE: comparing population average and mixed models for estimating the associations between neighborhood risk factors and health. *Epidemiology* 2010 Jul;21(4):467–74. <https://doi.org/10.1097/EDE.0b013e3181caeb90>.
31. Vinstrup J, Jakobsen MD, Madeleine P, Andersen LL. Biomechanical load during patient transfer with assistive devices: cross-sectional study. *Ergonomics* 2020 Sep;63(9):1164–74. <https://doi.org/10.1080/00140139.2020.1764113>.
32. Andersen LL, Vinstrup J, Villadsen E, Jay K, Jakobsen MD. Physical and Psychosocial Work Environmental Risk Factors for Back Injury among Healthcare Workers: Prospective Cohort Study. *Int J Environ Res Public Health* 2019 Nov;16(22):E4528. <https://doi.org/10.3390/ijerph16224528>.
33. Choi SD, Brings K. Work-related musculoskeletal risks associated with nurses and nursing assistants handling overweight and obese patients: A literature review. *Work* 2015;53(2):439–48. <https://doi.org/10.3233/WOR-152222>.
34. Murray E, Franche RL, Ibrahim S, Smith P, Carnide N, Côté P et al. Pain-related work interference is a key factor in a worker/workplace model of work absence duration due to musculoskeletal conditions in Canadian nurses. *J Occup Rehabil* 2013 Dec;23(4):585–96. <https://doi.org/10.1007/s10926-012-9408-7>.
35. Smedley J, Egger P, Cooper C, Coggon D. Manual handling activities and risk of low back pain in nurses. *Occup Environ Med* 1995 Mar;52(3):160–3. <https://doi.org/10.1136/oem.52.3.160>.
36. Hegewald J, Berge W, Heinrich P, Staudte R, Freiberg A, Scharfe J et al. Do Technical Aids for Patient Handling Prevent Musculoskeletal Complaints in Health Care Workers?—A Systematic Review of Intervention Studies. *Int J Environ Res Public Health* 2018 Mar;15(3):476. <https://doi.org/10.3390/ijerph15030476>.
37. Landsbergis P, Theorell T, Schwartz J, Greiner BA, Krause N. Measurement of psychosocial workplace exposure variables. *Occup Med* 2000 Jan-Mar;15(1):163–88.
38. Greiner BA, Krause N, Ragland DR, Fisher JM. Objective stress factors, accidents, and absenteeism in transit operators: a theoretical framework and empirical evidence. *J Occup Health Psychol* 1998 Apr;3(2):130–46. <https://doi.org/10.1037/1076-8998.3.2.130>.
39. van der Beek AJ, Frings-Dresen MH. Assessment of mechanical exposure in ergonomic epidemiology. *Occup Environ Med* 1998 May;55(5):291–9. <https://doi.org/10.1136/oem.55.5.291>.
40. van der Beek AJ, Dennerlein JT, Huysmans MA, Mathiassen SE, Burdorf A, van Mechelen W et al. A research framework for the development and implementation of interventions preventing work-related musculoskeletal disorders. *Scand J Work Environ Health* 2017 Nov;43(6):526–39. <https://doi.org/10.5271/sjweh.3671>.

Received for publication: 3 March 2021